


Fisheries and Environment
Canada

ENVIRONMENTAL
PROTECTION

AN ENVIRONMENTAL ASSESSMENT OF THE
EFFECTS OF CASSIAR ASBESTOS CORPORATION
ON CLINTON CREEK, YUKON TERRITORIES



Surveillance Report

EPS 5-PR-77-3

Pacific Region
June 1977

00 551 04
FILE
CASSIAR ASBES

AN ENVIRONMENTAL ASSESSMENT OF THE
EFFECTS OF CASSIAR ASBESTOS CORPORATION
ON CLINTON CREEK, YUKON TERRITORIES

Pollution Abatement Branch
Environmental Protection Service
Pacific Region

Report Number EPS 5-PR-77-3

June 1977

TABLE OF CONTENTS

PAGE

ABSTRACT

RÉSUMÉ

TABLE OF CONTENTS

LIST OF FIGURES

LIST OF TABLES

1	INTRODUCTION	1
2	CASSIAR ASBESTOS CORPORATION LIMITED - CLINTON CREEK MINE	1
3	DESCRIPTION OF STUDY AREA	4
4	METHODS AND MATERIALS	5
4.1	Sampling Stations	5
4.2	Water Quality Sampling	5
4.2.1	Water Chemistry	9
4.2.2	Asbestos Fibre Analysis	9
4.2.3	Toxicity	10
4.3	Biological Sampling	10
4.3.1	Invertebrates	10
4.3.2	Fish	11
4.3.3	Histological Assessment	12
5	RESULTS AND DISCUSSION	13
5.1	Water Chemistry	13
5.2	Asbestos Fibre Analysis	19
5.3	Toxicity	21
5.4	Invertebrates	21
5.5	Fish	24
5.6	Histological Assessment	27

	PAGE
6 CONCLUSIONS AND RECOMMENDATIONS	35
REFERENCES	37

LIST OF FIGURES

FIGURE		PAGE
1	LOCATION OF OPEN PIT, MILL AND MINE	3
2	STATION LOCATIONS	6
3	HUDGEON LAKE	8
4	WOLVERINE CREEK	8
5	WATER CHEMISTRY RESULTS - NON-METALS COLOUR, TURBIDITY, NON-FILTERABLE RESIDUE, TOTAL HARDNESS	16
6	WATER CHEMISTRY RESULTS - NON-METALS DISSOLVED OXYGEN, TOTAL ALKALINITY, CONDUCTIVITY	17
7	WATER CHEMISTRY RESULTS - METALS IRON, MAGNESIUM, CALCIUM	18
8	SEPARATION OF LAMELLAE EPITHELIUM	31
9	HYPERPLASM	32
10	ANEURYSM	32

LIST OF TABLES

TABLE		PAGE
1	CASSIAR ASBESTOS CORPORATION CLINTON CREEK MINE OPERATING SUMMARY 1969 - 1975	2
2	CASSIAR ASBESTOS CORPORATION LIMITED CLINTON CREEK MINE 1974 & 1975 SAMPLING STATIONS	7
3	WATER CHEMISTRY RESULTS - NON-METAL	14
4	WATER CHEMISTRY RESULTS - TOTAL METALS	15
5	ASBESTOS FIBRES IN WATER SAMPLES FROM HUDGEON LAKE AND FORTY MILE RIVER	20

TABLE		PAGE
6	TOXICITY EVALUATION - 96 Lr LC ₅₀ BIOASSAY RESULTS	22
7	MACRO INVERTEBRATE POPULATION AUGUST 1975 - CLINTON CREEK MINE	23
8	INVERTEBRATE SPECIES LIST AUGUST 1975	25
9	DIVERSITY AND EVENNESS RESULTS INVERTEBRATE POPULATION AUGUST 1975; CLINTON CREEK MINE	28
10	ELECTRO FISHING RESULTS AUGUST 6 - 7 1975	29
11	HISTOLOGICAL ASSESSMENT SLIDES ANALYSED	33
12	HISTOLOGICAL ASSESSMENT OF GILL TISSUE (ONCORHYNCHUS KISUTCH)	34

1 INTRODUCTION

There are only two asbestos mines north of the 60th parallel in Canada, one is the Asbestos Hill Mine in Putunig, Quebec, and the other is the Clinton Creek Mine in the Yukon Territory. The Clinton Creek Mine is located 97 kilometers northwest of Dawson City. The mine is owned and operated by Cassiar Asbestos Corporation Limited. It began producing in October, 1967 and is expected to cease operation in the summer of 1978.

In response to discussions with the Department of Indian and Northern Development (DIAND) and concerns expressed by citizens of Clinton Creek, Environmental Protection Service (Yukon District) conducted an aquatic environmental quality survey. A chemical and biological sampling program was initiated in the summer of 1974 and was repeated in 1975 to expand the data base. This report presents the results of these surveys. It should provide a baseline to which future environmental effects and possibly post-operational stabilization activities may be related.

2 CASSIAR ASBESTOS CORPORATION LIMITED - CLINTON CREEK MINE

The Cassiar Asbestos Corporation Limited, Clinton Creek Mine is an open pit asbestos mine located 64° 24.5' north latitude and 140° 36' west longitude. In 1975 it was the second highest mineral producer in the Yukon Territory.

The Clinton Creek deposit is an ultramafic body that occurs in metamorphic rocks that consist mainly of black phyllite, platy black limestone, grey argillite and brown weathering micaceous gritty quartzite (Sinclair, Gilbert, 1975). The asbestos occurs almost entirely as cross-fibre veins one quarter inch wide or less (Sinclair et al, 1976).


The production rate in 1968 was approximately 2500 tons per day (tpd) and in 1974 and 1975 it increased to 4596

TABLE 1

CASSIAR ASBESTOS CORPORATION - CLINTON CREEK MINE - OPERATING SUMMARY 1969-1975 (1)

	1969	1970	1971	1972	1973	1974	1975
Tons Milled	952,889	1,335,087	1,447,863	1,267,178	1,247,154	1,388,248	1,407,453
Rate (tons/day)	3,048	4,049	4,460	4,400	4,838	4,596	5,118
Grade (% recovery)	9.22	7.82	5.37	5.66	5.64	4.37	5.85
Reserves (Probable)			18,750,000	9,250,000	7,861,123	6,524,725	4,773,000
(Possible)				9,500,000	8,792,000	461,000	928,931

(1) (Craig, Laporte 1972; Sinclair, Gilbert 1 1975; Sinclair et al 1976)



tpd and 5118 tpd respectively (Table 1). In 1975, 1407453 tons of ore were milled with a recovery of 5.85 percent. At the close of 1975 the probable reserves were calculated at 4,773,000 tons. It is expected that the mine will cease operation in 1978 when the ore reserves will have been depleted.

By the end of 1975, after seven years of operation, an estimated 75,000,000 tons of waste have been removed from the mine.

3 DESCRIPTION OF STUDY AREA

Clinton Creek is a left bank tributary to the Forty-Mile River, 4.8 kilometers southwest of its confluence with the Yukon River. The town of Clinton Creek is located 86 kilometers northwest of Dawson City, Yukon Territories and is connected to the Yukon Territorial Highway System by a 30 kilometer access road. The open pit, mill and mine are approximately 11 kilometers northwest of the townsite (Figure 1).

The mine is located in hilly, unglaciated terrain. There are three open pits; Porcupine, the largest; Snowshoe and Creek Pit. The Porcupine and Snowshoe Pits are located on hills on the south side of Clinton Creek. Creek Pit is on the original alignment of Porcupine Creek.

The waste rock overburden from the pits, consisting mainly of argillite has been dumped in two main areas: Clinton Dump and Porcupine Dump. Clinton Dump is located south of Clinton Creek and contains several tens of millions of tons of waste rock. Porcupine Dump is much smaller and contains less than five million tons of waste. (Golder-Associates, 1976). Both dumps extend across the original channels of Clinton and Porcupine Creeks respectively as a result of waste material gradually creeping downslope. In the spring of 1974 the Clinton Creek channel was blocked, resulting in the formation of a lake (Hudgeon Lake) upstream of the toe of the dump.

The tailings pile from the mill is located on the west side of Wolverine Creek. In the spring of 1974, a large section of the tailings pile moved downslope and blocked the creek forming Wolverine Lake upslope of the toe of the slide. The impoundment was breached in the summer of 1974 and reformed in the spring of 1975.

The study area centered on Clinton Creek as the drainage from the open pit, waste rock piles and tailings all enter this stream.

4 METHODS AND MATERIALS

4.1 Sampling Stations

Nine stations were established for water quality and biological sampling. The stations are illustrated in Figure 2 and described in Table (2). All stations were marked by flagging nearby trees.

With the exception of stations one and eight, at the time of sampling these stations were located in shallow water with an average depth .45 - .60 meters. The water depth at stations one and eight was approximately one meter.

Sampling sites which differ from those described above are outlined in the appropriate sections.

4.2 Water Quality Sampling

The water quality of Clinton Creek was assessed by collecting samples for chemical analyses, metal and non-metal parameters; asbestos fibre content and toxicity (96Lr LC50)




FIGURE 2 STATION LOCATIONS

TABLE 2 CASSIAR ASBESTOS CORPORATION LIMITED
CLINTON CREEK MINE - 1974 and 1975 SAMPLING STATIONS

- Station 1 - On Clinton Creek above Hudgeon Lake.
- Station 2 - On Clinton Creek below waste rock pile but above the confluence with Wolverine Creek.
- Station 3 - On Wolverine Creek above Wolverine Lake.
- Station 4 - On Wolverine Creek just above the confluence with Clinton Creek.
- Station 5 - On Clinton Creek approximately 300 meters below the confluence of Wolverine Creek and Clinton Creek.
- Station 6 - On Clinton Creek Approximately 5 kilometers below station 5.
- Station 7 - On Clinton Creek, 100 meters above road leading to the townsite.
- Station 8 - On the Forty-Mile River, 300 meters below the confluence of Clinton Creek and the Forty-Mile River.
- Station 9 - On the Forty-Mile River, 100 meters above the confluence of Clinton Creek and the Forty-Mile River.




FIGURE 3 Hudgeon Lake (Formed by shifting of the waste rock - see arrow A).
Arrow B shows approximate sampling location
- histological assessment.




FIGURE 4 Wolverine Creek (Trees laden with asbestos fibers.)

4.2.1 Water Chemistry. Samples were collected during August 6 - 9, 1974, June 26 and August 8, 1975 for chemical analysis. All water samples were analyzed in the field for temperature, conductivity, dissolved oxygen and pH. Temperature and conductivity were measured with a Yellow Springs Instrument Company Salinity-Conductivity-Temperature Meter. Dissolved oxygen content was measured by the Azide Modification of the Winkler Method (Standard Methods, 1971). pH was determined using a Radiometer, Model 29B.

Additional samples were collected at all three sampling periods and analyzed for the following: pH, non-filterable residue (N.F.R.), total alkalinity, colour, total hardness, turbidity, and the metals; calcium (Ca), magnesium (Mg), cadmium (Cd), copper (Cu), iron (Fe), lead (Pb), molybdenum (Mo), manganese (Mn), potassium (K), and zinc (Zn).

The samples which were collected in June, 1975 were lost in transit and not recovered. In 1975, the turbidity measurements were changed from Jackson Turbidity Units (J.T.U.) used in 1974 to Formazin Turbidity Units (F.T.U.). In 1975, the samples were also analyzed for nickel (Ni), silver (Ag), and mercury (Hg). Manganese and potassium were not determined in 1975.

All samples were preserved and analyzed according to the methods described by Davidson et al, 1974, at the Environmental Protection Service, Chemistry Laboratory in West Vancouver.

4.2.2 Asbestos Fibre Analysis. In November, 1974, DIAND collected six water samples for asbestos fibre analysis. Three samples were collected from Hudgeon Lake by the pumphouse for the mine water supply and three were collected from the Forty-Mile River by the pumphouse for the Clinton Creek town-site drinking water supply.

A qualitative and quantitative analysis was done on two samples from each area by Dr. R.W. Durham at the Canada Centre for Inland Waters in Burlington, Ontario.

4.2.3 Toxicity. Water samples were collected at stations two and four on August 12, 1974 for static toxicity evaluation. The samples were collected in five gallon plastic jerry cans that were steam and acid cleaned in the laboratory, rinsed with sample water then filled to capacity.

Static bioassays were conducted using Coho salmon (*Oncorhynchus kisutch*) fry at the Environmental Protection Service Bioassay Laboratory in West Vancouver according to Davidson et al; 1974.

4.3 Biological Sampling

A biological assesement of Clinton Creek was conducted by obtaining qualitative and quantitative samples of aquatic invertebrates and fish. In addition, a histological analyses was conducted on gill tissue of Juvenile Coho salmon (*Oncorhynchus kisutch*) which had been subjected to bioassays using water collected from Hudgeon Lake.

4.3.1 Invertebrates. Samples of aquatic invertebrates were obtained at all stations in August, 1974 and again in August, 1975.

In 1974, a one square foot (366 sq. cm.) circular sample was used to sample the stream beds. The sampler was turned into the stream bed; the enclosed substrate agitated and the invertebrates removed from the screen. Three samples from each station were collected.

In 1975 artificial substrate samplers (A.S.S.) were used to sample the invertebrate populations. The A.S.S. were chrome plated barbecue chicken baskets with an exterior diameter of 17 cm., and length of 25 cm. They were filled with rocks of a uniform size from the creek flood plain or creek bottom that had been wiped clean of debris and vegetation. The surface area contained in each basket was

6000 \pm 1000 sq. cm. Three samplers were submerged in the creek at each station and secured to the bank by a rope. The samplers were placed on station on June 26, 1975 and retrieved on August 8, 1975.

All samples were preserved in 70% methanol. The organisms were sorted and grouped into families at the Environmental Protection Service Laboratory in Whitehorse. In 1974, the final identification and enumeration was done by W. Robson, Technician, Environmental Protection Service, and in 1975 it was done by Dr. C. Low of Envirocon Limited.

The data obtained were analyzed statistically on a Hewlett-Packard Model 9830A computer for diversity (\bar{d}) as described by Wilhm and Dorris, 1968, and for evenness of distribution (J) as described by Pielou, 1966. The formulas used in the calculations of these pollution indices are as follows:

$$\bar{d} = \sum_{i=1}^s \frac{n_i}{n} \log_2 \frac{n_i}{n}$$

$$J = \frac{-\sum_{i=1}^s \frac{n_i}{n} \log_2 \frac{n_i}{n}}{\log_2 a}$$

n_i = total number of individuals per taxon

n = total number of individuals per sample

s = total number of taxa

a = total number of species sampled

4.3.2 Fish. Fish populations were sampled in August 6 - 9, 1974 and August 6 - 7, 1975. A Smith-Root Type VIII Electro-Fisher and barrier nets were used at all stations. The barrier nets were placed downstream of the Electro-Fisher and approximately 30 meters of creek were fished before the nets were checked.

4.3.3 Histological Assessment.

In March, 1975, two water samples were collected from Hudgeon Lake by the pumphouse for the mine water supply. These samples were collected by DIAND in two 45 gallon drums and forwarded to the Environmental Protection Service Laboratory in West Vancouver. Subsamples were forwarded to Dr. R.W. Durham, Canada Centre for Inland Waters, Burlington, Ontario; for asbestos fibre analysis according to R.W. Durham and T.Pang, 1975.

The remaining sample was submitted to the Environmental Protection Service Bioassay Laboratory for bioassay determinations according to Davidson et al, 1974.

Juvenile Coho salmon (*Oncorhynchus kisutch*) were used in all bioassays. Five fish were placed in each aquarium and the aquaria were aerated. Food was supplied to all fish and a mixing apparatus was used to keep the water circulating and the asbestos fibres in suspension. The following bioassays were conducted:

- (a) 4 days at 100% concentration
- (b) 8 days at 100% concentration
- (c) 16 days at 100% concentration
- (d) 16 days at 100% concentration, then
16 days recovery in clean water.

After the termination of each bioassay, both the test and control fish were sacrificed. Gills were immediately dissected out and fixed in a 10% neutrally buffered formalin solution, for histological examination.

The examination of the gills for histological examination was done by Mr. J. Hollingdale, a research assistant at Simon Fraser University, this work was completed in April, 1975.

The gills were washed in tap water for twelve hours, and then dehydrated by rinsing through a series of graded ethanol concentrations; cleaned in xylene and embedded

in tissue mat at 60°C. Sixty-one serial sections, six microns in thickness, were cut on a rotary microtome, placed on slides and stained with haematoxylin and eosin dyes.

Histological examination and assessment was done in December, 1976, by J. McBride, Research and Resource Services Directorate, Fisheries and Marine Service, West Vancouver, B.C. Each slide was evaluated for the percentage of tissue showing the following histological disorders:

- (a) Separation of the Lamellae
- (b) Aneurysm
- (c) Hyperplasia

5 RESULTS AND DISCUSSION

5.1 Water Chemistry

The water chemistry data are presented in Tables 3 and 4 and Figures 5, 6 and 7. (Stations 2, 4, 5 and 6 appear to be affected by both the tailings pile on Wolverine Creek and the waste rock pile on Clinton Creek.) Elevated values for Calcium, Magnesium, Iron, Manganese and Potassium are evident at Stations 2, 4, 5 and 6. The high magnesium levels are indicative of the brucite ($MgO \cdot H_2O$) and magnesium carbonate in the orebody.

Corresponding higher values were obtained for hardness, and turbidity at the same stations. The values obtained at Station 4 on Wolverine Creek are comparable to those at Stations 2, 5 and 6 on Clinton Creek, indicating that both the tailings pile and waste rock pile are having similar effects on the water quality. However, these values are not significantly high to be considered harmful to aquatic organisms.

Dissolved oxygen levels were within the expected range of 7 - 10 (ppm) with the exception of two lower values which were obtained in the 1974 measurements; 6.0 ppm at Station 2 and 4.3 ppm at Station 6.

All other chemical parameters measured did not appear to be affected by the mine.

TABLE 3 WATER CHEMISTRY RESULTS - NON-METAL

Sample Date	Temp °C	Cond µmho/cm	D.O. PPM	pH (Field)	pH (Lab)	NFR mg/l	T. Alk. mg/l (CaCO3)	Colour C.U.	T. Hardness (mg/l CaCO3)	Turb. (1)
Stn 1	7/8/74	145	9.5	7.4	7.7	<2.5	210	70	230	<0.5
	26/6/75	550	9.5	7.4						
	8/8/75	172	10.3	7.7		6.	93.5	97	150	0.8
Stn 2	7/8/74	455	6.0	7.8	7.7	25.	170	45	1200	75.
	26/6/75	580	8.0	7.8						
	8/8/75	130	8.2	7.6		36.	96.8	126	180	4.7
Stn 3	7/8/74	252	7.2	7.0	7.6	35.	150.	60	100	<0.5
	26/6/75	400	9.5	8.1						
	8/8/75	195	10.2	7.8		3.	86.9	125	160	0.7
Stn 4	7/8/74	300	8.3	9.5	9.7	15.	240.	30	1100	46
	26/6/75	500	6.5	9.2						
	8/8/75	315	9.6	8.8		61.	131	93	360	10
Stn 5	7/8/74	385	7.4	8.3	9.0	4.	220	35	1400	48
	26/6/75	590	8.4	8.9						
	8/8/75	230	8.1	7.3		28.	99	112	250	3.5
Stn 6	7/8/74	375	4.3	7.0	7.2	<2.5	320	25	1200	9.7
	26/6/75	700	7.5	7.2						
	8/8/75	380	8.0	7.8		29.	125	115	290	13.
Stn 7	7/8/74	280	9.3	7.5	7.9	2.5	210	60	380	1.2
	26/6/75	600	9.5	8.0						
	8/8/75	285	9.1	7.8		62.	93.5	122	290	12.
Stn 8	7/8/74	220		7.9	7.7	<2.5	110	55	140	0.7
	26/6/75	300	9.0	8.1						
	8/8/75	240	8.7	7.8		52.	129	131	190	20.
Stn 9	7/8/74	132		7.8	7.2	2.8	59	25	34	<0.5
	26/6/75	130	8.2	8.1						
	8/8/75	80	8.7	7.6		17.	40.7	109	48	2.2

(1) 1974 Jackson Turbidity Units (JTU); 1975 Formazin Turbidity Units (FTU)

TABLE 4 WATER CHEMISTRY RESULTS - TOTAL METALS

Stn	Sample Date	Ca (ppm)	Mg (ppm)	Cd (ppm)	Cu (ppm)	Fe (ppm)	Pb (ppm)	Mo (ppm)	Ni (ppm)	Mn (ppm)	K (ppm)	Ag (ppm)	Zn (ppm)	Hg ⁽¹⁾ (ppb)
Stn 1	7/8/74	30	37	<0.1	<0.1	0.15	<0.1	<.3		<0.03	0.40		<0.01	
	26/6/75 8/8/75	32	16	<0.1	<0.1	0.23	<0.2	<.3	<0.05			<0.03	<0.01	<0.15
Stn 2	7/8/74	110	220	<0.1	<0.1	1.30	<0.2	<.3		0.59	2.50		<0.01	
	26/6/75 8/8/75	37	21.5	<0.1	<0.1	1.30	<0.2	<.3	<0.05			<0.03	<0.01	<0.15
Stn 3	7/8/74	39	2.7	<0.1	<0.1	0.28	<0.1	<.3		<0.03	0.71		<0.01	
	26/6/75 8/8/75	28	22.5	<0.1	<0.1	0.25	<0.2	<.3	<0.05			<0.03	<0.01	<0.15
Stn 4	7/8/74	5	270	<0.1	<0.1	2.10	<0.1	<.3		0.14	1.90		<0.01	
	26/6/75 8/8/75	46	60.5	<0.1	<0.1	1.60	<0.2	<.3	0.16			<0.03	<0.01	<0.15
Stn 5	7/8/74	63	210	<0.1	<0.1	1.90	<0.1	<.3		0.39	2.20		<0.01	
	26/6/75 8/8/75	48	31.5	<0.1	<0.1	.90	<0.2	<.3	<0.05			<0.03	<0.01	<0.15
Stn 6	7/8/74	5.9	280	<0.1	<0.1	0.66	<0.1	<.3		0.29	2.20		<0.01	
	26/6/75 8/8/75	48	42	<0.1	<0.1	1.40	<0.2	<.3	0.07			<0.03	<0.01	<0.15
Stn 7	7/8/74	40	70	<0.1	<0.1	0.14	<0.1	<.3		0.17	1.30		<0.01	
	26/6/75 8/8/75	46	42.5	<0.1	<0.1	1.50	<0.2	<.3	0.09			<0.03	<0.01	<0.15
Stn 8	7/8/74	16	24	<0.1	<0.1	0.12	<0.1	<.3		0.12	1.30		<0.01	
	26/6/75 8/8/75	31	27	<0.1	<0.1	1.10	<0.2	<.3	0.06			<0.03	<0.01	<0.15
Stn 9	7/8/74	12	0.7	<0.1	<0.1	0.07	<0.1	<.3		0.08	0.97		<0.01	
	26/6/75 8/8/75	27	4.3	<0.1	<0.1	0.6	<0.2	<.3	<0.05			<0.03	<0.01	<0.15




FIGURE 5 WATER CHEMISTRY RESULTS - NON-METALS
COLOUR, TURBIDITY, NON-FILTERABLE RESIDUE,
TOTAL HARDNESS






FIGURE 6

WATER CHEMISTRY RESULTS - NON-METALS
DISSOLVED OXYGEN, TOTAL ALKALINITY,
CONDUCTIVITY






FIGURE 7

WATER CHEMISTRY RESULTS - METALS
IRON, MAGNESIUM, CALCIUM

5.2 Asbestos Fibre Analysis

Water samples from Hudgeon Lake and the Forty Mile River were analyzed for asbestos fibre concentration in November 1974. The results obtained are presented in Table (5).

Asbestos fibre concentrations are reported in the number of chrysotile fibres per litre of water. Chrysotile is a fibrous form of ~~S~~erpentine and is the most common form found in the aquatic environment. The background level of asbestos in Canada and the northern U.S. surface waters varies between 1×10^5 fibres per litre and 1×10^7 fibres per litre (Lawrence, J; Zimmerman, H. Walter¹⁹⁷⁴).

In March, 1975 in conjunction with the histological assesement study an additional water sample from Hudgeon Lake was analyzed for asbestos fibres.

The results obtained on the Forty Mile River near the pumphouse for the Clinton Creek water supply are typical of most surface waters measured elsewhere in Canada.

The fibre concentration in the samples from Hudgeon Lake are appreciably higher than those from the Forty Mile River. The results obtained in 1974 are within the range of values that have been reported for raw water supplies. The fibre content found in 1975 ($300 \pm 80 \times 10^6$ fibres per litre) at Hudgeon Lake is high in comparison to the 1974 samples and in comparison to other background levels reported (Kay, G.H., 1973).

The mean fibre length of the 1974 samples ranged from 1 μ m to 1.5 μ m. This is considerably higher than natural fibre length averages of 0.2 μ m. This probably reflects the proximity to an airborne asbestos source (Durham, R.W., 1976). Fibre length determinations were not obtained for the 1975 sample.

TABLE 5

ASBESTOS FIBRES IN WATER SAMPLES FROM HUDGEON LAKE
AND FORTY MILE RIVER

<u>Station</u>	<u>Date & Time of sampling</u>		<u>Concentration No. of fibres per litre</u>	<u>mean fibre size</u>
Hudgeon Lake by pumphouse for mine water supply	22/10/74	12.45	36.4×10^6	1 μm
Duplicate	23/10/74	10.30	46.6×10^6	1 μm
Forty mile River by pumphouse for Clinton Creek water supply	22/10/74	14.40	$\sim 10^5$ (Detection limit)	-
Duplicate	23/10/74	08.30	3.0×10^6	1.5 μm
Hudgeon Lake by pumphouse for mine water supply	03/75		$300 \pm 80 \times 10^6$	

5.3 Toxicity

The bioassay results for water samples collected Stations 2 and 4 are presented in Table 6. The samples were found to be non-toxic at 100% concentration over 96 hours. These results confirm the water chemistry statistics.

Additional bioassays were conducted as part of the histological assessment study done in 1975. These bioassays were done on the Hudgeon Lake water sample and were also non-toxic at 100% concentration for over 96 hours. Subsequent bioassays were carried out for 8 and 16 days and were also non-toxic.

5.4 Invertebrates

The use of the modified circular samples for aquatic invertebrate collection in 1974 was not adaptable to the substrates encountered. Few organisms were collected and consequently the results were disregarded. The results of the sampling program in 1975 using artificial substrate samplers is considered representative of the invertebrate communities in the study area. The population summary and species list are presented in Tables 7 and 8.

A total of 32 species representing 2 plyla were found. Insects dominated the populations in terms of total numbers at all stations, with the exception of Station 5. The community structure at the control Stations 1 and 3 was similar. The cladoceran Moina Macropa was the dominant organism at Station 5.

The insect most prevalent was the midge corynoneura.

The absence of invertebrates at Station 4 is likely related to the substrate. The water from Wolverine Lake flows past the toe of the tailings site and is dispersed into the tailings which have been deposited in the creek bed. The water surfaces again approximately 100 meters downstream and flows through braided channels until the confluence with

TABLE 6

TOXICITY EVALUATION - 96 HR LC₅₀ BIOASSAY RESULTS

TEST FISH - Juvenile Coho Salmon (*Oncorhynchus kisutch*) acclimated to untreated Capilano Reservoir Tap Water at 11°C ± 1°C.
 Mean Fish Weight 4.91 gm
 Mean Fish Length 7.5 cm
 Test Tank Volume 20 kilo

Station	Station Location	Date Sampled	Date Bioassay	Initial Sample pH	Final Sample pH	Temp Range °C	Loading Density gm/l	% Survival 96 Hr	96 Hr LC ₅₀
2	Clinton Creek	Aug. 12/74	Aug. 16/74	8.93	8.21	12±1	1.22	100	Not Established
4	Wolverine Creek	Aug. 12/74	Aug. 16/74	9.74	8.69	12±1	1.22	100	Not Established

MACRO INVERTEBRATE POPULATION - AUGUST 1975 - CLINTON CREEK MINE

Species	Station			1			2			3			4			5			6			7			8			9		
	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C
Hydracarina sp.																														
Synurella sp.	1	14								9	6	10										3	5	2						
Moina macropa				3	3																									
Chaoborus sp.				1																										
Systemus sp.																														
Roederiodes sp.				1																										
Muscidae sp.																														
Corynoneura sp.	1	1	8				2															60	34	72		30	175			
Crictopus sp.																														
Spaniotoma sp.				16	6		3					17																		
Pentaneura sp.							1	10	4																					
Procladius sp.				14	1	6	8	6																						
Tendipes (=Chironomus) sp.							2																							
Tendipedidae sp pupa	1			1																										
Chrysops sp.				1																										
Longurio sp.	4	4	4																											
Tipula sp. b				1																										
Tipula sp. c				3																										
Tipula sp. d																														
Simuliidae sp.																														
Brachycera sp. pupa																														
Baetis sp.	2																													
Centroptilum sp.	1																													
Cinygmula sp.	2	6																												
Hemiptera sp nymph																														
Perloymia sp.																														
Nemoura sp.	2	4	1																											
Isogenus sp.	1																													
Arctopsyche sp.																														
Platycentropus sp.																														
Oligochaeta sp.				5																										

Clinton Creek. The tailings pile is composed of crushed serpentine (particle size \pm 0.6 cm) and asbestos fibres. The asbestos fibres are readily transported by the flowing waters of Wolverine Creek; leaving the serpentine to form the primary substrate. This substrate also shifts with increased water flows, making it a poor habitat for aquatic invertibrates.

The diversity-evenness results are presented in Table (9). Stations 1, 2, 3 and 7 had the greatest diversity. Stations 1 and 3 being slightly higher than 2 and 7. The low diversity results obtained for stations 5 and 6 can be attributed to the particulate serpentine which moves into Clinton Creek. The serpentine content of the substrate at Station 7 is far less than at Stations upstream. The higher diversity index reflects the improved habitat.

Varying quantities of asbestos fibres were present in the samplers at all stations except 1 and 3. Station 8 was plugged with fibres to a much greater extent than any other station. The Forty Mile River is a slower moving, deeper waterway than Clinton Creek and it is likely that the asbestos fibres settle out because of these characteristics. In addition to having the greatest concentration of asbestos fibres in the sampler, Station 8 had the greatest number of individuals.

The absence of organisms at Station 9 is due to the large fluctuations in water level at that location.

5.5 Fish

The electro fishing results are presented in Table (10). Fish were found at all stations with the exception of Stations 1, 3 and 4. The absence of fish at Station 1 is related to two factors. First, Clinton Creek upstream of Hudgeon Lake freezes during the winter and that the area between Station 2 and Hudgeon Lake is impassible. It is suspected that prior to the formation of Hudgeon Lake fish could have inhabited the upper reaches of Clinton Creek during

TABLE 8 INVERTEBRATE SPECIES LIST - AUGUST 1975

Phylum Arthropoda

Class Arachnida (Arachneidea)

Order Hydracarina

Hydracarina sp.

Class Crustacea

Order Amphipoda

Family Gammaridae

Synurella (bifurca or johanseni)

Order Cladocera

Sub Order Calyptomera

Family Daphnidae

prob. Moina macropa

Class Insecta

Order Diptera

Family Culicidae

Sub family Chaoborinae

Chaoborus sp.

Family Dolichopodidae

prob. Systemus sp.

Family Empididae

Hemerodromia sp.

Roederiodes sp.

Family Muscidae

Muscidae sp.

Family Tendipedidae

Sub family Hydrobaeninae

Corynoneura sp.

Cricotopus sp.

Spaniotoma sp.

Sub family Pelopiinae

Pentaneura sp.

Procladius sp.

Sub family Tendipedinae (Chironominae)

Tendipes (Chironomus) sp.

Adult Chironomidae

Tendipedidae sp. pupa

Family Tabanidae

Chrysops sp.

Family Tipulidae

Longurio sp.

Tipula sp. a

Tipula sp. b

Tipula sp. c

Tipula sp. d

Family Simuliidae (Melusinidae)

Simuliidae sp.

Brachycera pupa

Order Ephemeroptera

Family Baetidae

Baetis sp.

Centroptilum sp.

Family Heptageniidae

Cinygmula sp.

Order Hemiptera

Hemiptera sp. nymph

Order Plecoptera

Family Nemouridae

Sub family Leuctrinae

Perloymia sp.

Sub family Nemourinae

Nemoura sp.

Family Perlodidae

Sub family Isogeninae

Isogenus sp.

Order Tricoptera

Family Hydropsychidae

Arctopsyche sp.

Family Limnephilidae

Platycentropus sp.

Phylum Annelida

Class Oligochaeta

Oligochaeta sp.

the summer months. The absence of fish at Station 3 can be explained by similar circumstances. The absence of fish at Station 4 was due to the lack of water and the inability of the fish to pass through the culvert at the mouth of Wolverine Creek. The lack of food organisms and substrate characteristics being also contributing factors in explaining their absence.

The remaining stations were well populated by various species of fish. The most numerous and represented fish were the Longnose Sucker (*Catostomus catostomus*) and the Artic Grayling (*Thymallus arcticus*).

5.6 Histological Assessment

Tables 11 and 12 summarize the gill gissue structural changes sustained by the various experimental and control fish. The disorders identified and their significance are as follows:

(a) Separation of the Lamellae Epithelium

This signifys the dislodgement of the gill lamellae epithelia from the base membrane. It is a common response to toxicants in general (Figure 6).

(b) Aneurysm

This is a weakness in the blood vessel wall. The histopathology is unknown but this weakness is not commonly present in healthy fish.

(c) Hyperplasis

This signifys an increase in the epithelia cells of the gill lamellae. Also considered a common response to external irritants.

The sixteen day exposure group showed the highest incidence of separation of the lamellae epithelium. Eighty percent of the slides analyzed displayed 50 - 100 percent separation. Seventy percent of the slides of the 16 day exposure/16 day recovery group displayed 10% separation. These results indicate that this disorder became more prevalent with a

TABLE 9 DIVERSITY AND EVENNESS RESULTS
INVERTEBRATE POPULATION - AUGUST 1975 - CLINTON CREEK MINE

Station Number		Total Number of Individuals	Number of Taxa	Diversity Index	Evenness
1	A	14	8	2.807	0.936
	B	33	7	2.330	0.830
	C	29	7	1.975	0.703
2	A	36	9	2.460	0.776
	B	6	3	1.459	0.921
	C	12	5	1.896	0.817
3	A	40	8	2.622	0.874
	B	29	7	2.643	0.941
	C	48	8	2.236	0.745
4	A	0	0	0	0
	B	0	0	0	0
	C	0	0	0	0
5	A	4	2	0.811	0.811
	B	108	2	0.076	0.076
	C	10	5	2.122	0.914
6	A	22	4	0.957	0.478
	B	0	0	0	0
	C	0	0	0	0
7	A	88	6	1.412	0.546
	B	60	10	2.142	0.645
	C	94	6	1.141	0.442
8	A	0	0	0	0
	B	33	4	0.584	0.292
	C	186	4	0.417	0.209
9	A	0	0	0	0
	B	0	0	0	0
	C	3	2	0.918	0.918

TABLE 10: ELECTRO FISHING RESULTS, AUGUST 6 - 7, 1975

Scientific Name	Common Name	Total Number at Stations								
		1	2	3	4	5	6	7	8	9
Catostomus catostomus (Forster)	Longnose sucker		1			4	2	3	5	5
Coregonus clupeaformis (Mitchill)	Lake whitefish							1		
Cottus cognatus (Richardson)	Slimy sculpin		1				1	3	1	2
Oncorhynchus tshawytscha (Walbaum)	Chinook salmon									2
Prosopium cylindraceum (Pallas)	Round white fish		2					1		
Thymallus arcticus (Pallas)	Arctic grayling		14			2	1	2	2	2

greater exposure time and also that the fish recovered substantially after being exposed to clean water.

Both experimental and control fish displayed signs of hyperplasia and development of aneurysm and consequently these disorders cannot be attributed to the presence of asbestos fibres.

There has been limited research done on the effects of physical irritants to fish gill tissues. Smith, 1966, found that the Fathead minnow (*Pimephales promelas*) show an inverse relationship between oxygen uptake and the concentration of wood fibres; i.e., the higher the concentration of fibres, the lower the oxygen uptake. Smith, 1966, also found that fish which were previously exposed to wood fibres had metabolic levels similar to the control fish, indicating that previous exposure to wood fibres did not cause a permanently depressed metabolic level. Smith, 1966, also measured oxygen uptake and his results indicated that suspended wood fibres greatly affect a fish's ability to extract oxygen from water. When exposed to lethal concentrations of wood fibres, the fish showed signs of anoxia. When he investigated the gill tissue, it was found that the test fish had responded by increasing the production of mucus and that a large amount of mucus was present between the gill lamellae. In some instances the lamellae showed signs of malformation and fusion. Since gills serve as a respiratory and osmoregulatory organ, fish are required to circulate a large volume of water over their gills. Gills are affected and damaged when they are exposed to chemical toxicants and physical irritants (Smith, 1966; Cope, 1966; Gilderhus, 1966; Migaki, 1975; Skidmore, J.F.; 1970).

The present knowledge of the effects of asbestos fibres on aquatic organisms is very limited.



FIGURE 8 Arrows point to areas which
have separation of lamellae epithelium




FIGURE 9 Arrows indicate hyperplasia of lamellae epithelium.




FIGURE 10 Arrows shows site of the Aneurysm.

CONCLUSIONS AND RECOMMENDATIONS

The water chemistry of the area near to the tailings and waste rock pile has been affected as indicated by elevated levels of Calcium, Magnesium, Iron Manganese, Potassium, Turbidity and Hardness at Stations 2, 4, 5 and 6. The values obtained are not significantly high to be considered harmful to aquatic organisms. These parameters should be continually monitored.

The samples analyzed for asbestos fibre concentration in 1974 were within the expected range. However, the sample collected from Hudgeon Lake in 1975 had a higher asbestos fibre content in comparison to other background levels reported elsewhere in Canada. The fibre length determinations done in 1974 probably reflect the proximity to an airborne asbestos source. It is recommended that the asbestos fibre content of Hudgeon Lake, Clinton Creek and The Forty Mile River receive continual monitoring including fibre length determinations. The information obtained and presented in this report is limited but clarifys the need to do additional and more complete studies.

Invertebrate communities have been adversely affected by substrate alteration particularly in the vicinity of the tailings pile (Station 4). The effects of the serpentine content of the substrate are evident as far as Station 7 on Clinton Creek.

TABLE 1: HISTOLOGICAL ASSESSMENT
- SLIDES ANALYSED


Treatment	Seperation of Lamellae Epithelium % of Gill Tissue Affected				Hyperplasm Present	Aneurysm Present
	0%	10%	10-50%	50-100%		
4 days control						
Slide 13	X					
16	X					
23	X					
28	X					
33	X					
37	X					
41	X					
42	X					
51	X					
61	X					
4 days exposure						
slide 4	X					
10	X					
17	X					
29		X				
38		X				
43			X			
47		X				
50	X					
8 days control						
slide 1	X					
5	X					X
21		X				
24	X				X	
30			X		X	X
40	X					
44	X				X	
53	X					
55	X					
57	X					
8 days exposure						
slide 3		X			X	
6		X				
11			X			
15			X			
20		X				
25		X				
31		X				
34		X				
36		X				
39		X				
45		X				
48		X				
52	X					
16 days exposure						
slide 8				X		
9				X		
14				X		
18			X			
22				X		
26				X		
49				X		
56				X		
58				X		
60		X				
16 days exposure and 16 days recovery						
slide 2	X				X	X
7		X			X	
12			X			
19		X				
27		X				
32		X				
35		X				
46		X				
54	X					
59		X				

TABLE 12: HISTOLOGICAL ASSESSMENT OF GILL TISSUE (ONCORHYNCHUS KITSUTCH

Percentage of Slides Showing the Following Histological Disorders						
Treatment	Seperation of Lamellae Epithelium				Hyperplasia	Aneurysm
	0%	10%	10-50%	50-100%		
4 days control	100					
4 days exposure	50	37.5	12.5			
8 days control	80	10	10		30	20
8 days exposure	7.7	76.9	15.4		7.7	
16 days exposure		10	10	80		
16 days exposure with 16 days recovery	20	70	10		20	20

The fish populations have also been adversely affected by the activities of the Clinton Creek Mine. The absence of fish at Stations 1, 3 and 4 has been attributed to the impoundments responsible for the creation of Wolverine Lake and Hudgeon Lake. In addition, substrate alterations and subsequent lack of food organism being contributing factors in explaining their absence.

The results of the histological studies indicate an additional detrimental effect on the fish populations directly attributable to the presence of asbestos fibres. The greatest number of fish showing separation of the lamellae epithelium of the gill tissue occurred in the greatest exposure group (16 day^s). It is recommended that these studies be continued.



REFERENCES

- Cope, O.B., 1966. Contamination of the freshwater ecosystem by pesticides. J. Appl. Ecol. 3 (Suppl) L 33-44.
- Craig, Laporte, 1972. North of 60 Mineral Industry Report, 1973, Yukon Territory EGS 1975-7. Northern Economic Development Branch, Department of Indian Affairs and Northern Development. Government of Canada pp 90.
- Davidson et al, 1972. Pollution Sampling Handbook. Environment Canada. Fisheries Service, Pacific Region.
- Durham, R.W. and T. Pang, 1975. Asbestos Fibers in Lake Superior. American Society Testing Materials, Special Publication No. 573.
- Durham, R.W., 1976. Personal communications between R.W. Durham and G. Hebert 1976.
- Gilderhus, P.A., 1966. Some effects of sublethal concentrations of the sodium arsenite on bluegills and the aquatic environment. Trans. Amer. Fish. Soc. 95: 289-296.
- Golder & Associates, 1976. Report to Cassiar Asbestos Corporation, Ltd. on Stabolization Measures for Mine Closure. Clinton Mine, Yukon Territories Draft, May 1976.
- Kay, G.H., 1973. Asbestos in Drinking Water. Journal Water Pollution Control Federation, August 1973.
- Lawrence, J. and H. Walter Zimmermann, 1977. Asbestos in Water: Mining and Processing Effluent Treatment. Canada Center for Inland Waters, Burlington, Ontario pp. 156-160.
- Migaki, G. and W.E. Ribelin, 1975. The Pathology of Fishes. The University of Wisconsin Press.

- Pielou, E.C. 1966. The Measurement of Diversity in Different Types of Biological Collection. J. Theoret, Biol. 13 pp. 131-144.
- Sinclair, Gilbert 1975. North of 60, Mineral Industry Report, 1975. Yukon Territory EGS 1975-7. Northern Economic Development Branch, Department of Indian Affairs and Northern Development, Government of Canada. pp. 69-72.
- Sinclair, Morin, Craig, Marchand 1976. North of 60, Mineral Industry Reprot 1975, Yukon Territory EGS 1976-15. Northern Economic Development Branch, Department of Indian Affairs and Northern Development, Government of Canada pp. 71-73.
- Skidmore, J.F. 1970. Respiration and Osmoregulation in Rainbow Trout with Gills damaged by Zinc Sulphate. J. Exp. Biol. 52: 481-494.
- Smith, L.L. 1966. Effect of Pulpwood Fiber on Oxygen consumption and swimming Endurance of the Fathead Minnow, Pemephales promelas. Trans. Am. Fish. Soc. 95 (1): 71-84.
- Standard Methods 1971. Landucci, Robert, development of Standard Methods System and Detailed Hypothsis of the Periphyton Modification of the Bermolli Principle. pp. 467-483.
- Wilhm, J.L. and T.C. Dorris 1968. Biological Parameters for Water Quality Criteria. Bioscience 18, 477-481.