

**LEGEND**  
Coloured legend blocks indicate map units that appear on this map. This legend is common to maps 1876A-1879A.

**CENOZOIC**

**QUATERNARY**

**HOLOCENE - POST-McCONNELL GLACIATION**

- O** Bog, fen, and swamp deposits: undivided; thickness < 1 m to 10 m
- Ap** **Alluvial deposits:** gravel to silt size sediments deposited by streams
- At** **Alluvial terrace sediments:** gravel, cobble to pebble; massive to thick bedded capped by sand and silt; flat lying; includes lacustrine and organic deposits in alluvial channels and backswamp areas subject to periodic inundation and reworking by floods; thickness 1 to 5 m
- Al** **Alluvial fan sediments:** gravel, cobble to pebble with a sandy matrix; massive to thick bedded; capped by sand and silt; sediments are of flood plain origin now isolated from flooding by stream incision; thickness 1 m to 10 m or more
- Af** **Alluvial fan sediments:** gravel, sand, silt, and diamicton, poorly sorted; thick bedded to massive; sediments from fan-shaped landforms at the confluence of tributary streams with lower gradient trunk streams; subject to flooding accompanied by sudden stream migration and inundation by debris flows on fans with gradients in excess of 4%; thickness up to 10 m or more
- Au** **Alluvial sediments:** undivided sediments forming floodplains, fans, and terraces as above that cannot be subdivided at this map scale

**PLEISTOCENE AND HOLOCENE (UNDIVIDED)**

- Eb** **Eolian sands:** sand, well sorted; massive; forms crescent shape and linear dunes and features; gently undulating inter-dune eolian plains; thickness 1 to 5 m
- Cu** **Colluvial deposits:** stony diamicton resulting from the physical and chemical breakdown of bedrock and reworking and transportation by creep, solifluction, debris flow, snow avalanching, and rockfall; also includes diamicton created by landsliding. Colluvial deposits may contain reworked glacial sediments within the limits of ice cover during the Reid and McConnell glaciations. Colluvial deposits beyond the limits of the McConnell Glaciation ice cover are likely the product of continuous formation and reworking over a significant part of the Pleistocene
- Cb** **Colluvial blanket sediments:** diamicton, stony with a sandy matrix; massive; surface conforms to underlying bedrock or buried glacial deposits; thickness > 1 m to 50 m or more in large landforms
- Cv** **Colluvial veneer sediments:** diamicton, stony with a sandy matrix; massive; thickness < 1 m to discontinuous over bedrock
- Ca** **Colluvial apron sediments:** diamicton, bouldery diamicton and bouldery sandy gravel, poorly sorted; massive; sediments form a wedge-like slope-ice complex of small steep slopes; flow and avalanche-dominated fans and solifluction deposits; thickness < 1 m at top and down slope limit up to 5 m or more in the thickest part of the apron
- bCa** **Rockfall sediments:** boulders, angular; massive; deposits form as rockfall accumulations along the bases of bedrock slopes; thickness ranges from < 1 m at top to up to 10 m

**LATE PLEISTOCENE (WISCONSINAN) - McCONNELL GLACIATION**

- Gp** **Glaciolacustrine deposits:** well stratified sand, silt, clay, and minor silt and clay, moderately to well sorted and becomes finer downward; massive to thick bedded; planar surface, deposit of delta form in plan view; thickness > 5 m
- Gt** **Glaciolacustrine terrace sediments:** pebble to cobble gravel; massive to thick bedded; incised into flights of terraces by glacial streams; thickness 1 to > 10 m
- Gd** **Glaciolacustrine delta sediments:** sand, gravel, and minor silt and clay, moderately to well sorted, texture becomes finer downward; massive to thick bedded; deposit has a planar surface and delta form in plan view; thickness > 5 m
- Gx** **Glaciolacustrine ice stagnation complex sediments:** gravel, sand, diamicton, poorly to moderately sorted, and minor silt and clay; bedding thick to massive and commonly folded and faulted from syndepositional ice meltout; surface consists of hummocks, kettles, esker and crevasse-fill ridges with minor elements of units G<sub>1</sub>, G<sub>2</sub>, G<sub>3</sub>, and G<sub>4</sub>
- Gu** **Discontinuous glaciolacustrine sediments:** gravel and sand including elements of units Gp and Gx, discontinuously distributed in areas of units Mb and Mv
- Mb** **Moraine deposits (TILL):** glacial diamicton, mainly silt, generally consisting of a matrix ranging from sand to clay that supports clasts ranging from boulders to pebbles in size, deposited either directly from glacial ice or by gravity flow from glacial ice
- Mv** **Till veneer:** diamicton, stony with a silty, sandy matrix; massive to crudely stratified; may contain extensive areas of thin (< 1 m) to patchy colluvium over bedrock

**MIDDLE PLEISTOCENE - PRE-McCONNELL GLACIATION (UNDIVIDED)**

- A<sup>pl</sup>** **Alluvial fans:** single fans or aprons of coalesced fans formed of gravel and sand, poorly to moderately sorted, thick bedded. Sediments disturbed by cryoturbation and clasts commonly wind scouted. Thickness up to 10 m or more
- A<sup>mx</sup>** **Alluvial complex sediments:** gravel and sand, poorly to moderately sorted; thin to thick bedded; interstratified with colluvial diamicton, reworked beds, peat, and woody debris; sediments underlie the floors and margins of narrow upland valleys and grade laterally upslope into colluvial blankets. They contain segregated ice lenses and ice wedges and are normally capped by blanket bog; sediments may represent several depositional cycles; thicknesses may exceed 10 m in mid-valley locations

**MIDDLE PLEISTOCENE - REID GLACIATION**

- A<sup>rl</sup>** **Alluvial deposits:** complexes of nonglacial and fan sands and gravels deposited by streams that flowed from ice-free areas toward Reid Glaciation ice margins. These sands and gravels locally overlie older interglacial gravels that contain glacial gold
- A<sup>rl</sup>** **Alluvial terrace sediments:** gravelly micaceous sand and gravel, moderately sorted, clasts angular to subangular; bedding is thin to massive and terraced gravel clasts are commonly frost shattered and wind scouted. Sediments have been incised into flights of terraces. Sediments are commonly cut by ice wedge waters pseudomorphs over their upper 2 m (includes terrace gravels along Klaza River possibly deposited by outlet waters from a lake dammed by a glacial margin during Reid Glaciation). Thickness 1 to 15 m
- A<sup>rlx</sup>** **Alluvial complex sediments:** gravel and sand, poorly to moderately sorted; thin to thick bedded; interstratified with colluvial diamicton, reworked beds, peat, and woody debris; sediments underlie the floors and margins of narrow upland valleys and grade laterally upslope into colluvial blankets. They contain segregated ice lenses and ice wedges and are normally capped by blanket bog; sediments may represent several depositional cycles; thicknesses may exceed 10 m in mid-valley locations
- L<sup>rp</sup>** **Glaciolacustrine plain:** sand, silt, and clay, with minor dropstones; thinly bedded to laminated; thickness 1 to > 5 m
- G<sup>lp</sup>** **Glaciolacustrine terrace sediments:** gravel and sand, moderately to well sorted; thick bedded to massive; planar surface; thickness 1 to 10 m or more
- G<sup>lt</sup>** **Glaciolacustrine terrace sediments:** pebble to cobble gravel; massive to thick bedded; incised into flights of terraces by glacial streams; thickness 1 to > 10 m
- G<sup>ld</sup>** **Glaciolacustrine delta sediments:** sand, gravel, and minor silt and clay, moderately to well sorted and becomes finer downward; massive to thick bedded; planar surface, deposit of delta form in plan view; thickness > 5 m
- G<sup>lx</sup>** **Glaciolacustrine ice stagnation complex sediments:** gravel, sand, diamicton, poorly to moderately sorted, and minor silt and clay; bedding thick to massive and commonly folded and faulted from syndepositional ice meltout; surface consists of hummocks, kettles, esker and crevasse-fill ridges with minor elements of units G<sub>1</sub>, G<sub>2</sub>, G<sub>3</sub>, and G<sub>4</sub>
- M<sup>rb</sup>** **Moraine deposits (TILL):** glacial diamicton, mainly silt, generally consisting of a matrix ranging from sand to clay that supports clasts ranging from boulders to pebbles in size, deposited either directly from glacial ice or by gravity flow from glacial ice
- M<sup>rv</sup>** **Till blanket:** diamicton, stony, silty sandy matrix, massive; conforms to underlying topography; thickness 1 to 5 m
- M<sup>rv</sup>** **Till veneer:** diamicton, stony, silty sandy matrix, massive; discontinuous and may contain extensive areas of thin (< 1 m) and patchy colluvium over bedrock

**EARLY PLEISTOCENE - YOUNGER PRE-REID GLACIATION**

- G<sup>mp</sup>** **Glaciolacustrine plain sediments:** gravel and sand, deeply weathered; forms an unsorted plain
- G<sup>mt</sup>** **Glaciolacustrine terrace sediments:** gravel and sand, deeply weathered; incised into flights of terraces
- M<sup>rv</sup>** **Moraine deposits (TILL):** glacial diamicton, mainly silt, generally consisting of a matrix ranging from sand to clay that supports clasts ranging from boulders to pebbles in size, deposited either directly from glacial ice or by gravity flow from glacial ice
- M<sup>rv</sup>** **Till veneer:** patchy, deeply weathered diamicton. Matrix sandy silty clay. Formerly tillcap-rich stones are weathered to clay

**EARLY PLEISTOCENE**

- V** **VOLCANIC ROCK AND INTERSTRATIFIED SEDIMENTS**  
**Pleistocene volcanics (undivided):** basalt, breccia, volcanic ejecta and tephra; andesite, rhyolite, and dacite; andesite and rhyolite of the Selkirk volcanics erupted during the early and late Pleistocene or early Holocene epochs in the Fort Selkirk area. Cumulative basalt flow thicknesses exceed 100 m where they have filled valleys. Deposits of the two known pre-Reid glaciations and at least one nonglacial period are locally interstratified with the volcanics and are exposed only in sections

**PALEOZOIC AND MESOZOIC**

- R** **PRE-QUATERNARY BEDROCK:** basalt, andesite, gneiss, schist, gneiss, greywacke, granodiorite and monzonite; includes areas of thin colluvial cover, blockfields, sorted stone polygons in alpine areas
- R-A** **AVAILANCHE MODIFIED PRE-QUATERNARY BEDROCK:** bedrock areas subject to rapid mass wasting processes (rockfall and snow avalanches)

**SYMBOLS**

Note: pR - pre-Reid Glaciation, r - Reid Glaciation, pm - pre-McConnell Glaciation, (no designator, assume McConnell Glaciation)

- Geological boundary
- Blanket bog covering generally less than 1 m thick
- Discontinuous eolian sands or silts, thickness locally up to 2 m
- Open system pingo, collapsed open system pingo
- Thermokarst collapse activity
- Landslide, arrow(s) indicate direction of movement
- Cirque: degraded cirque active prior to McConnell Glaciation
- Arête: degraded arête active prior to McConnell Glaciation
- Streamlined glacial bedforms: ice flow direction known, unknown
- Meltwater channel: large, small ice-walled channel, arrow indicates flow direction
- Esker: flow direction defined, unknown
- End moraine
- Recessional moraine
- Ice-contact face in stratified drift (teeth on ice side)
- Ice limit
- Cryoplanation terrace
- Tor
- Vertebrate fossil locality
- Stratigraphic contact

**LEGEND FOR STRATIGRAPHIC SECTIONS**

**QUATERNARY**

**POSTGLACIAL**


- Alluvial (Holocene) sands and gravels

**MIDDLE TO LATE PLEISTOCENE**

- Glaciolacustrine sediments, McConnell Glaciation
- Palaeosol developed between Reid and McConnell Glaciation
- Stream deposited sediments (pre-McConnell Glaciation) of nonglacial origin (includes terrace gravels along Klaza River possibly deposited by outlet waters from a lake dammed by a glacial margin along the southern margin of Dawson range during Reid Glaciation)

**EARLY TO MIDDLE PLEISTOCENE**

- Till, Reid Glaciation
- Glaciolacustrine sediment, Reid Glaciation
- Glaciolacustrine sediments, Reid Glaciation
- Alluvial deposits, deposited between the younger pre-Reid glaciation and Reid Glaciation



Surficial geology by L.E. Jackson Jr. 1988-1992, Geological Survey of Canada  
Digital cartography by J.D. Naraway, Geoscience Information Division  
Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

MAP 1878A  
SURFICIAL GEOLOGY  
**GRANITE CANYON**  
YUKON TERRITORY

Scale 1:100 000 - Échelle 1/100 000

Kilometres 0 2 4 6 8 Kilomètres

Universal Transverse Mercator Projection / Projection transverse de Mercator  
© Crown copyrights reserved / © Droits de la Couronne réservés

Mean magnetic declination 1997, 28°28' E, decreasing 12.1' annually. Readings vary from 28°11' E in the SW corner to 28°44' E in the NE corner of the map

Elevations in feet above mean sea level

Base map assembled and modified by the Geoscience Information Division from maps 115-010 (1986), 115-019 (1982), 115-015, 16 (1961) published at the scale of 1:50 000 by the Surveys and Mapping Branch

Copies of the topographical editions covering this map area may be obtained from the Canada Map Office, Natural Resources Canada, Ottawa, Ontario, K1A 0G9

115-013	115-014	115-015	115-016
1877A	1878A	1879A	
115-012	115-011	115-010	115-009
115-015	115-016	115-017	115-018
1876A	1878A		
115-014	115-013	115-012	115-011

ESIC CIST  
JUL 17 1997  
Earth Sciences Sector / Secteur des sciences de la Terre  
MAP LIBRARY / CARTOTHEQUE

NOT TO BE TAKEN FROM LIBRARY / NE PAS SORTIR DE LA BIBLIOTHÈQUE

1878A