



Copies of this map may be obtained from the Geological Survey of Canada, 601 Booth Street, Ottawa, Ontario, K1A 0E8. 3563-33rd Street, N.W., Calgary, Alberta T2L 2A7.



Surficial geology by L.E. Jackson, Jr. 1988-1992, Geological Survey of Canada.  
 Digital cartography by J.D. Nansaway, Geoscience Information Division.  
 Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada.

MAP 1879A  
 SURFICIAL GEOLOGY  
**TANTALUS BUTTE**  
 YUKON TERRITORY  
 Scale 1:100 000 - Échelle 1/100 000  
 Kilometers 2 4 6 8 Kilometers  
 Universal Transverse Mercator Projection / Projection transverse de Mercator  
 © Crown copyright reserved

Base map assembled and modified by the Geoscience Information Division from maps 115-47 (1984), 115-42 (1989), 115-11 (1985), 115-48 (1981) published at the scale of 1:50 000 by the Surveys and Mapping Branch.  
 Copies of the topographical editions covering this map area may be obtained from the Canada Map Office, Natural Resources Canada, Ottawa, Ontario, K1A 0E9.  
 Mean magnetic declination 1997, 28°06' E, decreasing 11.9' annually. Readings vary from 27°50' E in the SW corner to 28°22' E in the NE corner of the map.  
 Elevations in feet above mean sea level

115-013	115-014	115-015	115-016
1877A	1878A	1879A	
115-012	115-011	115-010	115-009
115-005	115-006	115-007	115-008
1876A	1876B	1876C	
115-004	115-003	115-002	115-001

### LEGEND

Coloured legend blocks indicate map units that appear on this map. The legend is common to maps 1876A-1879A.

#### CENOZOIC

##### QUATERNARY

###### HOLOCENE - POST-McCONNELL GLACIATION

- O** Bog, fen, and swamp deposits: undivided, thickness < 1 m to 10 m
- Ap** Floodplain sediments: gravel, cobble to pebbles; massive to thick bedded; capped by sand and silt; flat lying; includes lacustrine and organic deposits in abandoned channels and backswamp areas subject to periodic inundation and reworking by floods; thickness 1 to 5 m
- At** Alluvial terrace sediments: gravel, cobble to pebbles with a sandy matrix; massive to thick bedded; capped by sand and silt; sediments are of flood plain origin now isolated from flooding by stream incisions; thickness 1 to 10 m or more
- At'** Alluvial fan sediments: gravel, sand, silt, and diamicton, poorly sorted; thick bedded to massive; sediments form fan-shaped landforms at the confluence of tributary streams with lower gradient streams; subject to flooding accompanied by sudden stream migration and truncation by debris flows on fans with gradients in excess of 4%; thickness up to 10 m or more
- Au** Alluvial sediments, undivided: sediments forming floodplains, fans, and terraces as above that cannot be subdivided at this map scale

###### PLEISTOCENE AND HOLOCENE (UNDIVIDED)

- Eb** Eolian deposits: well sorted medium sand to coarse silt transported and deposited by wind action during the early Pleistocene and McConnell Glaciation. Thin deposits of very fine sand and coarse silt < 1 m thick are deposited discontinuously throughout low lying areas (see SYMBOLS). Below:
  - Eb** Eolian sands: sand, well sorted, massive; forms crescent-shaped and linear dunes and features or gently undulating inter-dune eolian plains; thickness 1 to 5 m
- Cv** COLLUVIAL DEPOSITS: stony diamicton resulting from the physical and chemical breakdown of bedrock and reworking and transportation by creep, solifluction, debris flow, snow avalanching, and rockfall. It also includes diamicton created by landsliding. Colluvial deposits may contain reworked glacial sediments within the limits of ice cover during the Reid and McConnell glaciations. Colluvial deposits beyond the limits of the McConnell Glaciation ice cover are likely the product of continuous formation and reworking over a significant part of the Pleistocene
- Cb** Colluvial blanket sediments: diamicton, stony with a sandy matrix; massive; surface conforms to underlying bedrock or buried glacial deposits; thickness > 1 m to 50 m or more in large landslides
- Cv** Colluvial veneer sediments: diamicton, stony with a sandy matrix; massive; deposit conforms to underlying topography; thickness 1 to 5 m
- Ca** Colluvial apron sediments: diamicton, bouldery diamicton and bouldery sandy gravel, poorly sorted; massive; sediments form a wedge-like slope toe complex of small steep-sided fans and avalanche dominated fans and solifluction deposits; thicknesses < 1 m at up and down slope limit to up to 5 m or more in the thickest part of the apron
- bCa** Rockfall sediments: boulders, angular, massive; deposits form as rockfall accumulations along the bases of steep bedrock slopes; thickness ranges from < 1 m to meters to 10 to 100 m

###### LATE PLEISTOCENE (WISCONSINAN) - McCONNELL GLACIATION

- Lp** Glaciolacustrine plain: sand, silt, and clay with minor dropstones; thinly bedded to laminated; thickness > 5 m
- Lb** Glaciolacustrine blanket: silt and clay with minor sand; thinly bedded to laminated; deposit conforms to underlying topography; thickness 1 m to 5 m
- Lv** Glaciolacustrine veneer: silt and clay with minor sand; thinly bedded to laminated; deposit conforms to underlying topography; thickness < 1 m to discontinuous
- Lx** Ice-contact glaciolacustrine complex: sand, silt, and clay; laminated to medium bedded with up to 10 percent textural beds of gravel and diamicton and dropstones; surface is hummocky, pitted, and ridged; thickness > 5 m
- Gp** GLACIOFLUVIAL DEPOSITS: sand, gravels and minor silts > 1 m thick deposited by streams flowing away from, or in contact with, glacial ice including deltas graded to former glacial lakes. Sorting ranges from good to poor and stratification from thin bedded to massive. Sediments commonly display evidence of syndepositional collapse due to meltout of buried or supporting ice
- Gt** Glaciolacustrine terrace sediments: pebble to cobble gravel; massive to thick bedded; capped by sand and silt; planar surface; thickness 1 to > 10 m
- Gt'** Glaciolacustrine terrace sediments: pebble to cobble gravel; massive to thick bedded; incised into flights of terraces by glacial streams; thickness 1 to > 10 m
- Gd** Glaciolacustrine delta sediments: sand, gravel, and minor silt and clay; moderately to well sorted; texture becomes finer downward; massive to thick bedded; deposit has a planar surface and delta form in plan view; thickness > 5 m
- Gx** Glaciolacustrine ice stagnation complex sediments: gravel, sand, diamicton, poorly to moderately sorted, and minor silt and clay; bedding thick to massive and commonly folded and faulted from syndepositional ice meltout; surface consists of hummocks, kettle, esker and crevasse-fill ridges with minor elements of units Gp, Gd, and Gt
- Gu** Discontinuous glaciolacustrine sediments: gravel and sand including elements of units Gp and Gx; discontinuously distributed areas of units Mb and Mv
- Mb** MORAINAL DEPOSITS (TLL): glacial diamicton, mainly silt, generally consisting of a matrix ranging from sand to clay that supports clasts ranging from boulders to pebbles in size; deposited either directly from glacial ice or by gravity flow from glacial ice
- Mv** Till blanket: diamicton, stony with a silty, sandy matrix; massive to crudely stratified; surface conforms to the underlying topography; thickness 1 to 5 m
- Mv'** Till veneer: diamicton, stony with a silty, sandy matrix; massive to crudely stratified; may contain extensive areas of thin (< 1 m) to patchy colluvium over bedrock

###### MIDDLE PLEISTOCENE - PRE-McCONNELL GLACIATION (UNDIVIDED)

- A<sup>m</sup>** ALLUVIAL DEPOSITS: gravel and sand deposited by streams that were not tied to glacial meltwaters. Sediments may represent several cycles of alluviation and erosion. Sediments are not presently correlative to past glaciations but presumably predate McConnell Glaciation due to the presence of McConnell age loess overlying them. Basal gravels within these sediments commonly contain placer gold in basins draining Cretaceous granitoids and andesites
- A<sup>m</sup>** Alluvial fans: single fans or aprons of coalesced fans formed of gravel and sand, poorly to moderately sorted, thick bedded. Sediments disturbed by cryoturbation and clasts commonly wind sculpted. Thicknesses up to 10 m or more
- A<sup>m</sup>** Alluvial complex sediments: gravel and sand, poorly to moderately sorted; thin to thick bedded, interstratified with colluvial diamicton, reworked loess, peat, and woody detritus; sediments underlie the floors and margins of narrow upland valleys and grade laterally (upslope) into colluvial blankets. They contain segregated ice lenses and ice wedges and are normally capped by blanket bog sediments; may represent several depositional cycles; thicknesses may exceed 10 m in mid-valley locations

###### MIDDLE PLEISTOCENE - REID GLACIATION

- A<sup>r</sup>** ALLUVIAL DEPOSITS: complexes of nonglacial and fan sands and gravels deposited by streams that flowed from ice-free areas toward Reid Glaciation ice margins. These sands and gravels locally overlie older interglacial gravels that contain placer gold
- A<sup>r</sup>** Alluvial terrace sediments: gravelly micaceous sand and gravel, moderately sorted; clasts angular to subangular; bedding thin to massive and lenticular; gravel clasts are commonly flat shattered and wind sculpted; sediments have been incised into flights of terraces. Sediments are commonly cut by ice wedge water pseudomorphs over their upper 2 m; includes terrace gravels along Klaxa River possibly deposited by outlet waters from a lake dammed by a glacial margin during Reid Glaciation. Thickness 1 to 15 m
- A<sup>r</sup>** Alluvial fans: single fans or aprons of coalesced fans formed of gravel and sand, poorly to moderately sorted; thick bedded. Sediments disturbed by cryoturbation and clasts commonly wind sculpted. Thickness up to 10 m or more
- A<sup>r</sup>** Alluvial complex sediments: gravel and sand, poorly to moderately sorted; thin to thick bedded, interstratified with colluvial diamicton, reworked loess, peat, and woody detritus; sediments underlie the floors and margins of narrow upland valleys and grade laterally (upslope) into colluvial blankets. They contain segregated ice lenses and ice wedges and are normally capped by blanket bog sediments; may represent several depositional cycles; thicknesses may exceed 10 m in mid-valley locations
- L<sup>p</sup>** GLACIOFLUVIAL DEPOSITS: well stratified sand, silt, and minor gravel and diamicton deposited in lakes ponded by glacial ice. Glaciolacustrine silts and clays commonly contain aggregated ground ice and are affected by cryoturbation and reworking by floods; thickness 1 to > 5 m
- L<sup>p</sup>** Glaciolacustrine plain: sand, silt, and clay, with minor dropstones; thinly bedded to laminated; thickness 1 to > 5 m
- G<sup>p</sup>** GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from, or in contact with, glacial ice
- G<sup>p</sup>** Glaciolacustrine terrace sediments: gravel and sand, moderately to well sorted; thick bedded to massive; planar surface; thickness 1 to > 10 m or more
- G<sup>t</sup>** Glaciolacustrine terrace sediments: pebble to cobble gravel; massive to thick bedded; incised into flights of terraces by glacial streams; thickness 1 to > 10 m
- G<sup>d</sup>** Glaciolacustrine delta sediments: sand, gravel, and minor silt and clay; moderately to well sorted and becomes finer downward; massive to thick bedded; deposit has a planar surface and delta form in plan view; thickness > 5 m
- G<sup>x</sup>** Glaciolacustrine ice stagnation complex sediments: gravel, sand, diamicton, poorly to moderately sorted, and minor silt and clay; bedding thick to massive and commonly folded and faulted from syndepositional ice meltout; surface consists of hummocks, kettle, esker and crevasse-fill ridges with minor elements of units G<sup>p</sup>, G<sup>d</sup>, and G<sup>t</sup>
- M<sup>b</sup>** MORAINAL DEPOSITS (TLL): glacial diamicton, mainly silt, generally consisting of a matrix ranging from sand to clay that supports clasts ranging from boulders to pebbles in size; deposited either directly from glacial ice or by gravity flow from glacial ice
- M<sup>v</sup>** Till blanket: diamicton, stony, silty sandy matrix; massive; conforms to underlying topography; thickness 1 to 5 m
- M<sup>v</sup>** Till veneer: diamicton, stony, silty sandy matrix; massive; discontinuous and may contain extensive areas of thin (< 1 m) to patchy colluvium over bedrock

###### EARLY PLEISTOCENE - YOUNGER PRE-REID GLACIATION

- G<sup>l</sup>** GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice margins during the early Pleistocene. They are deposited to massive, clasts, except for quartz, quartzite, and chert are disaggregated or weathered to clay over the upper 2 m of the sediments where they underlie the surface; clasts near the surface of the unit are intensely wind sculpted and the internal's cut by ice wedge pseudomorphs and sand wedges; thickness 1 m to > 5 m
- G<sup>m</sup>** Glaciolacustrine plain sediments: gravel and sand, deeply weathered; forms an unincised plain
- G<sup>t</sup>** Glaciolacustrine terrace sediments: gravel and sand, deeply weathered; incised into flights of terraces
- M<sup>v</sup>** MORAINAL DEPOSITS (TLL): glacial diamicton, mainly silt, generally consisting of a matrix ranging from sand to clay that supports clasts ranging from boulders to pebbles in size; deposited either directly from glacial ice or by gravity flow from glacial ice
- M<sup>v</sup>** Till veneer: patchy, deeply weathered diamicton. Matrix silty sandy clay. Formerly kelpar-rich stones are weathered to clay

###### EARLY PLEISTOCENE

- V** VOLCANIC ROCK AND INTERSTRATIFIED SEDIMENTS
  - Pleistocene volcanics (undivided): basalt, trachyte, volcanic ejecta and tephra; Pleistocene or early Holocene epochs in the Fort Selkirk area. Cultivite basal flow thicknesses exceed 100 m where they have filled valleys. Deposits of the two known pre-Reid glaciations and at least one nonglacial period are locally interstratified with the volcanics and are exposed only in sections

###### PALEOZOIC AND MESOZOIC

- R** PRE-QUATERNARY BEDROCK: basal andesite, gneiss, schist, quartzite, gneiss, granitoid and monzonite; includes areas of thin colluvial cover, blockfields, sorted stone polygons in aprone areas
- R-A** AVALANCHE MODIFIED PRE-QUATERNARY BEDROCK: bedrock areas subject to rapid mass wasting processes (rockfall and snow-avalanches)

### SYMBOLS

Note: pR - pre-Reid glaciation; r - Reid Glaciation; pM - pre-McConnell Glaciation; (no designator, assume McConnell Glaciation)

- Geological boundary
- Blanket bog covering generally less than 1 m thick
- Discontinuous eolian sands or silts, thickness locally up to 2 m
- Open system pingo, collapsed open system pingo
- Thermokarst collapse activity
- Landslide, arrow(s) indicate direction of movement
- Cirque; degraded cirque active prior to McConnell Glaciation
- Arête; degraded arête active prior to McConnell Glaciation
- Streamlined glacial bedforms: ice flow direction known, unknown
- Meltwater channel; large, small, ice-walled channel; arrow indicates flow direction
- Esker; flow direction defined, unknown
- End moraine
- Recessional moraine
- Ice-contact face in stratified drift (teeth on ice side)
- Ice limit
- Cryoplanation terrace
- Tor
- Vertebrate fossil locality
- Stratigraphic section