






# Fieldguide

## List of Figures



**Figure 1: Geological map of southwestern British Columbia (1:500,000) with field trip route and location of stops**

**Figure 2A: Morphogeological belts of the Canadian Cordillera.**

**Figure 2B: Simplified metamorphic map of the Canadian Cordillera; this map displays the geologic character of the belts in Figure 2A; two belts of granitic and high-grade metamorphic rocks separates three tracts of little metamorphosed and subgreenschist facies rocks.**

**Figure 3: Simplified terrane map of the Canadian Cordillera;**

**Figure 4: Distribution of ages of Mesozoic and Tertiary granite rocks in the Coast Belt (simplified from Wheeler and McFeely, 1991).**

**Figure 5: Major Cretaceous and early Tertiary structures within and flanking the Coast Belt.**

**Figure 6: Terrane map of southwestern British Columbia (1:500,000) and accompanying crustal profile across the southern Coast and Intermontane Belts.**

**Figure 7: Evolving concepts of Coast Belt tectonics.**

**Figure 8: Correlation chart showing the stratigraphy and lithology of terranes and plutonic rocks in the southern Coast Belt and adjacent regions.**

**Figure 9: Map derived from 1:1 000 000 scale Tectonic Assemblage map of the southern Coast Mountains showing distribution of terranes, plutonic and metamorphic complexes, and overlap assemblages.**

**Figure 10: Amounts of uplift in southwestern British Columbia in the last 10 million years and present rates of uplift in millimetres per year.**

**Figure 11: Times and natures of terrane linkages, southwestern British Columbia**

# Fieldguide

## List of Figures








Figure 12: Summary tectonic evolution of southwestern British Columbia.

Figure 13: Parts of southeastern Coast and northwestern Cascade mountains, showing correlative (?) terranes and rock units that span the physiographic boundary between the two

Figure 14: Geological map of the Coast Belt thrust system near Harrison Lake, and index to U-Pb sample locations and interpreted dates cited in the text.

Figure 15: Geometry, kinematics and timing of the Coast Belt thrust system.

Figure 16: P-T diagram summarizing conditions of M1 and M2 metamorphic conditions in the southern Coast Belt

Figure 17: Some faults associated with the Fraser-Straight Creek fault system.

Figure 18: Terranes and Structures of the southeast Coast Mountains, Intermontane Belt and adjoining parts of the Cascade Ranges, used to demonstrate possible offsets on Fraser-Straight Creek Fault.

Figure 19: Southwestern margin of the Intermontane Belt, showing location of Stop 1-7.

Figure 20: Geologic map of the eastern coast Belt.

Figure 21: Structural and stratigraphic framework of the eastern Coast Belt.

Figure 22: Measured structural-stratigraphic sections of clastic successions overlying Bridge River and Cadwallader terranes in the southeastern Coast Belt.

Figure 23: Geology of the southern Shulaps Range, based on work of Schiarizza et al. (1990a,b), Coleman (1990), Journey and Northcote (1992), and previous mapping of Roddick and Hutchison (1973), Woodsworth (1977), and Monger (1989a).

# Fieldguide

## List of Figures







Figure 24: Fault systems of the southeastern Coast Belt.

Figure 25: Matrix-supported conglomerate of the Brew Group, exposed in the core of a southwest-vergent antiformal syncline, south of Mount Brew.

Figure 26: Thin-bedded phyllitic quartzite of the lower Brew Group exposed on the ridge southwest of Mount Brew.

Figure 27: Asymmetric fault zone fabrics in the lower plate of the Cayoosh Creek Fault

Figure 28: Geology of the Cayoosh Range (92J/8, J/9).

Figure 29: Pillowed greenstone of the Bridge River Complex exposed on ridge near headwaters of Downtown Creek.

Figure 30: Thin-bedded chert of the Bridge River Complex exposed at Twin Lakes, near headwaters of Melvin Creek.

Figure 31: Proposed correlations, terrane linkages and paleogeographic relationships for rocks of the southeastern Coast Mountains.

Figure 32: Asymmetric folds of foliation in footwall of southwest-vergent thrust faults cut by the Downtown Creek Fault. Photo taken along ridge crest at head of Downtown Creek.

Figure 33: View northeast from Duffey Lake Road along contact between the Bridge River Complex and conformably overlying Cayoosh assemblage

Figure 34: Ridge section of volcanic sandstone, siltstone and shale of the Cayoosh assemblage exposed in headwater region of Melvin Creek.

Figure 35: Schematic stratigraphic section of the Cayoosh assemblage, as compiled from measured structural sections in the Melvin and Mount Brew regions.

# Fieldguide

## List of Figures








Figure 36: Thinly laminated turbiditic siltstones of the lower Cayoosh assemblage.

Figure 37: Graded volcanoclastic sandstone and siltstone of the Cayoosh assemblage along ridge crest north of Duffey Lake.

Figure 38: Graded feldspathic wacke of the Cayoosh assemblage (Unit III) exposed near headwaters of Downtown Creek.

Figure 39: Thinly bedded phyllitic quartzite of the upper Cayoosh assemblage featuring complex intraformational (?) deformation structures, similar to those described by Monger (1991a) in the Settler Schist and Darrington Phyllite.

Figure 40: Asymmetric fault zone fabrics along the roof thrust of the Twin Lakes fault zone. Hanging wall rocks are sheared talc-carbonate "listwanite" schists of the Bralorne-East Liza Complex.

Figure 41: Staurolite-garnet-andalusite schists of the Chism Creek Schist exposed along ridge north of Duffey Lake.

Figure 42: Thinly laminated phyllitic quartzite of the Chism Creek Schist. Minor structures record a polyphase history of folding and fabric development exposed along ridge crest north of Duffey Lake.

Figure 43: Lithoprobe seismic reflection data and interpretations of Lines 14 and 17 along the Carpenter Lake-Hurley River Road, showing integration of geological and geophysical data. Interpretation by Journey in Monger et al. (1990).

Figure 44: Lithoprobe seismic reflection data and interpretations of Line 13 along the Duffey Lake Road, showing integration of geological and geophysical data. Interpretation by Journey in Monger et al. (1990).

Figure 45: Aerial view to the southeast, showing Lillooet Lake along strike of the Coast Belt thrust belt imbricate zone, and the boundary between eastern and western coast belts.

# Fieldguide

## List of Figures







Figure 46: The four "tracts" of the southwestern Coast Mountains, with location of stops 2-9 to 2-11.

Figure 47: Lithoprobe seismic reflection data and interpretations of Line15 illustrating high amplitude reflections beneath the Mount Cayley volcanic complex.

Figure 48: Sigmoidal structure in foliated amphibolite indicating movement to southwest (top to left). Corrie Ridge, 6.5 km east-north-east of the Black Tusk, Garibaldi Park.

Figure 49: Northeast-vergent, southwest -dipping mylonite/sheared granite rock; From mid-Cretaceous granite exposed near summit of Mount Roderick

Figure 50: Strike-normal view of pseudotachylite (ultramylonite) dyke in quartz diorite; these dykes dip northeast and have southwest-vergence; From mid-Cretaceous granite exposed near summit of Mount Roderick.