

**LEGEND**

Concentration of cobalt, less than 7 ppm  
in stream sediments ..... in spring sediments ..... 1

Concentration of cobalt, 7 ppm to 15 ppm  
in stream sediments ..... in spring sediments ..... 2

Concentration of cobalt, 15 to 30 ppm  
in stream sediments ..... in spring sediments ..... 3

Concentration of cobalt, 30 to 50 ppm  
in stream sediments ..... in spring sediments ..... 4

Concentration of cobalt, 50 ppm or greater  
in stream sediments ..... in spring sediments ..... 5

Location of known veins ..... 6

Mineral occurrence ..... 7

Mineral deposit ..... 8

**Mineral Symbols**

Arsenic ..... As Silver ..... Ag  
Antimony ..... Sb Tungsten (lode) ..... W  
Copper ..... Cu Tungsten (placer) ..... W(P)  
Gold (lode) ..... Au Tin (lode) ..... Sn  
Gold (placer) ..... Au(P) Tin (placer) ..... Sn(P)  
Lead ..... Pb Zinc ..... Zn  
Molybdenum ..... Mo

- INDEX TO MINES AND PROSPECTS**
- |   |                             |                          |
|---|-----------------------------|--------------------------|
| 1. Elsa                                 | 23. Sadio-Friendship        | 46. No. 1                |
| 2. Dixie                                | 24. Ladie                   | 47. Gambler              |
| 3. Coral and Wigwam                     | 25. Bellekono               | 48. Main fault and Nabob |
| 4. Arctic and Mastiff                   | 26. Mount Keno (Hogan vein) | 49. Lake View            |
| 5. Ruby                                 | 27. Ankeno                  | 50. Nabob No. 2          |
| 6. No Cash                              | 28. Mount Keno (Raner vein) | 51. Helen Fraction       |
| 7. Betty                                | 29. Dorothy                 | 52. Gold Hill No. 2      |
| 8. Cream                                | 30. Kijo                    | 53. Ladie Fraction       |
| 9. No Cash                              | 31. Croesus No. 1           | 54. Danam                |
| 10. Calumet                             | 32. Blank Cap and Shepherd  | 55. Silver Basin         |
| 11. Dragon (UN)                         | 33. Lucky Queen             | 56. Gold Queen           |
| 12. Forno                               | 34. Lake                    | 57. Carbon               |
| 13. Galikeno (McLeod vein)              | 35. Vanguard                | 58. Alice                |
| 13a. Galikeno (Sims and Sugiyama veins) | 36. Apex                    | 59. Carbon               |
| 14. Eagle                               | 37. Shamrock                | 60. Divide               |
| 15. Fisher Creek                        | 38. Highlander              | 61. Devon                |
| 16. Bluebird                            | 39. Cub and Beany           | 62. Faith                |
| 17. Tin Can                             | 40. Stone                   | 63. Silver King          |
| 18. Rio                                 | 41. Homestake               | 64. Gerlitsky            |
| 19. Duncan Creek                        | 42. No. 6                   | 65. Shungite             |
| 20. Moth                                | 43. Porcupine-Kimnan        | 66. Lookout              |
| 21. Onak                                | 44. Comstock                | 67. Rex                  |
| 22. Klondyke-Keno                       | 45. No. 9                   | 68. Peas Silver          |

Field work by C. F. Gleeson, W. M. Tupper, A. Suparnan, K. Demai, M. Shalqilah, J. A. Colwell, J. R. Deighton, C. H. Yurehak, J. K. Worth, H. R. James, A. G. Troup, G. Wasi, L. Hogg, and F. R. Campbell

Analyses by C. C. Durham

Compilation and text by C. F. Gleeson

Geological cartography by the Geological Survey of Canada, 1967

- Roads, all weather .....  
Other roads .....  
Trail .....  
Intermittent lake and stream .....  
Horizontal control point .....  
Elevation in feet above mean sea-level ..... 2000



Base-map cartography by the Geological Survey of Canada, 1966 from maps published by the Surveys and Mapping Branch and by the Army Survey Establishment, R. C. E.

Approximate magnetic declination, 34° 45' East, decreasing 4.2' annually

Published, 1968  
Copies of this map may be obtained from the Director, Geological Survey of Canada, Ottawa



MAP 54-1965  
COBALT CONTENT OF STREAM AND SPRING SEDIMENTS  
KENO HILL AREA  
YUKON TERRITORY



**Introduction**

The reconnaissance geochemical survey of Keno Hill area, Yukon Territory was started and completed in the summer of 1964. The creeks not accessible by roads were reached by helicopter. An attempt was made to maintain a sample interval of 1,000 feet along all rivers, creeks, and their tributaries. The data on this map are based on 5,500 samples of stream sediment collected from the channels of the streams and on the sediments and precipitates in the vicinity of springs from an area of approximately 1,500 square miles. Where possible the active channel was sampled; however as work progressed it was found that moss on the creek banks below the water line had trapped considerable amounts of fine sediment suitable for sampling. The wet sediments and waters were analyzed at the sample site for cold citrate-soluble heavy metals. The results of this work have been published in a series of 14 preliminary maps (Gleeson, et al., 1965). Field observations on the character of the stream, composition of the sediment, pH and temperature of the water, and rock types in the vicinity of the sample station were entered in code on special geochemical field cards. Subsequently, this information was punched on cards for electronic data processing.

The wet sediment was dried in the field at a temperature of about 60°C and sieved through an 80 mesh stainless steel screen. The sieved samples were shipped to Ottawa where they were ground to minus 100 mesh in a ceramic ball mill.

**Analysis**  
Cobalt was analyzed spectrographically by total energy D.C. arc semi-quantitative method using a Jarrell-Ash optical spectrograph with a 1.5 metre grating. A 10 milligram sample of ground stream sediment was mixed with 20 milligrams of graphite, packed into a carbon electrode, and capped with a 20 milligram buffer mixture of calcium carbonate and graphite. The loaded electrode was preheated at 450°C to oxidize the organic matter in the sample and thus allow the arc to proceed smoothly without loss of material from the electrode cavity. The electrode was then removed from the furnace after 45 minutes and cooled. Two drops of a saturated solution of magnesium nitrate in absolute ethyl alcohol were added in order to promote the smooth burning of the sample. The electrode was placed under an infrared lamp for at least five minutes to evaporate the alcohol. The samples were arced at 15 amps, and the spectra recorded on 35 mm Kodak Spectrum Analysis Film Number 1. The unknown spectra were then compared with a synthetically prepared series of spectra; the limit of detectability for cobalt was 7 ppm.

**General Geology**  
The regional geology has been described by Bostock (1947, 1954), and Green and Roddick (1962). More detailed geological studies have been made by Kindie (1962), McTaggart (1969), Poole (1965), and Green (1967, 1959). The geology, geochemistry, and origin of the mineral deposits in Keno Hill and Dalhousie areas have been described by Boyle (1965). Reports by Aho (1964) and Cockfield (1922) provide further information on mineral deposits of the area. The map-area is underlain by a series of metamorphosed sedimentary rocks, mainly quartzites, phyllites, slates, chlorites, sericitic and granitic schists, also gneiss and minor limestone. The age of these rocks is uncertain and appears to range from Precambrian to Mesozoic (Poole, 1965; Tempelman-Kluit, 1966). A dolomitic and limestone unit outcrops in the northeast part of the area. Fossils from these rocks range in age from Late Cambrian to Late Silurian or Early Devonian (Green and Roddick, 1962).

Mafic igneous sills and lenses now altered to greenstones are inter-layered with the metamorphosed sediments. Quartz-feldspar porphyry sills and lamprophyre dykes are present locally. Granitic stocks outcrop in the metamorphosed sediments east and north of Mayo Lake, northwest of Hanson Lake, south of Dalhousie and in the vicinity of Mount Halidane. Sill zones containing schistellite, occur in the vicinity of some of the granitic masses particularly around Dalhousie, Mount Halidane and east of Mayo Lake. Most of the lead-zinc-silver ore deposits in the Keno - Galena Hills area occur along northeasterly striking vein faults in thick-bedded quartzite and occasionally in greenstone (Boyle, 1965). In the Dalhousie area quartz arsenopyrite-gold veins with a general northeast strike are present near the contacts of the granitic stocks. Also easterly striking vein faults are mineralized with siderite, jasperite, boulangerite, pyrite, arsenopyrite, galena, tetrahedrite, and chalcopyrite. Two cassiterite-tourmaline veins occur on the right limit of Dalhousie Gulch near its mouth (Boyle, 1965; Poole, 1965). Also northerly striking lead-zinc-silver veins are present in Davidson Range (Cockfield, 1922; Aho, 1964). Fluorite has been recovered from Dalhousie Gulch, Haggart Creek, and Danam Creek since 1898. The area has undergone several stages of glaciation. Thick glacial deposits occupy the major valleys and hill slopes below an elevation of 3,000 feet. Permafrost is present throughout the area.

**Results**  
Statistical studies using electronic computation have yet to be completed and until they are adequate assessment of the results is difficult. However, cumulative distribution curves have been constructed from the information supplied by the computer. The curve for cobalt is illustrated on this map. A break in the slope of the curve suggests two distributions both distributed log-normally. Values for cobalt range from less than 7 ppm to 1000 ppm. For this map the samples have been grouped as follows: less than 7 ppm, 7 to 15 ppm, 15 to 30 ppm, and greater than 30 ppm. The majority of the high cobalt values are coincident with high nickel values (map 54-1965) and as in the case of nickel they are thought to be related to greenstones which are ultramafic in composition. There does not seem to be any correlation between cobalt in stream and spring sediments and the known silver veins of the area. Further follow up field work is warranted to explain the distribution of cobalt in the stream and spring sediments of the Keno Hill area.

Aho, A. E.: Mineral potential of the Mayo district, Western Miner, vol. 37, No. 10, pp. 90-98 (1964).  
Bostock, H. S.: Mayo, Yukon Territory; Geol. Surv. Can., Map 890A (1947).  
Cockfield, W. E.: McQuesten, Yukon Territory; Geol. Surv. Can., Map 1143A (1922).  
Boyle, R. W.: Geology, geochemistry, and origin of the lead-zinc-silver deposits of Keno Hill - Galena Hill area, Yukon Territory; Geol. Surv. Can., Bull. 111 (1965).  
Cockfield, W. E.: Silver-lead deposits of Davidson Mountains, Mayo District, Yukon Territory; Geol. Surv. Can., Summ. Rept. 1921, pt. A, pp. 1A-6A (1925).  
Gleeson, C. F., et al.: Heavy metal content of stream and spring sediments; Heavy metal content of stream and spring waters, Keno Hill area, Yukon Territory; Geol. Surv. Can., Maps 15-1964 to 21-1964 (1965).  
Green, L. H.: Mayo Lake, Yukon Territory; Geol. Surv. Can., Map 5-1956 (1957).  
Green, L. H.: McQuesten Lake and Scougale Creek map-areas, Yukon Territory; Geol. Surv. Can., Paper 58-4 (1958).  
Green, L. H., and Roddick, J. A.: Dawson, Larson Creek, Nash Creek map-areas, Yukon Territory; Geol. Surv. Can., Paper 62-7 (1962).  
Kindie, E. D.: Keno Hill, Yukon Territory; Geol. Surv. Can., Map 1102A (1962).  
McTaggart, R. C.: The geology of Keno and Galena Hills, Yukon Territory; Geol. Surv. Can., Bull. 58 (1960).  
Poole, W. H.: Report of activities: field, 1964; Geol. Surv. Can., Paper 65-1, pp. 32-34 (1965).  
Tempelman-Kluit, D.: Report of activities, May to October, 1965; Geol. Surv. Can., Paper 66-1, pp. 44-49 (1966).

