

The regional geochemical trend map displayed above utilized a moving weighted average using an inverse distance function (1/d²) to filter out minor irregularities and emphasize broad-scale regional features. Single point anomalies may be suppressed or eliminated, however, geological units which are chemically enriched, or large metallic deposits undergoing weathering would be expected to produce identifiable anomalies

Geological Survey of Canada
 Resource Geophysics and Geochemistry Division
 CONTRACTORS
 Sample collection by Rogers Exploration Services Ltd., Whitehorse
 Sample preparation by Golder Associates, Ottawa
 Gold analysis by Chemex Labs Limited, Vancouver, B.C.
 Sediment chemical analyses by Barringer Magenta Ltd., Rexdale, Ontario
 Water chemical analyses by Barringer Magenta Laboratories (Alberta) Ltd., Calgary

This map forms one of a series of maps released by the Geological Survey of Canada, Open Files 1217 to 1220. Each Open File consists of maps of various geochemical variables: 21 for stream sediment, 3 for stream water and 1 sample site location

Copies of map material and listings of field observations and analytical data, from which the material was prepared, may be available at users expense by application to:

K.G. Campbell Corporation
 880 Wellington St.
 Bay 238
 Ottawa, Ontario
 K1R 6K7



The data are also available in digital form. For further information please contact:

The Director
 Computer Science Centre
 Department of Energy, Mines and Resources
 Ottawa, Ontario
 K1A 0E4

- Glaciolacustrine deposits
- Undivided surficial deposits; alluvium, glacial till and moraine, outwash and ice contact deposits, volcanic ash, loess, colluvium
- Bedrock exposures; includes discontinuous veneer of undivided glacial drift, local alpine glaciation features

- SYMBOLS
- Surficial deposit boundary
 - Meltwater channels, outwash deposits, indicating direction of flow
 - Glaciation lineation parallel to ice flow direction, includes fluting, crag and tail, roches moutonnées and drumlinoid forms, direction of flow known, unknown
 - Drumlinoid form, direction of movement inferred, not inferred
 - Glacial striae, direction inferred
 - Esker and/or kame complex
 - Boulder train, direction of movement

Sources of information:
 Hughes, D.L., Campbell, R.B., Muller, J.E., and Wheeler, J.O. (1968) Glacial Map of Yukon Territory, Geological Survey of Canada, Map 6-1968, (1:1 000 000 scale) to accompany GSC Paper 68-34
 Prest, V.K., Grant, D.R., and Rampton, V.N. (1967) Glacial Map of Canada, Geological Survey of Canada (1:5 000 000 scale)
 Wheeler, J.O. (1960) Geology - WHITEHORSE, Yukon Territory, Geological Survey of Canada, Map 1095A (1:253 440 scale)




pH in water
 GSC OPEN FILE 1218
 REGIONAL GEOCHEMICAL RECONNAISSANCE MAP 83-1985
 CANADA-YUKON
 MINERAL DEVELOPMENT AGREEMENT (1984-89)
 STREAM SEDIMENT AND WATER GEOCHEMICAL SURVEY
 SOUTHERN YUKON TERRITORY, 1985

Elevation in feet above mean sea level

Mean magnetic declination 1986, 29°10' East, decreasing 14.2' annually. Readings vary from 29°06' E in the SE corner to 22°29' E in the NW corner of the map area

Scale 1:250 000
 Universal Transverse Mercator Projection
 © Crown Copyrights reserved

Base map at the same scale published by the Surveys and Mapping Branch in 1979. Streams were revised by the Geological Survey of Canada for this edition



LEGEND

QUATERNARY	33 RMC 64*	MILES CANYON: Basalt
	32 QS 64	Glacial and surficial deposits
TERTIARY	31 TOM 62	Quartz monzonite, granodiorite
LATE TERTIARY	30 LTG 62	Rhyolite porphyry, granite, granodiorite
PLIOCENE	29 PV 62	Basalt
Eocene	28 EMN 59	Acid to intermediate tuff, breccia
MOUNT NANSEN GROUP	27 ESK 59	Andesite, basalt, breccia
SKUKUM GROUP	26 ESL 59	Rhyolite, trachyte
SLOKO GROUP	25 KTG 56	Granite, quartz monzonite
CRETACEOUS AND TERTIARY	24 KTGD 56	Granodiorite, quartz diorite
	23 KTQD 56	Tonalite
CRETACEOUS	22 KY 52	Syenite, monzonite
	21 KQM 52	Quartz monzonite, granodiorite; CASSIAR quartz monzonite, alaskite
	20 KGD 52	Granodiorite
	19 KV 52	Basalt, andesite, quartz dacite
JURASSIC AND CRETACEOUS	18 JKDI 51	Diorite, hornblende diorite
JURASSIC	17 JKT 51	TANTALUS: Conglomerate, siltstone, arkose, coal
	16 JL 47	Greywacke, arkose, conglomerate
TRIASSIC AND JURASSIC	15 TJS 46	Argillite, sandstone, siltstone
TRIASSIC	14 TGMN 42	Foliated hornblende granodiorite, quartz
UPPER TRIASSIC	13 UTLW 45	Greywacke, argillite, conglomerate
	12 UTC 45	Limestone
	11 UTLV 45	Andesite, basalt
MESOZOIC UNDIVIDED	10 MGD 41	Granodiorite, quartz monzonite
	9 MGDN 41	Foliated hornblende granodiorite, quartz monzonite
	8 MV 41	Andesite, basalt, tuff
PALEOZOIC UNDIVIDED	7 PGDN 09	PELLY GNEISS: Foliated to gneissic granodiorite
CARBONIFEROUS AND PERMIAN	6 CPH 35	HORSEFEED: Limestone
	5 CPK 35	KEDAHDA: Chert, argillite
	4 CPV 35	Andesite, basalt, chert, tuff
	3 CPUB 35	Serpentine, diorite, pyroxenite, peridotite
HADRYNIAN AND CAMBRIAN	2 HCSN 08	Schist, gneiss, quartzite
HADRYNIAN	1 HC 07	Crystalline limestone

*A mnemonic code assigned to rock types and recorded as part of field observations

Geological boundary

Fault

No analytical result

Geological base and legend are derived from: Map 1398A, MACMILLAN RIVER, YUKON - DISTRICT OF MACKENZIE - ALASKA, NTS SHEET 105, 115. Compiled by H. Gabrielsen, D.J. Tempelman-Kluit, S.L. Blusson and R.B. Campbell, Geological Survey of Canada, Energy, Mines and Resources Canada, 1980. 1:1 000 000 scale