GEOLOGICAL SURVEY OF CANADA OPEN FILE 1686

MINERAL AND ENERGY RESOURCE ASSESSMENT OF THE TLOGOTSHO PLATEAU, NAHANNI KARST, RAGGED RANGES AND ADJACENT AREAS UNDER CONSIDERATION FOR EXPANSION OF NAHANNI NATIONAL PARK RESERVE, NORTHWEST TERRITORIES

Edited by C.W. Jefferson¹ and W.A. Spirito¹

Contributing Authors:

C.W. Jefferson¹, K.G. Osadetz², W.A. Spirito¹, S.M. Hamilton³, Z. Chen², D.W. Morrow², F.A. Michel⁴, D. Power-Fardy⁵ and D. Paré⁶

¹ Geological Survey of Canada – Mineral Resources Division, 601 Booth St., Ottawa ON K1A 0E8

² Geological Survey of Canada – Calgary, 3303 –33rd St. N.W., Calgary, AB T2L 2A7

³ Ontario Geological Survey, Ministry of Northern Development and Mines, Willet Green Miller Centre, 933 Ramsey Lake Road, 7th Floor, Sudbury ON P3E 6B5

⁴ Department of Earth Sciences, 2240 Herzberg Laboratories, 1125 Colonel By DriveCarleton University, Ottawa, ON K1S 5B6

⁵ 28 Tanglewood Dr., Ottawa, ON K2H 6P3

⁶ Consorminex Inc., 11-1695 Atmec, Gatineau, PQ, J8P 7G7

Jefferson, C.W., Spirito, W.A. (eds.) 2003. Non-renewable mineral and energy resource potential of proposed extensions to Nahanni National Park Reserve, Northern Cordillera, N.W.T. Geological Survey of Canada, Open File 1576, CD-ROM.

Table of Contents

	page
Table of Contents	iii
1 Executive Summary	1 1
1. Executive Summary	1-1 Mataau
1.1. Minerai Resource Assessment of the Raggea Ranges and Nananni Rarst – Hogoisno F Study Areas, South Nahanni Piyar Pasion	1 2
Broken Domein	1-2
Bioken Domain	1-0
Homonald Domain	1-0
MacDonald Domain	1-0
Matteen Demoin	1-0
Mattson Domain	1-/
Memeur Domain	1-/
<u>Ragged Domain</u>	1-/
Somore Domain.	1-/
1.2. On and Gas Resource Assessment of the Ragged Ranges and Nananni Karst – Hogois,	no 1 7
Plateau Study Areas, South Nananni Kiver Kegion	1-/
1.5. Caveats	1-9
2. Introduction	2-1
2.1 MERA Terms of Reference	2-2
2.1. Millian Fernis of Regenered Study Areas and Features of National Park Interest	2-4
2.2. Ocographic Setting, million Study in case and 1 earliers of Hattonal 1 and interest million 2.3 History and Method of NNPR MERA Studies	2-6
2.5. Thistory and interior of this Assessment	2-7
2.7. Confidence in this Assessment.	2-12
2.5. Organization of this Report and Responsibilities of the Mathors	2-12
2.0. Acknowledgmenis	2-12
3. Geological Setting	3-1
3.1. Rock Packages	3-5
3.1.1. Basinal shale-dominated and transitional Neoproterozoic strata	3-6
3.1.2. Platformal and transitional carbonate - clastic Neoproterozoic strata	3-7
3.1.3. Basinal shale-dominated & transitional Early Paleozoic strata	3-7
3.1.4. Platformal carbonate and clastic Early Paleozoic strata	3-9
3.1.5. Basinal shale-dominated and transitional Late Devonian and younger strata.	3-9
3.1.6. Platformal clastic-dominated Late Devonian and vounger strata	3-10
3.1.7. Cretaceous intrusions	3-10
3.1.8. Quaternary - unconsolidated glacial and alluvial deposits	3-11
3.2. Metamorphism	3-14
3.3. Tectonism and Paleogeography	3-14
3.4 Definition of resource assessment domains for minerals and coal	3-15
3.5. Geological features of public interest	3-18
4. Regional Surficial Geochemistry	4-1
4.1. Introduction and Background	4-1
4.1.1. History of geochemical studies in Nahanni region	4-1
4.1.2. Principles of stream sediment sampling	4-2

Stream silts	4-3
Heavy mineral concentrates (HMCs)	4-3
4.2. Sample Collection, Preparation and Analysis	4-5
4.2.1. Sample collection for the South Nahanni River mineral resource assessment	4-5
Collection of silts for silt and HMC analysis	4-9
Collection of pebbly sands for HMC analysis	. 4-10
Collection of bulk samples	. 4-10
4.2.2. Sample preparation in the laboratory	. 4-10
Preparation of silt samples	. 4-10
Preparation of HMCs from silt samples	. 4-11
Preparation of HMCs from pebbly sands	. 4-13
Preparation of bulk samples	. 4-13
4.2.3. Analytical methods	. 4-14
Precision and accuracy	4-14
The risk of contamination	. 4-14
Treatment of gold grains	4-15
4.3. Statistical analysis	4-15
4.3.1. Statistical methods for South Nahanni River data	4-15
4.3.2 Geochemical data below detection limit	4-16
4 3 3 Maximum likelihood estimation (MLE)	4-16
How are MLEs derived?	4-17
Maximum likelihood (ML) estimation of South Nahanni River data	4-18
4.3.4 Elements used in statistical analysis	4-18
4.3.5 Tests for normality	4-18
4.3.6. How sample mean and standard deviation can be used	4-18
4.3.7 Calculation of percentile values to classify the original data	4-19
4.3.8 Regression analysis	4-28
A linear first-order regression model	4-28
Application to South Nahanni River stream sediment geochemical data	4-29
The significance of regression results	4-32
Predicted background values for 100% of each rock type	4-32
4.4 Spatial analysis	4-33
4.4.1 Geographic information systems (GIS)	4-33
4 4 2 Catchment hasin analysis	4-34
4 4 3 Removing the effect of rock type	4-35
4.5 Gold Results	4-36
4.5.1 HMCs from Silt Samples	4-36
4.5.2 Placer Gold from nebbly sand samples	4-36
4 5 3 Placer Gold from Silts	4-37
4 5 4 Heavy Mineral Point Count	4-39
4.5.5 Sources of gold	4-41
4.6 Discussion of surficial geochemistry results	4-42
4.6.1 Mineral notential from geochemistry - Ragged Ranges	4-43
4.6.2 Mineral potential from geochemistry - Nahanni Karst / Tlogotsho Plateau	4-56
4.7 Conclusions from stream sediment geochemistry studies in South Nahanni River region	4-67
4.7.1 Stream silts vs. heavy mineral concentrates	4-67
4.7.2. Statistical techniques	4-75
4.7.3 Mineral notential from stream sediment geochemistry	4-75
1.7.5. Minerai porennai from su cam seanneni zeoenemisir y	. 175

5. Spring Water Geochemistry and Hydrogeology	5-1
5.1. Introduction	5-1
5.1.1. Background - spring sampling as an assessment tool	5-1
5.1.2. Previous hydrogeologic studies in the Nahanni region	5-1
5.1.3. Hydrogeologic objectives and scope	5-3
5.2. Hvdrogeologic setting	5-4
5.2.1. Controls on groundwater recharge, flow and discharge	5-4
5.2.2. Hvdrochemical influences	5-5
5.3. Hvdrogeology methods and data collected	5-5
5.4. Classification of springs	5-6
5.4.1. Introduction to classification methods	5-6
5.4.2. Field classification	5-7
5.4.3. Kleiner-Hartigan Trees - grouping by hydrogeochemical pattern recognition.	5-7
5.4.4. Cluster analysis	5-9
5.4.5. Discussion of individual classifications	5-14
5.4.6. Comparisons among groups	5-15
5.5. Final spring groups, hydrogeochemical processes and resource assessment in the study	areas
	5-15
5.5.1. Introduction	5-15
5.5.2. Group A – Kraus Hot Springs at Clausen Creek	5-16
5.5.3. Group B - Acidic, high $SO_4^{2^2}$ springs	5-19
Subgroup B1 - Metalliferous springs in the Meilleur River area	5-20
Subgroup B2a - Metalliferous springs in Ragged Ranges study area	5-22
Subgroup B2b - Metalliferous springs discharging from the Mattson Format	tion
	5-24
5.5.4. Group C - High pCO_2 springs in the Flat River Valley	5-25
5.5.5. Group D - High discharge TDS springs in carbonate terrane	5-28
Subgroup D1a - Fault localized thermal springs	5-29
Subgroup D1b - Other D1 springs	5-31
Subgroup D2 - Moderate TDS springs in carbonate terrain	5-32
5.5.6. Group E - Thermal springs related to quartz monzonite	5-32
5.5.7. Group F - Low TDS spring and surface waters	5-34
5.6. Hydrogeochemical conclusions	5-35
5.6.1. Classification of springs	5-35
5.6.2. Details of final spring water classification (see also Table 5.5)	5-36
5.6.3. Anomalous springs and areas of special interest	5-36
6. Mineral Occurrences and Geochemical Anomalies	6-1
6.1. Mineral showings of proposed extensions to NNPR	6-1
6.2. Gemstones – new to the region	6-1
6.3. Geochemical Anomalies	6-2
6.4. Ragged Ranges Metallotects	6-9
6.5. Nahanni Karst - Tlogotsho Plateau: Placer Gold	6-16
6.6. Nahanni Karst - Tlogotsho Plateau: Sedex Zinc-Lead, Meilleur River Area	6-16
6.7. Nahanni Karst - Tlogotsho Plateau: other geochemical associations	6-17
6.8. Summary of Metallotects, Proposed Expansions to Nahanni National Park Reserve	6-18

7. Mineral Resource Assessment of the Tlogotsho Plateau, Nahanni Karst, Ragged Ranges and Adjacent Areas Under Consideration for Expansion of Nahanni National Park Reserve, N.W.T.

7.1. Resource Assessments - Past and Present	
7.2. Qualitative Assessments of Mineral Potential by Deposit Models in Domains	
7.2.1 Placer gold	
7.2.2 Coal	
7.2.3 Sedex lead-zinc-silver	
7.2.4 Sedimentary nickel sulphides	7-3
7.2.5 Mississippi Valley-type lead-zinc	
7.2.6 Lode gold	
7.2.7 Vein silver-lead-zinc	
7.2.8 Skarn tungsten	
7.2.9 Skarn lead-zinc-silver	7-7
7.2.10 Emerald associated with bituminous sedimentary rocks	
7.2.11 Emeralds associated with skarns	
7.2.12 Other Gems	7-9
7.2.13 Granitic pegmatite-hosted rare metals	7-10
7.3. Mineral deposit types of low to very low potential in the S Nahanni R. region	7-10
7.4. Mineral deposit types of highly uncertain potential in the South Nahanni River region	on 7 - 11

8. Petroleum Resource Assessment of the Tlogotsho Plateau, Nahanni Karst, and Adjacent Areas Under Consideration for Expansion of Nahanni National Park Reserve, Northern Cordillera and District of Mackenzie

ict of Mackenzie	8-1
8 1. Abstract	8-1
8.2. Hvdrocarbon assessment background	
8.3 Geology of the Liard hydrocarbon assessment domain	
8.3.1 Neoproterozoic successions	
8.3.2 Cambrian, Ordovician and Silurian	
8.3.3 Devonian	
8.3.4 Carboniferous	
8.3.5 Permian and Triassic	
8.3.6 Cretaceous	
8.3.7 Structure	
8.4 Petroleum Systems	
8.5 Exploration History	
8.6 Petroleum Occurrences	
8.6.1 Discovered reserves	
8.6.2 Petroleum shows	
8.6.3 Previous petroleum potential assessments	
8.7 Petroleum Resource Assessment	
8.7.1 Method	
8.7.2 Petroleum systems analyses	
8.7.3 Play-based assessments of petroleum potential	

8.7.4 The Geo-anchored method	8-22
8.7.5 Historical vindication of sequential sampling assessment methods	8-22
8.7.6 Reservoir volume methods	8-23
8.7.7 Play definition	8-26
8.7.8 Sequential sampling method (geo-anchored discovery history analysis)	8-26
8.7.9 Play reservoir volume and multivariate analysis calculations	8-27
8.8 Discussion	8-31
8.9 Implications and Recommendations for NNPR and Proposed Extensions	8-34
8.9.1. Specific Areas Outside and On the Margins of the Proposed Park Reserve	
Extensions with High Natural Gas Potential	8-34
8.9.2. Factors relating to lower hydrocarbon potential of Nahanni Karst –	
Tlogotsho Plateau Study Areas	8-34
8.9.3 Access for scientific and engineering studies in the park reserve	8-36
8.10 Conclusions	8-36

9. References	
Appendix 1 - Mineral Occurrence and Deposit Data	A1-1
A-1.1 Introduction to Appendix – Mineral Occurrences of the South Nahanni Ri	ver RegionA1-1
Figure A-1(i). Mineral occurrences of the South Nahanni River region	A1-2
Figure A-1(ii). Labelled showings in the Howards Pass – Lened area	A1-3
Figure A-1(iii). Labelled showings in the Tungsten area	A1-3
Figure A-1(iv) Labelled showings in the Prairie Creek – Nahanni Karst area	A1-3
Table A-1.1. Partial listing of mineral occurrences in the South Nahanni River	area
	nOcc.xls and A1-4
Table A-1.2 Nahanni mineral occurrences locations	ni MinOccLoc.xls
Selected Mineral Deposit Synopses from the South Nahanni River Region	A1-19
A-1.2 Cantung Tungsten Deposit Synopsis	A1-19
A-1.3 Howards Pass Lead-Zinc Deposit Synopsis	A1-20
A-1.4 Lened Tungsten and Emeralds Deposit Synopsis	A1-20
A-1.5 Little Nahanni Pegmatite Group Lithium Synopsis	A1-21
A-1.6 Mawer Carbonate-Hosted Lead Zinc	A1-22
A-1.7 O'Grady Aplite-Pegmatite Gemstones	A1-23
A-1.8 Prairie Creek Vein-Carbonate-Shale-Hosted Silver-Lead-Zinc	A1-24
A-1.9 Selena Creek Placer Gold Synopsis	A1-25
A-1.10 Vulcan Sedex Lead-Zinc-Barite Synopsis	A1-26
Appendix 2 - Geochemistry of bedrock samples	A2-1
A-2.1: List of Samples, Coordinates and Field Data (descriptions etc.)	A2-1
Table A-2.1(i): Listing in chronologic order by groups, of 110 rock samples	. Table A-2.1-i.xls
Table A-2.1(ii). Analytical results of 100 representative rock samples from	
South Nahanni River region (includes major and trace elements)	Table A-2.1-ii.xls

Appendix 3 – Stream Sediment Geochemical Data and Statistical Notes

Appendix 3.1 Analytical Results for Stream Silts	Appendi	\mathbf{x} 3.1.xls
Appendix 3.2 Gold Properties and Analytical Result	s for HMCs Derived from Silts	A3.2-1
Table A-3.2(i) Samples for MI separation, mour	nting and point counting	A3.2-1
Table A-3.2(ii) Descriptive terms for gold grain	S	A3.2-1
Table A-3.2(iii) Gold grains recuperated at sha	king table; <177µm fraction	A3.2-2
Table A-3.2(iv) List of mineral codes used in po	int counting heavy minerals	A3.2-3
Figure A-3.2(i) Histograms of 300-grain counts	of selected HMCs from stream pebbly sai	nds
yielding anomalous gold or other meta	ellotects of interest	A3.2-3
Table A-3.2(v) Heavy mineral analysis (% abun	dance), s.g. 3.3, 300 grain count	A3.2-4
Table A-3.2(vi) INAA results for heavy mineral	concentrates from silts, elements Na to Th	,
		d A3.2-7
Table A-3.2(vi) cont'd INAA results for heavy	mineral concentrates from silts, elements	La to U,
Mass		A3.2-11
Appendix 3.3 Analytical Results for HMCs from Pet	obly Sands Appendi	ix 3.3.xls
Annendiy 3.4 Analytical Results for Sample Duplica	tas Bulk Samples and Standards	
Appendix 5.4 Analytical Results for Sample Duplica	Annendi	ix 34 xls
A-3 4 1 Silt	Worksheet "A-3 4 1 Silts" in Appendi	x 3.4 xls
A-3 4 2 Heavy Mineral Concentrates	Worksheet "A-3 4 2 HMCs" in Appendi	x 3.4 xls
		.A 9. 1.A15
Appendix 3.5 Analytical Techniques and Laborator	ies	A3.5-1
A-3.5.1 Heavy Mineral Concentrates 1985. 198	6 and 1987	A3.5-1
Table A-3.5.1(i) Detection limits for INAA/AAS	analyses of HMCs	A3.5-1
A-3.5.2 Stream Silts 1985, 1986, 1987	· · · ·	A3.5-1
Table A-3.5.2(i) Detection limits for 1985 silt at	nalyses	A3.5-1
Table A-3.5.2(ii) Detection limits for 1986 silt a	inalyses	A3.5-2
Table A-3.5.2(iii) Detection limits for 1987 silt	analyses	A3.5-3
A-3.5.3. Preparation of HMCs from bulk stream	n silt samples	A3.5-3
Table A-3.5.3(i) Silt HMC-mag standards for II	VAA analysis	A3.5-5
Table A-3.5.3(ii) Sand HMC-mag standards for	· INAA analysis	A3.5-5
Figure A-3.5.3(i) Procedures used in preparing	; most HMCs from South Nahanni River si	lt
samples		A3.5-6
Figure A-3.5.3(ii) Preparation of standards for	INAA from 86-JPW-123-S	A3.5-6
Figure A-3.5.3(iii) Preparation of standards for	r INAA from 87-JPW-61-S	A3.5-6
Figure A-3.5.3(iv) Preparation of standards for	· INAA from 87-JPW-66-S	A3.5-7
Table A-3.5.3(iii) HMC processing results from	shaking table and methylene iodide	A3.5-8
Annondix 3.6 Estimation of Distribution Parameter	s from Data with Observations Polow I	Dotostion
I imit	s nom Data with Observations Delow I	A 2 6 1
Appendix 3.7 Input/Output for ML Estimation of M	lean and Variance	A3.7-1

Appendix 3.8 Good Estimators of α and β ²	A3.8-1
Appendix 3.9 ML Regression	A3.9-1
Appendix 3.10 Predicted Values for Elements Based on Basins Underlain by 100% of R	ock Type
	A3.10-1
A-3.10.1. Ragged Ranges	A3.10-1
A-3.10.2. Nahanni Karst-Tlogotsho Plateau	A3.10-16
Appendix 4 – Analytical Data, Background Information and Detailed Methodology for S Water Geochemistry and Hydrogeology.	pring
Appendix 4.1. Field Data and Classification of Springs	A4.1-1
A-4.1.1. Group P1 - Tufa springs	A4.1-1
A-4.1.2. Group P2 - Iron springs	A4.1-1
A-4.1.3. Group P3 - Hot springs in quartz monzonite terrane	A4.1-2
A-4.1.4. Group P4 - Clausen Creek hot springs	A4.1-2
A-4.1.5. Group P5 - High discharge cold springs	A4.1-2
A-4.1.6. Group P6 – Miscellaneous	A4.1-2
<i>Table A-4.1(i)</i>	A4.1-3
Appendix 4.2. Laboratory Data for Nahanni Spring Waters and Precipitates	A4.2-1
Table A-4.2(i) Analytical techniques and detection limits	A4.2-1
Table A-4.2(ii) Major ions in spring waters	A4.2-2
Table A-4.2(iii) Trace elements in spring waters	A4.2-5
Table A-4.2(iv) Isotopes in spring waters	A4.2-9
A-4.2.1. Analytical methods for stable isotopes in Nahanni water samples	A4.2-10
Table A-4.2(v) Major elements from selected precipitates from springs	A4.2-12
Table A-4.2(vi) Trace elements from selected precipitates from springs	A4.2-13
Appendix 4.3. Additional Notes on Kleiner-Hartigan Tree Classification	A4.3-1
A-4.3.1. Introduction to major and trace-element classification	A4.3-1
A-4.3.2. Kleiner-Hartigan methodology	A4.3-1
A-4.3.3. Results and discussion of Kleiner-Hartigan Trees groupings	A4.3-1
Appendix 4.4. Additional Notes on K-Means Cluster Analysis Classification	A4.4-1
A-4.4.1. Introduction to cluster analysis	A4.4-1
A-4.4.2. Data development and methodology	A4.4-1
Data input	A4.4-1
Data manipulation	A4.4-1
Figure A-4.4(i) Flow chart to show potential routes for inputting, manipulating and	
clustering the data	A4.4-2
A-4.4.3. Clustering method	A4.4-3
A-4.4.4. Discussion and results of cluster analysis	A4.4-3
Figure A-4.4(ii) Output of cluster analysis on samples from groups C7t, C8t and C9t	A4.4-4
Table A-4.4(i) Results of K-means cluster analysis	A4.4-5

Figure 1.2. Summary of resource assessment domains and mineral potential indicators of the	
Ragged Ranges study area	. 1-3
Figure 1.3. Summary of resource assessment domains and mineral potential indicators of the	
Nahanni Karst and the Tlogotsho Plateau study areas	. 1-4
Figure 1.4. Play area map showing the location of the three structures, Etanda Dome, Twisted	ĺ
Mountain Anticline and Mattson Anticline	. 1-8
Figure 2.1. Geographic, political and economic context of NNPR	. 2-1
Figure 2.2. Proposed interim land withdrawal, draft as of February 2003	. 2-3
Figure 3.1 Summary geology of the South Nahanni River region	. 3-2
Figure 3.2. Simplified geology, Ragged Ranges study area	. 3-4
Figure 3.3. Simplified geology of the Nahanni Karst - Tlogotsho Plateau study area	. 3-5
Figure 3.4. Typical Road River Group shales	. 3-9
Figure 3.5. Paleokarst solution collapse breccias	3-10
Figure 3.6 Fog occupying the South Nahanni River valley near the Vulcan property	3-12
Figure 3.7 Knob and kettle terrain on the north side of South Nahanni River	3-12
Figure 3.8 Glacier Lake here viewed looking east from Mount Harrison Smith	3-12
Figure 3.9 Ice-cored rock glacier facing I ened	3-13
Figure 3.10. Slumped alluvium north side of Nahanni River hank near Rabbitkettle River	3_13
Figure 3.11 Bedded fluvial sands and gravels	3_13
Figure 3.12a Polie #1 looking north full of water	3 13
Figure 3.12a. Polje #1 looking north, ampty of water	2 12
Figure 2.12. I offe #1 looking north, empty of water	3^{-13}
Figure 3.13. Lateral train note on north side of police #1	3 - 14 2 1 /
Figure 2.15. West flenk of Nehenni Putte looking East	2 15
Figure 3.16. Assessment domains of the South Nahanni River region	3 17
Figure 3.17. Virginia Falls (centre) are carved into the delemitic Suppleed Formation	3-17 2 10
Figure 2.18. Pabbitkettle tufe travertine springs	2 10
Figure 2.10. White Aster (86 IDH 082: Meilleur P. area)	2 20
Figure 3.19. White Aster (objf fi-062, Meineur K. area)	2 20
Figure 2.21 Winter LandSet game of the sector NNDD area	2 21
Figure 5.21. White Landsat scene of the eastern NNFK area	3-21
Figure 4.1a. Sample Location Map, Ragged Ranges Study Area	. 4-/
Figure 4.10. Sample Location Map, Nananin Kaist Study Area	. 4-0
Figure 4.2. Sleving Peddy Sand Samples for Heavy Mineral Concentrates	4-10
Figure 4.3a. Gold Grain from Brophy Creek, Ragged Ranges	4-15
Figure 4.50. Gold Grain from wretched Creek, Nananni Karst	4-15
Figure 4.4: Calchment basins for Ragged Ranges	4-20
Figure 4.5a. Ragged Ranges, Au in HMC using MLE – Raw Data	4-22
Figure 4.50. Ragged Ranges, Au in HMC using MLE – Predicted Data	4-22
Figure 4.0a. Ragged Ranges, Au in HMC using $SUB - Raw Data$	4-23
Figure 4.60. Ragged Ranges, Au in HMC using SUB – Predicted Data	4-23
Figure 4. /a. Ragged Ranges, W in HMC using MLE – Raw Data	4-24
Figure 4.7b. Ragged Ranges, W in HMC using MLE – Predicted Data	4-24
Figure 4.8a. Ragged Ranges, W in HMC using SUB – Raw Data	4-25
Figure 4.8b. Ragged Ranges, W in HMC using SUB – Predicted Data	4-25
Figure 4.9a. Ragged Ranges, Cr in HMC using MLE – Raw Data	4-26
Figure 4.9b. Ragged Ranges, Cr in HMC using MLE – Predicted Data	4-26
Figure 4.10a. Ragged Ranges, Cr in HMC using SUB – Raw Data	4-27
Figure 4.10b. Ragged Ranges, Cr in HMC using SUB – Predicted Data	4-27

Figure 4.11a. Nahanni Karst, Hf in silt using MLE – Raw Data	4-31
Figure 4.11b. Nahanni Karst, Hf in silt using MLE – Predicted Data	4-31
Figure 4.12: Digitized elements: streams, catchment basins, park boundary, study area and	sample
locations for Nahanni Karst/Tlogotsho Plateau	4-34
Figure 4.13: Catchment basin map for Nahanni Karst/Tlogotsho Plateau	4-35
Figure 4.14: Location of Gold Flakes from HMCs in Ragged Ranges and Nahanni Karst	4-37
Figure 4.15. Locations of catchment basins (numbered polygons outlined by solid lines) w	vhich
are anomalous in one or more elements in the eastern part of the South Nahanni R	iver
resource assessment region.	4-38
Figure 4.16a. Ragged Ranges, Au in silt using MLE – Raw Data	4-44
Figure 4.16b. Ragged Ranges, Au in silt using MLE – Predicted Data	4-44
Figure 4.17a. Ragged Ranges, As in HMC using MLE – Raw Data	4-45
Figure 4.17b. Ragged Ranges, As in HMC using MLE – Predicted Data	4-45
Figure 4.18a. Ragged Ranges, Cu in silt using MLE – Raw Data	4-46
Figure 4.18b. Ragged Ranges, Cu in silt using MLE – Predicted Data	4-46
Figure 4.19a. Ragged Ranges, Mo in HMC using MLE - Raw Data	4-47
Figure 4.19b. Ragged Ranges, Mo in HMC using MLE – Predicted Data	4-47
Figure 4.20a. Ragged Ranges, Ni in HMC using MLE – Raw Data	4-48
Figure 4.20b. Ragged Ranges, Ni in HMC using MLE – Predicted Data	4-48
Figure 4.21a. Ragged Ranges, Ce in HMC using MLE – Raw Data	4-49
Figure 4.21b. Ragged Ranges, Ce in HMC using MLE – Predicted Data	4-49
Figure 4.22a. Ragged Ranges, W in silt using MLE – Raw Data	4-50
Figure 4.22b. Ragged Ranges, W in silt using MLE – Predicted Data	4-50
Figure 4.23a. Ragged Ranges, Cd in silt using MLE – Raw Data	4-51
Figure 4.23b. Ragged Ranges, Cd in silt using MLE – Predicted Data	4-51
Figure 4.24a. Ragged Ranges, Ni in silt using MLE – Raw Data	4-52
Figure 4.24b. Ragged Ranges, Ni in silt using MLE – Predicted Data	4-52
Figure 4.25a. Ragged Ranges, Pb in silt using MLE – Raw Data	4-53
Figure 4.25b. Ragged Ranges, Pb in silt using MLE – Predicted Data	4-53
Figure 4.26a. Ragged Ranges, Zn in silt using MLE – Raw Data	4-54
Figure 4.26b. Ragged Ranges, Zn in silt using MLE – Predicted Data	4-54
Figure 4.27a. Ragged Ranges, Zn in HMC using MLE – Raw Data	4-55
Figure 4.27b. Ragged Ranges, Zn in HMC using MLE – Predicted Data	4-55
Figure 4.28a. NK/TP, Zn in HMC using MLE – Raw Data	4-56
Figure 4.28b. NK/TP, Zn in HMC using MLE – Predicted Data	4-56
Figure 4.29a. NK/TP, Sb in HMC using MLE – Raw Data	4-57
Figure 4.29b. NK/TP, Sb in HMC using MLE – Predicted Data	4-57
Figure 4.30a. NK/TP, Pb in silt using MLE – Raw Data	4-58
Figure 4.30b. NK/TP, Pb in silt using MLE – Predicted Data	4-58
Figure 4.31a. NK/TP, Zn in silt using MLE – Raw Data	4-59
Figure 4.31b. NK/TP, Zn in silt using MLE – Predicted Data	4-59
Figure 4.32a. NK/TP. Co in silt using MLE – Raw Data	4-60
Figure 4.32b. NK/TP, Co in silt using MLE – Predicted Data	4-60
Figure 4.33a. NK/TP, Co in HMC using MLE – Raw Data	4-61
Figure 4.33b. NK/TP. Co in HMC using MLE – Predicted Data	4-61
Figure 4.34a. NK/TP, Ba in silt using MLE – Raw Data	4-62
Figure 4.34b. NK/TP, Ba in silt using MLE – Predicted Data	4-62
Figure 4.35a. NK/TP,Ni in HMC using MLE – Raw Data	4-63

Figure 4.35b. NK/TP, Ni in HMC using MLE – Predicted Data	. 4-63
Figure 4.36a. NK/TP, Cr in HMC using MLE – Raw Data	. 4-64
Figure 4.36b. NK/TP, Cr in HMC using MLE – Predicted Data	. 4-64
Figure 4.37a. NK/TP, Cu in silt using MLE – Raw Data	. 4-65
Figure 4.37b. NK/TP, Cu in silt using MLE – Predicted Data	. 4-65
Figure 4.38a. NK/TP, Cd in silt using MLE – Raw Data	. 4-66
Figure 4.38b. NK/TP, Cd in silt using MLE – Predicted Data	. 4-66
Figure 4.39a. NK/TP, Au in HMC using MLE – Raw Data	. 4-67
Figure 4.39b. NK/TP, Au in HMC using MLE – Predicted Data	. 4-67
Figure 4.40a. Ragged Ranges Mineral Potential Summary Map	. 4-76
Figure 4.40b. Ragged Ranges Anomalous Basin Sample Numbers	. 4-77
Figure 4.41a. NK/TP Mineral Potential Summary Map	. 4-78
Figure 4.41b. NK/TP Anomalous Basin Sample Numbers	. 4-79
Figure 5.1. Ragged Ranges spring sample locations on simplified geology base	5-2
Figure 5.2. Nahanni Karst and Tlogotsho Plateau sample locations on simplified geology	
base	5-3
Figure 5.3. Tree plot configuration produced by using all 1986 and 1987 geochemical data.	5-8
Figure 5.4. Geochemistry of spring water samples from the Ragged Ranges study area	. 5-10
Figure 5.5. Geochemistry of spring samples from the Nahanni Karst and Tlogotsho Plateau	
study areas	. 5-11
Figure 5.6. Samples in the Kleiner-Hartigan tree classification which were left ungrouped	. 5-12
Figure 5.7 Results of the hierarchical cluster analysis	. 5-13
Figure 5.8 Mean pH for each group in the final classification (Table 5.5)	. 5-16
Figure 5.9. Mean water temperature for each group	. 5-16
Figure 5.10 Mean total dissolved solids concentration, calculated from major ion data	. 5-16
Figure 5.11 ¹⁸ O and 2H plot of spring waters in the study areas	. 5-17
Figure 5.12 Piper diagram showing major ion chemistry for Group A	. 5-17
Figure 5.13 Schematic explanation for the localization and chemistry of the springs in	
Group A	. 5-18
Figure 5.14. Mean of Fe and Mn for each group, as listed in Table 5.5	. 5-19
Figure 5.15 Mean Zn and Ni concentrations for each group as listed in Table 5.5	. 5-20
Figure 5.16 Stability field for iron in solution (from Hem, 1985)	. 5-20
Figure 5.17 Piper plot showing the major ion chemistry of springs	. 5-21
Figure 5.18 Scatter plot of log(Zn+Ni) vs. log(Fe) for the three subgroups from Group B	. 5-21
Figure 5.19 Piper plot of the major ion chemistry of waters in Group C	. 5-25
Figure 5.20. Calculated partial pressure of CO ₂ for all samples in the non-acidic groups	. 5-25
Figure 5.21 $\delta^{13}C(\text{gas})$ calculated from $\delta^{13}C(\text{DIC})$ for springs in various groups	. 5-26
Figure 5.22 Cross section of the Flat River valley showing the possible model for the origin	of
CO2 rich fluids	. 5-28
Figure 5.23 Piper plot of major ion chemistry of Sub-groups D1 and D2	. 5-28
Figure 5.24 Piper plot showing major ion chemistry for springs in Group E	. 5-32
Figure 5.25 Piper plot of the very low TDS waters of Group F	. 5-34
Figure 5.26. View looking north over "White Aster hot spring"	. 5-38
Figure 6.1a. Mineral occurrences of the South Nahanni River region on geology	6-3
Figure 6.1b. Regional mineral occurrences on regional mineral resource assessment	
domains	6-4
Figure 6.2. Regional mineral resource assessment domains (Fig. 6.1B) in Ragged Ranges stu	ıdy
area with summary data	6-5

Figure 6.3. Regional mineral resource assessment domains (Fig. 6.1B) in Nahanni	Karst –
Tlogotsho Plateau study area	6-11
Figure 8.1. Study area and key geographic locations	
Figure 8.2: Diagram illustrating stratigraphic relationships in the main gas produci	ng interval
in the Devonian	
Figure 8.3: Seismic structure section from Kotaneelee Field	
Figure 8.4: Discovered major gas fields of the Liard Plateau	8-14
Figure 8.5: Illustration of the high rate of exploratory success and performance of the high rate of the hig	the Liard
Plateau gas play	
Figure 8.6: An illustration of historical analysis using the sequential sampling type	of
petroleum assessment model	
Figure 8.7: Predicted natural gas field sizes, as a function of accumulation size	
Figure 8.8. Predicted 15 largest natural gas field sizes	8-27
Figure 8.9. Predicted natural gas field sizes as a function of accumulation size ran	k
for the Geo-anchored resource assessment model	8-28
Figure 8 10: Predicted natural gas field sizes as a function of accumulation size ra	nk for the
reservoir volume resource assessment model that assumes 100 accumulation	8_{-31}
Figure 8 11: Predicted natural gas field sizes, as a function of accumulation size ra	nk for the
Figure 6.11. Fledicted flatural gas field sizes, as a function of accumulation size fa	
Figure 9.12. Desdicted network can field sizes as a function of accumulation size re	
Figure 8.12. Predicted natural gas field sizes, as a function of accumulation size ra	
Figure 9.12. Place and the second state of the three structures. Etcade De	
Figure 8.13: Play area map showing the location of the three structures, Etanda Do	me, I wisted
Mountain Anticline and Mattson Anticline	8-35
Table Captions (*note: tables from appendices are listed in Table of Contents with the appropriate Table 1.1. Resource potential ratings of 5 and above for assessment domains locat Ragged Ranges and Nahanni Karst-Tlogotsho Plateau study areas	ed only within 1-5
Table 1.2. Explanation of mineral and energy potential rating categories	
Table 3.1 Rock types of South Nahanni River region	
Table 4.1: Data Types to Which ML Estimation Can Be Applied	4-17
Table 4.2: Calculation of Mean and Standard Deviation Using MLE and Substituti	on 4-17
Table 4.3: Critical Values to Calculate Percentiles	4-20
Table 4.4 Statistical Elements Which Remain Unchanged/Change in Log-transform	ned and Non-
transformed Environments:	4 20
Table 4.5: Equations Used to Calculate Predicted Values for 100% of a Rock Type	
Table 4.6: Analyses of Gold Grains Found in Heavy Mineral Concentrates made fr	
gravels	
Table 4.7: Comparison of gold from silt heavy mineral concentrates with gold from	e
i wore i. , . Comparison of gold from sht heavy infineral concentrates with gold from	4-20 24-33 24-33 24-33 24-36 n standard silt
samples and from sieved-gravel HMCs	4-20 4-33 rom sandy
samples and from sieved-gravel HMCs	4-20 4-33 rom sandy 4-36 n standard silt 4-39 gravel were
Table 4.8: Samples for which HMCs and INAA analyses from bulk silts vs sand & compared.	4-20 4-33 rom sandy 4-36 n standard silt 4-39 gravel were 4-40
Table 4.8: Samples for which HMCs and INAA analyses from bulk silts vs sand & compared. Table 4.9: Element Associations and Inferred Deposit Types	4-20 4-33 rom sandy 4-36 n standard silt 4-39 gravel were 4-40 4-42
 Table 4.9: Element Associations and Inferred Deposit Types Table 4.10: Summary data used to determine anomalous catchment basins (shown 	4-20 4-33 rom sandy 4-36 n standard silt 4-39 gravel were 4-40 4-42 in Figure
 Table 4.8: Samples for which HMCs and INAA analyses from bulk silts vs sand & compared. Table 4.9: Element Associations and Inferred Deposit Types Table 4.10: Summary data used to determine anomalous catchment basins (shown 4.40a). Ragged Ranges area 	4-20 4-33 rom sandy 4-36 n standard silt 4-39 gravel were 4-40 4-42 in Figure 4-68
 Table 4.1: Comparison of gold from site field y finite a concentrates with gold from samples and from sieved-gravel HMCs Table 4.8: Samples for which HMCs and INAA analyses from bulk silts vs sand & compared. Table 4.9: Element Associations and Inferred Deposit Types Table 4.10: Summary data used to determine anomalous catchment basins (shown 4.40a), Ragged Ranges area Table 4.11: Summary data used to determine anomalous catchment basins (shown 4.40a) 	4-20 4-33 rom sandy 4-36 n standard silt 4-39 gravel were 4-40 4-42 in Figure 4-68 in Figure
 Table 4.1: Comparison of gold from site fleavy inneral concentrates with gold from samples and from sieved-gravel HMCs Table 4.8: Samples for which HMCs and INAA analyses from bulk silts vs sand & compared. Table 4.9: Element Associations and Inferred Deposit Types Table 4.10: Summary data used to determine anomalous catchment basins (shown 4.40a), Ragged Ranges area Table 4.11: Summary data used to determine anomalous catchment basins (shown 4.41a) Nahanni Karst/Tlogotsho Plateau areas 	4-20 4-33 rom sandy 4-36 n standard silt 4-39 gravel were 4-39 gravel were 4-40 4-42 in Figure 4-68 in Figure 4-72
 Table 4.11: Summary data used to determine anomalous catchment basins (shown 4.40a), Ragged Ranges area Table 4.11: Summary data used to determine anomalous catchment basins (shown 4.41a), Nahanni Karst/Tlogotsho Plateau areas Table 5 1 Results of the field classification of spring waters based on physical 	4-20 4-33 rom sandy 4-36 n standard silt 4-39 gravel were 4-40 4-42 in Figure 4-68 in Figure 4-72
 Table 4.11: Comparison of gold from site field y finite a concentrates with gold from samples and from sieved-gravel HMCs Table 4.8: Samples for which HMCs and INAA analyses from bulk silts vs sand & compared. Table 4.9: Element Associations and Inferred Deposit Types	4-20 4-33 rom sandy 4-36 n standard silt 4-39 gravel were 4-40 4-42 in Figure 4-68 in Figure 4-72

Table 5.2. Results of classification by pattern recognition of "Kleiner-Hartigan Trees"	5-8
Table 5.3. Results of classification by hierarchical cluster analysis	. 5-13
Table 5.4. Evolution of classifications in this study, ordered by final groupings	. 5-15
Table 5.5. Final Classification of spring waters in the three South Nahanni River	
study areas	. 5-17
Table 5.6 Major ion and selected trace element concentrations for Group A springs	5-18
Table 5.7. Major ion and selected trace element concentrations for Group B springs	5-23
Table 5.8. Major ion and selected trace element concentrations for Group C springs	5-26
Table. 5.9 Major ions and selected trace element concentrations for Group D springs	5-29
Table. 5.10. Major ion and selected trace element concentrations for Group E springs	5-33
Table 5.11. Major ion and selected trace element concentrations for Group F springs	. 5-34
Table 5.12. Summary of interesting and/or geochemically anomalous ground waters in the	
Nahanni River Region	. 5-39
Table 6.1. Summary of anomalous data in Ragged Ranges Area	6-6
Table 6.2. Summary of anomalous data in Nahanni Karst - Tlogotsho Plateau areas	. 6-12
Table 7.1. Some tectonic classes of mineral deposits in South Nahanni region	7-1
Table 8.1. Gas field size and discovery year	. 8-13
Table 8.2. GSC 1992 estimate of Liard Fold and Thrust Belt undiscovered petroleum	
potential	. 8-17
Table 8.3. Results of previous NEB studies of Liard Fold and Thrust Belt	. 8-18
Table 8.4. Results from the Geo-anchored method (unit bcf)	. 8-26
Table 8.5. Parameters and assessment results from the regional geological approach	. 8-28
Table 8.6. Probability distribution of reservoir parameters(scenario I)	. 8-29
Table 8.7. Probability distribution of reservoir parameters (scenario II)	. 8-29
Table 8.8. Probability distribution of reservoir parameters (scenario III)	. 8-30
Table 8.9. Assessment results from volumetric approach based on scenarios I, II and III	. 8-30
Table 8.10. Summary Results of Various Calculations of Undiscovered Petroleum	
Potential in the Liard Fold and Thrust Belt performed for this study	. 8-32