LEGEND

Note: This legend is common to National Geochemical Reconnaissance Mp. 14-1976, Open File 414, Mp. 15-1976, Open File 415 and Map 16-1976, Open File 416

Canada
Department of Energy, Mines and Resources
Geological Survey of Canada

Parts of 104 D, E and 114 A, H

QUATERNARY
- 31 Unconsolidated glacial and alluvial deposits

MESOZOIC
CRETACEOUS
- 30 Sandstone, shale and conglomerate (Ft. SUNDAY DOME FORMATION)
- 29 Sandstone, shale, coal, minor limestone (Ft. KOTAKA FORMATION) - fine grained, dark and variegated; Ft. KEELE FORMATION - shale and siltstone
- Hornblende and hornblende-biotite syenite, commonly porphyritic
- Hornblende-biotite granite
- 28 Granite and gabbro
- 27 Mottled green and maroon shale and minor siltstone
- 26 SEVEN-O-ALL FORMATION - massive quartzites, minor slate and phyllite conglomerate

TRIASSIC
- 25 LOWER SLEEPY FORMATION - quartzite, slate, phyllite and minor quartz-pebble conglomerate and breccia

PERMIAN
- 24 GRANDAGE FORMATION - grey shale and limestone; Mbr. DONALD CREEK FORMATION - limestone, shale, carbonaceous and clastics, undivided

CARBONIFEROUS TO PERMIAN
- 23 ETHEL FORMATION - shale and limestone; ANNE RIVER FORMATION - shale, siltstone and limestone; Lismore Group undivided - bio, limestone, shale, siltstone, minor sandstone, conglomerate, argillaceous limestone, limestone, clastics and coal
- 22 SIR JOHN FORMATION - shale; undivided conglomerate

DEVONIAN
- 21 DEVONIAN TO CARBONIFEROUS
- 20 LOWER FORMATION - black shale; NATION RIVER FORMATION - shale, siltstone and limestone; Mbr. DONALD CREEK FORMATION - limestone, shale, siltstone and minor chert-calcrete conglomerate and quartzite

UNITS TO MIDDLE DEVONIAN
- 19 MIDDLE FORMATION - limestone, shale; LIMESTONE FORMATION - limestone; MIDDLE FORMATION - limestone and minor dolomite; Mbr. DONALD CREEK FORMATION - limestone, shale, siltstone

ORDOVICIAN AND SILURIAN
- 18 ROAD RIVER FORMATION - shale, limestone, black chert and graptolitic shale; minor quartzite and chert-arenite conglomerate
- 17 Limestone, dolomite and limestone breccias and dolomites; dark to black chert

CAMBRIAN
- 16 Limestone and dolomites, minor red shale; undivided clastics Mb. quartzite, sandstone, conglomerate, shale and local arkosic and feldspathic sandstone; Mbr. CHARLIE LAKE SANDSTONE
- 15 Limestone, dolomite, shale, quartzite, minor sandstone and conglomerate

PRECAMBRIAN AND/OR CAMBRIAN
- 14 Dark green volcanic rocks, breccia, tuff, agglomerate, slate, chert, siltstone and limestone
- 13 Quartzite, sandstone, quartz-pebble conglomerate, narrow and green phyllite, slate, minor dark calcite, quartz-arenite schist, feldspathic and chert clasts

PRECAMBRIAN
- 12 Intrusive carbonates and clastics
- 11 GRAYLING GROUP: mudstone, limestone, iron formation and dolomite

PROTEROZOIC
- 10 SAPOSKOW FORMATION: sandstone and dolomite; FOLSTOW FORMATION: sandstone, conglomerate, siltstone, shale, minor dolomite; QUEEN
- 9 Orange dolomite, sands, quartzite, grey dolomite, shale and narrow shale, quartzite, conglomerate, limestone, black shale; undivided clastics
- HELIKIAN AND OLDER ROCKS
- 8 Argillite, slate, phyllite quartzite, dolomite, conglomerate and siltstone; limestone

Drift-covered area.....
Geological contact.....
Fault.....

Geology generalized from geochemical maps by W.D. Goodfellow from North Creek (116G) and Sheldon Lake (116H) sheets, by J.G. Fyles from Mt. Sidney Williams and Anne River (105P) and Hart River (116B) Sq. E.J. Wheeler (G.S.C. Open File 276, 1973)

Geological cartography by the Geological Survey of Canada

Base-map assembled by the Geological Survey of Canada from maps published at the same scale by the Surveys and Mapping Branch in 1964, 1973

Mean magnetic declination 1973, 34°02.7' East, decreasing 0.3' annually. Readings vary from 34°25.5' in the SE corner to 34°38.4' in the NW corner of the map-area

Elevations in feet above mean sea-level

GEOCHEMICAL SURVEY AND DATA PRESENTATION

The absolute background and anomalous concentrations, and the contrast between them will vary regionally depending on factors such as the physiography, geology, the nature of the drainage and the type of sediment sampled.

Physiographical regions within the survey area include the Hornaday, Ogilvie and Richardson Mountains, and the Eagle Plains and Bonnet Plume basin. Most of the sediments sampled in the mountainous areas are colluvium, residual soil and bedrock. Close compliance of the underlying rocks and has been transported as particulates during the heavy spring run-off. Stream sediments range in size from fine silt to boulders with local concentrations of sand and gravel. Sediments in the mountainous areas of the Ogilvie and Dorset Plane basins are commonly clay-rich, and may also be saline-rich, and deposit organic-rich sediment. Under these conditions, organic matter may be expected to be high. Fe, Mn, Co, Ni, Cu, Zn, Pb, Ag, Cd, Hg, Mo, Sb, Bi, As, Au, Ba, Sr, Ca, Mg, Na, Be, Sc, Ti, Y, La, Ce, Nd, etc.), and may therefore produce spurious anomalies. It is suggested that field observations, such as sediment composition, present in the data (Hall 1970) should be considered when evaluating geochemical survey data.

The geology of the survey area is represented by sedimentary, volcanic and intrusive rocks that range in age from lower(?) Proterozoic to Upper Cenozoic, with almost every period represented. To date, it has been reported to occur in breccias associated with the Proterozoic sedimentary and volcanic rocks of the Wernecke Mountains, and in the Cretaceous sediments of the Eagle Plains basin. Areas of potential in the survey area include the Hollowing- the Mcnamir siltstone shales and batholithic, the folkosaic shales as a source of Sn areas and high tonnage Is the Cretaceous sediments of the Eagle Plains basin. Areas of potential in the survey basins and structures such as faults and unconformities which may serve as favorable traps for deposition of U.

Because of the geological derivation of stream sediments from mountainous terrain, the geochemistry is strongly influenced by the chemistry of the underlying rocks. For example shales, or their metamorphic equivalents, which are common in Proterozoic, Paleozoic and Mesozoic rocks, are known to have relatively high background for a number elements determinant which is reflected in the geochemistry of the stream sediments. Therefore, it is suggested that each stream system be evaluated in terms of the local background, and that anomalies be based on local background.

The element associations expected for particular types of U and base metal mineralization should be considered when evaluating geochemically anomalous stream sediments. For example, Cu, Co, Ni and W is a known association, Pb, Y, and W are quasi-common. The reference to element associations amongst the data in a broad sense would be expected to be accompanied by anomalous values referred to above. Furthermore, element associations in stream sediments and waters will be useful in identifying regional geochemical trends that may be, at least in part, controlled by bedrock lithology. Regional geochemical trends in the data should be carefully examined when be evaluated for possible mineralization on the basis of the weak to strong the anomalous concentrations but also element interrelations.

The data in this publication are presented to be assessed using different criteria than for stream sediment maps. The data for the field duplicate pairs, presented in the data listings, give a measure of the combined field and analytical error. Each sample site was visited only once, therefore temporal variations, variations over a short time period in order to minimize seasonal (climatic) effects. Seasonal variations in the magnitude of anomalies have been detected. However, in terms of contrast between background and anomalous concentrations, the anomalies should be unaffected by over longer periods of time. Considering the above, caution must be exercised when using the geochemistry of stream waters to delineate regional trends, because only limited data from Fe and Mn are not necessarily indicative of mineralized zones. The chemistry of the coating of bedrock and recycling of U and Fe in hydrous oxides can lead to the leaching of bedrock and recycling of U and Fe in hydrous oxides can be expected to give a qualitative indication of the chalcherty. The data should be interpreted in terms of the drainage basin geology, i.e. potential primary source and secondary source. Areas of U and base metal potential should be evaluated on a regional basis only to support a particular model of uranium provenance.

The concentration of an element at a sample site is graphically represented as one of a series of specific symbols, each of which is assigned to a range of values plotted. The symbols are arranged so that they first increase in size to the largest symbol and then increase in blackness to the fifteenth. The two unit crosses at the lowest end of the scale. The class limits are determined by the use of a geometric interval detection limit, or in the data group containing the majority of the values. The data are grouped on a semi-logarithmic scale, i.e. 1,2,5,10,20,50,100 etc. Five decades are usually sufficient to accommodate the total range of the data. Sometimes, however, a series of maps quantitating one element has been chosen for the contouring. The choice of symbols and the data groups they represent is based on the histogram and cumulative frequency plot of the data. Single sample data from one, or more contiguous, open file sheets covered in one field season. Each symbol represents a class interval, and the class limits are listed in each group usually includes the median of the data as defined by the 0.5 (50%) point on the cumulative frequency plot. Some, or all, of the remaining 14 symbols are chosen to emphasize the anomalous data.

The map data symbol maps are only intended to assist the rapid inspection of the data for gross regional features. To fulfil the need for a more specific and detailed interpretation of the data, the original numerical values for each sample should be consulted. To assist in the appraisal of the data in terms of the symbol map bedrock geology, a table of summary statistics for the drainage samples distributed according to rock units underlying the drainage basins is presented on the maps below the histogram. In many instances, the table will also illustrate, more clearly than the maps, the dependence of many geochemical levels on bedrock type. It may be noted that the data for each rock unit appears to approximate a log-normal distribution the data fit both the arithmetic and logarithmic units appears to approximate a normal distribution except where the concentration of the element is near the detection limit for a large number of samples (e.g. U and F in water, Au, Mo and Sn in sediment). In these situations, the frequency distribution will be positively skewed regardless of the type of scale used. The values in the table of statistics must be examined when using the table of summary statistics to establish background and anomalous concentration ranges for a given element.

Other factors, such as pH, Eh, specific gravity, geological, environmental and recorded data should be utilized. The data separation by bedrock type can often be improved by constructing new data subsets and deriving local threshold levels based on the statistical data and up-to-date knowledge available.


RICHARDSON
MOUNTAINS
MACKENZIE
MOUNTAINS
Dempster Highway
Eagle Plains
Bonnet Plume Basin
Ogilvie Mountains
Hornaday River
Peel River
Arctic Red River
Mackenzie River
Fort McPherson
Inuvik
Aklavik
Tuktoyaktuk
Tuktoyaktuk Peninsula
Mackenzie Delta
Beaufort Sea
Yukon Territory
Northwest Territories
Alaska
Dawson
Whitehorse
Mayo
Keno Hill
Stewart River
Klondike River
Sixtymile River
Fortymile River
Yukon River
Porcupine River
Old Crow
Old Crow Flats
Bell River
Rat River
Blackstone River
Blackstone Uplands
Tombstone Mountains
Tombstone Range
Tombstone Plateau
Tombstone Tablelands
Tombstone Pass
Tombstone Gap
Tombstone Mountain
Tombstone Valley
Tombstone Creek
Tombstone Lake
Tombstone Pond