

STRATIGRAPHIC LEGEND

Note: All stratigraphic units from the legend are present on topographic maps.

HOLOCENE (< 0.010 Ma)

Modern soil

Stratified sand and gravel, alluvial sediments

LATE PLEISTOCENE (< 0.125 Ma)

Silt (organic) rich, peat, and organic debris, and extensive interstratified and sagging silt, collectively called muck

Massive to stratified siltstone; colluvial sediments (they locally date to Middle Pleistocene)

Massive to stratified silt and fine sand, siltstone sediments

Stratified silt and sand, reworked/modified colluvial sediments locally interstratified with alluvial sediments

Stratified sand and gravel, alluvial sediments

MIDDLE PLEISTOCENE (0.125 - 0.780 Ma)

Pebbles developed in fluvial and younger pre-fluvial glacial/coluvial sediments

Stratified sand and gravel, glacial/coluvial sediments deposited during first glaciation

Stratified sand and gravel, glacial/coluvial sediments deposited during younger pre-fluvial glaciations

Massive to stratified silt and fine sand, primary and reworked/modified colluvial sediments

LATE PLEISTOCENE TO EARLY PLEISTOCENE (0.7 - 0.780 Ma)

Pebbles developed in fluvial and younger pre-fluvial glacial/coluvial and non-glacial sediments

Stratified sand and gravel, glacial/coluvial sediments deposited during older pre-fluvial glaciations

Stratified to massive siltstone; all deposited during one of several older pre-fluvial glaciations

Fine sand, silt and clay; boulders or slack water fluvial sediments

PLEISTOCENE (pre-glacial) - 2.7 - 5 Ma)

Basalt and basalt breccia

Stratified to massive gravel and sand; White Channel Gneiss and equivalent units underlying regional glacial/coluvial sediments; Late Fluvial sediment


PRE-PLEISTOCENE (> 5 Ma)

Massive and felsic bedrock

SYMBOL

Topo: identification and age, if known, described on stratigraphic log

▲ Ice wedge pseudomorph or sand wedge



LEGEND

Note: Map units listed below occur within one metre of the surface. Where organic or colluvial sediments < 1 m thick underlie these, a pattern is overlaid upon the map unit. Along some valleys, colluvial or alluvial sediments > 1 m thick overlie older alluvial gravels that contain coarse pebbles. In order to approximate these potentially erodible deposits, a contour map unit is presented, e.g., Cd-v. This means that colluvial or alluvial sediments overlie alluvial gravels. The pattern is taken from the Fluvial in age. This legend is part of a larger regional study hence coloured boxes indicate units that occur on this map. In addition, not all symbols in the legend are represented on this map.

CENOZOIC

QUATERNARY HOLOCENE

Modern Level: peat, muck, and silt

ORGANIC DEPOSITS: peat and organic silt formed predominantly by the accumulation of vegetative material in bogs, fens, and swamps situated on valley bottoms, generally to commonly encountered within 1 m of the surface. The most common is peat.

Organic Blanket: undrained; thickness > 1 m to 5 m

Organic Veneer: blanket bog generally < 1 m thick

ALLUVIAL DEPOSITS: gravel to silt size sediments, well stratified, deposited by streams

Floodplain Sediments: gravel, cobble to pebble, massive to well stratified, capped by sand and silt. No lying, includes lacustrine and organic deposits abandoned channels and backwater areas, subject to periodic inundation and reworking by floods; thickness 1 to 5 m

Alluvial Fan Sediments: gravel, sand, silt, and siltstone, massive to well stratified; sediments form fan-shaped fans or complexes of colluvial lobes and terraces at the confluence of tributary streams; may be subject to flooding accompanied by sudden stream migration and inundation; thickness up to 10 m

Alluvial Sediments Complex: sediments forming floodplains, fans, and terraces as above that cannot be subdivided at the map scale

HOLOCENE AND PLEISTOCENE (UNDIVIDED)

COLLUVIAL DEPOSITS: siltstone and sandstone resulting from the physical and chemical breakdown of bedrock and subsequent reworking and transportation by creep, solifluction, and landsliding; colluvial deposits may contain reworked glacial/coluvial and fluvial sediments within the limits of pre-fluvial ice cover and reworked siltstone sediments; colluvial deposits are products of formation and reworking over a significant part of the Pleistocene and Holocene epochs

Colluvial Blanket and Veneer Sediments: siltstone, stony with a sandy matrix; massive to poorly stratified; colluvial blankets generally conform to underlying bedrock and exceed 1 m in thickness, veneers are < 1 m in thickness and are commonly discontinuous over bedrock

Colluvial Apron Sediments: siltstone and stony sandy gravel, poorly sorted, massive sediments form a wedge-like slope complex of small steep debris flow and solifluction deposits; thickness is 1 m at the upper and lower slope limit to up to 5 m or more in the thickest part of the apron

Landslide Sediments: silt to boulders, poorly sorted to unsorted, massive clasts are subangular to angular and are locally derived; thickness varies greatly

Colluvial Complex Sediments: areas of intergrading colluvial and alluvial sediments which are too complex to subdivide at the scale of mapping; unit may include colluvial and alluvial fan, colluvial blanket, landslide sediments and colluvial drift within the limits of glaciation; the unit commonly occurs along the lower slopes of valley margins

Colluvial Eolian Apron (muck): primary deposits of siltstone, fine sand and silt reworked/modified with organic silt, and detrital, alluvial fan gravel and sand and variable amounts of stony colluvial detritus; forms aprons along valley bottoms through reintergradation of siltstone sediments from siltstone to valley floor, commonly preserved on north-facing slopes; thickness 1 to 20 m; commonly contains granitic bodies of ice and buried ice wedges

MIDDLE TO LATE PLEISTOCENE (UNDIVIDED)

ALLUVIAL DEPOSITS: gravel and sand deposited by streams that were not fed by glacial meltwater; sediments may have experienced several cycles of alluviation and erosion, but are now inactive due to burial or fluvial incision; basal gravels within these sediments commonly contain pebble grit

Alluvial Terrace Sediments: gravel, cobble to pebble with a sandy matrix; massive to well stratified; capped by sand and silt; sediments are of flood plain origin now isolated from flooding by stream incision; thickness 1 m to 10 m

Alluvial Fan Sediments: single fans or aprons of colluvial fans formed of gravel and sand, poorly to moderately sorted, now isolated from water and debris floods due to fluvial incision; sediments disturbed by cryoturbation; thickness up to 10 m

Alluvial Complex Sediments: silt, sand and gravel, poorly to moderately sorted; thin to thick bedded; interstratified with colluvial detritus; sediments occupy the floors and margins of narrow upland valleys and grade laterally up slope into colluvial blankets; sediments may represent several depositional cycles; thickness may exceed 10 m in mid-valley locations

EOLIAN DEPOSITS: well sorted medium sand to silt initially transported and deposited by wind action during glaciations and commonly reintergraded through fluvial and colluvial processes; deposits of very fine sand and coarse silt that are distributed discontinuously throughout low lying areas

Eolian Blanket: fine sand and silt, well sorted, massive; may form present shape and their dunes and features or gently undulating near-dune ridges; thickness 1 to 5 m

Eolian Veneer: thin deposits of very fine sand and coarse silt distributed discontinuously throughout low lying areas; thickness < 1 m

LATE PLEISTOCENE - MCDONNELL GLACIATION

GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice, deposits display poor soil development with rare cryoturbation

Glaciofluvial Terrace Sediments: gravel and sand, unweathered, forming one or more terraces

MIDDLE PLEISTOCENE - REID GLACIATION

GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice, deposits display moderate soil development with signs of cryoturbation; soil thickness < 0.5 m

Glaciofluvial Terrace Sediments: gravel and sand, moderately weathered, forming one or more terraces

LATE PLEISTOCENE TO MIDDLE PLEISTOCENE - PRE-REID GLACIATIONS (UNDIVIDED)

GLACIOACRISTINE DEPOSITS: well stratified sand, silt, clay, deposited in lakes ponded by glacial ice

Glacioacristine Undivided: sand, silt, and clay; undifferentiated at the scale of mapping

GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice or meltwater; channels and outwash plains; massive to well stratified. Surface soils may extend to 2 m depth with well developed clay skins on clasts. Frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

Glaciofluvial Terrace Sediments: gravel and sand, deeply weathered, incised into flights of terraces; thickness 1 to > 5 m

MORANIAN DEPOSITS (TILL): glacial detritus, mainly till generally consisting of a matrix ranging from sand to clay that supports clasts ranging from boulders to pebbles in size; deposited either directly from glacial ice or by gravity flow from glacial ice; surface soils may extend to 2 m depth with well developed clay skins on clasts. Frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

Till Blanket: siltstone, stony, silt-sand matrix, massive, conforms to underlying topography; thickness > 1 m; extensively colluviated on slopes

Till Veneer: siltstone, stony, silt-sand matrix, massive, discontinuous and may contain extensive areas of fine (< 1 m) colluvium

ALLUVIAL DEPOSITS: Gravel and sand deposited by streams that were not fed by glacial meltwater; sediments may have experienced several cycles of alluviation and erosion, but are now inactive due to burial or fluvial incision; basal gravels within these sediments commonly contain coarse pebbles

Alluvial Terrace Sediments: sandy pebbles and cobble gravel deposited by streams having a fluvial source but graded to the margins of pre-fluvial glacial or glacial drainage; thickness 1 to 5 m

UNDIFFERENTIATED DRIFT: siltstone, gravel, sand, silt and clay deposited from glacial meltwater; sediments may have experienced several cycles of alluviation and erosion and permafrost make differentiation into component glacial sediments difficult; thickness commonly exceeds 10 m and mass occupying broad topography; commonly colluviated and intergraded with colluvium; surface soils may extend to 2 m depth with well developed clay skins on clasts. Frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

Drift: silt to gently sloping

Drift Modified by Landsliding: drift translated along failure planes into irregular steps and sub-parallel scarp

Fluvially Inclined Drift: formerly extensive areas of drift incised by closely spaced stream valleys

LATE PLEISTOCENE

Basal: columnar alkali feldspar and feldspar, erosional remnants of formerly valley filling flows underlying terraces along lower Roubidou Creek; thickness 10 m

PLEISTOCENE AND LATE PLEISTOCENE

ALLUVIAL DEPOSITS: principal gravel and sand, highly dissected and deeply weathered

Pediment and Bajaze Sediments: inclined fluvial surfaces which are found at a mid-valley position in unglaciated drainage systems; locally thicker than 5 m; formed as a result of limited aggradation of stream gravel and significant colluviation; composition of fine, stony sandy gravel that contains both locally derived subangular stream gravel deposits and angular bedrock fragments

High Level Terrace Sediments (includes White Channel Gneiss and equivalent sediments): weathered pebbles to cobble gravel > 1 m thick; surface soils may extend to 2 m depth with well developed clay skins on clasts. Frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering; terraces above the 500 m contour may be remnant features from the southward flowing paleo-Yukon drainage system

PRE-LATE TERTIARY

Bedrock: siltstone, granite, ultramafic, gneiss, mica-schist, and basalt; includes areas of thin colluvial cover, blockfields, and sorted slope polygons in alpine areas

SYMBOLS

Geologic contact: defined, approximate, inferred

Open system pingo, collapsed open system pingo

Terrestrial collapse activity

Landslide movement direction in bedrock and colluvium

Scars created by widespread landslide movement in drift

Terrace scarp (folds on steep side)

Degraded Arnie: active during pre-fluvial glaciations

Meltwater channel: flow direction, unknown flow direction

Meltwater channel: large

At the (pre-fluvial) glacial limit: defined, inferred

Cryoturbation terrace

Ter

Landsform Shaped by glacial ice

Vestibule foot locally

Stratigraphic section

Radiocarbon date in years (BP/2σ)

A-A' radiometric age on basalt

Fault trace

Lineaments (fault, fracture, joint system) defined by linear drainage courses, aligned gaps in ridges, or aligned breaks in bedrock ridges

Abandoned valley; paleoflow defined

Abandoned valley; paleoflow undefined

Paleoflow; suspected buried valley

Geology by C.A. Hucroft, 1999, 2000

Co-ordinated through the auspices of the Ancient Pacific Margin NATMAP

Digital cartography by K. Shimamura, Terrain Sciences Division

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map from data compiled by Geomatics Canada, modified by Terrain Sciences Division

Magnetic declination 2002, 261°E, decreasing 18.6W annually

Elevations in feet above mean sea level

Contour interval 100 feet

Section 1

Overlying sediments removed by mining

Bouldery cobble gravel, disorganized to weakly stratified, moderately sorted, underlain by subangular clasts, clay supported, clasts composed of high-grade metamorphic lithologies and very rare volcanic lithologies; coarse matrix near upper surface, open framework near base of exposure; base not seen

Section 2

Covered

Silty fine sand, laminated, unit fine upward; disturbed by roots, contains rare isolated angular pebbles, as well as thin gravel lenses; several discontinuous thin beds of fine sand are slightly enriched with humified organics (paleo-A horizon); sharp lower contact

Very fine to coarse sand and silt; organic rich matrix; laminated with rare irregular oxidized coarse sand lenses; thin sand beds unstratified around coarse sand lenses; normally indurated; base not seen

Section 3

Silty fine sand, massive, round pebbles, brown (D YR); lower contact is abrupt and planar

Cobble pebble gravel, stratified, clay supported, poorly sorted; coarse sand matrix; coarse rare boulders; rounded to subangular clasts; vertical to upper boundary; dark lenses (2 A YR) soil development with common moderate thick clay skins on clasts; a thin thick carbonate accumulation on bottom of clasts; 10% of clasts are fractured; 15% are weathered to sand; base not seen

Pebbly gravel, clay bearing; dark yellowish brown (10 YR); many chemically weathered clasts; base not seen

Section 4

Silty fine sand, dark brown to dark reddish brown soil development; massive, abrupt and planar lower contact

Cobble pebble gravel, stratified, clay supported, poorly sorted; coarse sand matrix; coarse rare boulders; rounded to subangular clasts; vertical to upper boundary; dark lenses (2 A YR) soil development with common moderate thick clay skins on clasts; a thin thick carbonate accumulation on bottom of clasts; 10% of clasts are fractured; 15% are weathered to sand; base not seen

Section 5

Cobbly rubble, massive, matrix supported (80% clasts); monomictic; contains rare boulders; very angular, very poorly sorted; micaceous poorly sorted silty sand matrix; base not seen

Section 6

Silty fine sand; massive; brown to yellowish brown (10 YR); lower contact is abrupt and planar

Very fine to coarse sand; interstratified, moderately well sorted; rare laminations and thin beds of greenish-brown sand and gravels; abrupt and planar lower contact

Cobbly pebble gravel, stratified, subangular clasts, with moderately sorted interstratified sand; lower contact is sharp and planar but truncated by ice wedge pseudomorphs in unit below

Cobbly, matrix supported pebble gravels; the long axis of most clasts are vertically oriented; clasts are rounded, granitic and mica-bearing metamorphic clasts are highly weathered and disaggregated; unit is commonly cut by landscape pseudomorphs with clay supported rounded pebbles; matrix is homogeneous, mostly silt matrix with areas of coarse sand; lower contact is gradual and irregular

Boulder gravel; rounded clasts; poorly sorted matrix; lower contact is gradual and irregular

Section 7

Overlying sediments and soil removed by mining

Interstratified silty sand and organic material; rich in black to dark brown humified organic material and contains peat and wood; most stems have well cross-section; base of unit has abundant poorly sorted sand lenses; lower contact is glacial

Cobble pebble gravel; weakly stratified, poorly sorted angular to subangular clasts; poly-lithic; clay supported, coarse sand matrix; lower contact is abrupt and planar

Interbedded fine sand and coarse sand; cross-stratified, well sorted; lower contact is sharp and planar

Organic rich silt, stratified; contains wood; very steep lower contact

Weakly weathered, poorly sorted, isolated cobble pebble gravel; angular to subangular clasts; lower contact covered

OPEN FILE 4345
SURFICIAL GEOLOGY
BRITANNIA CREEK
YUKON TERRITORY

Scale 1:50 000 Échelle 1:50 000

Universal Transverse Mercator Projection
North American Datum 1983
Projection transverse universelle de Mercator
Système de référence géodésique nord-américain, 1983
© Her Majesty the Queen in Right of Canada, 2002

119075	119076	119077	119078	119079	119080
119081	119082	119083	119084	119085	119086
119087	119088	119089	119090	119091	119092
119093	119094	119095	119096	119097	119098
119099	119100	119101	119102	119103	119104
119105	119106	119107	119108	119109	119110
119111	119112	119113	119114	119115	119116
119117	119118	119119	119120	119121	119122

NATIONAL TOPOGRAPHIC SYSTEM REFERENCE



OPEN FILE DOSSIER PUBLIC

4345

Open file products are available through the GSC Internet website.

Les dossiers publics sont disponibles sur le site Internet du GSC.

2002

Recommended citation:
Hucroft, C.A.
2002. Surficial Geology, Britannia Creek, Yukon Territory (115110).
Geological Survey of Canada, Open File 4345, scale 1:50 000.