



OPEN FILE 4581
SURFICIAL GEOLOGY
LOS ANGELES CREEK
YUKON TERRITORY
Scale 1:50 000/Echelle 1/50 000

Universal Transverse Mercator Projection / North American Datum 1983
Projection Transverse Mercator de Méridien / Système de référence géodésique nord américain, 1983
© Her Majesty the Queen in Right of Canada 2005. / © Sa Majesté la Reine du chef du Canada, 2005.

Co-ordinated through the auspices of the Ancient Pacific Margin NATMAP

115 N13	115 N16	115 O13	115 O16	115 O19
OF479	OF480	OF481	OF482	OF483
115 N10	115 N9	115 O11	115 O18	115 O18
OF478	OF477	OF488	OF487	OF486
115 N17	115 N4	115 O3	115 O2	115 O18
OF475	OF476	OF482	OF483	OF484
115 N2	115 N1	115 O4	115 O5	115 O1
OF474	OF473	OF480	OF489	OF487
115 K15	115 K16	115 J13	115 J15	115 J15
OF454	OF455	OF456	OF457	OF458

Digital cartography K. Shimamura, Terrain Sciences Division
Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada.
Digital base map from data compiled by Geomatics Canada, modified by Pam Drees, Earth Sciences Sector Information Division (ESS/ID).
Magnetic declination 2005, 25°14' E, decreasing 18.4' annually
Elevations in feet above mean sea level
Contour interval 100 feet

Geology by L.E. Jackson, Jr. (1989-2002)

Open file no. 4581
Geological Survey of Canada
This dossier public no. 4581
2005

Recommended citation:
Jackson, L.E., Jr., 2005. Surficial Geology, LOS ANGELES CREEK, Yukon Territory; Geological Survey of Canada, Open File 4581, scale 1:50 000.

LEGEND

Note: Map units listed below occur within or meet at the surface. Where organic or siltstone sediments < 1 m thick overlie these, a pattern is overlaid upon the map unit. Along some valleys, colluvial or alluvial sediments > 1 m thick overlie older alluvial gravels that could contain placer gold. In order to accurately show potentially exploitable deposits, a compound map unit is presented, e.g., CwM1. This means that colluvial complex sediments overlie alluvial terrace sediments thought to be late Tertiary in age. This legend is part of a larger regional study hence coloured boxes indicate units that appear on the map. In addition, not all symbols in the legend are represented on the map.

CENOZOIC
QUATERNARY
HOLOCENE
m **Made Land:** placer mines, roads, and airstrip

ORGANIC DEPOSITS: peat and organic silt formed predominantly by the accumulation of vegetative material in bogs, fens, and swamps situated on valley bottoms; peat/organic silt commonly encountered within 1 m of the surface. Thermokarst collapse is common.

ORGANIC BLANKET: undivided; thickness > 1 m to 5 m

ORGANIC VENEER: blanket bog generally < 1 m thick

ALLUVIAL DEPOSITS: gravel to silt size sediments, well stratified, deposited by streams

Floodplain Sediments: gravel, cobble to pebbles; massive to well stratified, capped by sand and silt; fill lying; includes lacustrine and organic deposits in abandoned channels and backswamp areas; subject to periodic inundation and reworking by floods; thickness 1 to 5 m

Alluvial Fan Sediments: gravel, sand, silt, and siltstone, massive to well stratified; sediments form fan-shaped aprons or complexes of coalesced fan-shaped landform at the confluence of tributary streams; may be subject to flooding accompanied by sudden stream migration and inundation; thickness up to 10 m

Alluvial Sediments Complex: sediments forming floodplains, fans, and terraces that cannot be subdivided at this map scale

HOLOCENE AND PLEISTOCENE (UNDIVIDED)
COLLUVIAL DEPOSITS: stony clastic material resulting from the physical and chemical breakdown of bedrock and subsequent reworking and transportation by creep, solifluction, and landsliding; colluvial deposits include glacial till and moraine sediments within the limits of pre-Ried ice-cover and reworked siltstone sediments; colluvial deposits are products of formation and reworking over a significant part of the Pleistocene and Holocene epochs; surface is commonly hummocky or undulating

Colluvial Blanket and Veneer Sediments: clastic, stony with a sandy matrix; massive to poorly stratified; colluvial blankets generally conform to underlying bedrock and exceed 1 m in thickness; veneers < 1 m in thickness and are commonly discontinuous over bedrock

Colluvial Apron Sediments: blocky clastic and blocky sandy gravel, poorly sorted; massive to moderately sorted; colluvial blankets; thickness 1 to 10 m

Landslide Sediments: silt loam to boulders, poorly sorted to unsorted; massive; clasts are subangular to angular and are locally derived; thickness varies greatly

Colluvial Complex Sediments: areas of intergrading colluvial and alluvial sediments which are too complex to subdivide at the scale of mapping; unit may include colluvial and alluvial fan, colluvial blanket, landslide sediments and colluvial drift within the limits of glaciation; the unit commonly occurs along the lower slopes of valley margins

Colluvial/Eolian Apron (muck): primary deposits of eolian fine sand and silt reworked and interstratified with organic silt, and detrital, alluvial fan gravel and sand; variable amounts of stony colluvial clastics; forms aprons along valley bottoms through reindimentation of eolian sediments from valley sides to valley floor; commonly present on north-facing slopes; thickness 1 to 20 m; commonly contains aggregated bodies of ice and buried ice wedges

MIDDLE TO LATE PLEISTOCENE (UNDIVIDED)
ALLUVIAL DEPOSITS: gravel and sand deposited by streams that were not fed by glacial meltwater; sediments may have experienced several cycles of alluviation and erosion, but are now inactive due to burial or fluvial incision; basal gravels within these sediments commonly contain placer gold

Alluvial Terrace Sediments: gravel, cobble to pebbles with a sandy matrix; massive to well stratified; capped by sand and silt; sediments are of flood plain origin now isolated from flooding by stream incision; thickness 1 to 10 m

Alluvial Fan Sediments: single fans or aprons of coalesced fan formed of gravel and sand; poorly to moderately sorted, now isolated from water and debris floods due to fluvial incision; sediments disturbed by cryoturbation; thickness up to 10 m

Alluvial/Colluvial Complex Sediments: silt, sand and gravel, poorly to moderately sorted; with thick bedded, interstratified with colluvial clastics; sediments underlie the floors and margins of narrow upland valleys and grade laterally up slope into colluvial blankets; sediments may represent several depositional cycles; thickness may exceed 10 m in mid-valley locations

EOLIAN DEPOSITS: well sorted medium sand to silt initially transported and deposited by wind action during glacial periods and commonly reworked through fluvial and colluvial processes; deposits of very fine sand and coarse silt < 1 m thick are distributed discontinuously throughout low lying areas

Eolian Blanket: fine sand and silt, well sorted; massive; may form crescentic shape and linear dunes and features to gently undulating inter-dune eolian plains; thickness 1 to 5 m

Eolian Veneer: thin deposits of very fine sand and coarse silt distributed discontinuously throughout low lying areas; thickness < 1 m

LATE PLEISTOCENE - MCCONNELL GLACIATION
GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice; deposits display poor soil development with rare cryoturbation

GLACIOFLUVIAL Terrace Sediments: gravel and sand, unweathered, forming one or more terraces

MIDDLE PLEISTOCENE - REED GLACIATION
GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice; deposits display moderate soil development with signs of cryoturbation; soil thickness < 0.5 m

GLACIOFLUVIAL Terrace Sediments: gravel and sand, moderately weathered, forming one or more terraces

LATE PLEISTOCENE TO MIDDLE PLEISTOCENE - pre-Reed GLACIATIONS (UNDIVIDED)
GLACIOFLUVIAL DEPOSITS: well stratified sand, silt, clay, deposited in lakes ponded by glacial ice

GLACIOFLUVIAL UNDIVIDED: sand, silt, and clay, unstratified at this scale of mapping

GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice in meltwater channels and outwash plains; massive to well stratified; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

GLACIOFLUVIAL Terrace Sediments: gravel and sand, deeply weathered; incised into flights of terraces; thickness 1 to > 5 m

MOORIAL DEPOSITS (TLL): glacial clastic, mainly silt, generally consisting of a matrix ranging from sand to clay that supports coarse ranging from boulders to pebbles in size; deposited either directly from glacial ice or by gravity flow from glacial ice; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

Till Blanket: clastic, stony, silt-and-matrix, massive, conforms to underlying topography; thickness > 1 m; extensively colluviated on slopes

Till Veneer: clastic, stony, silt-and-matrix, massive, discontinuous and may contain extensive areas of < 1 m colluvium

ALLUVIAL DEPOSITS: gravel and sand deposited by streams that were not fed by glacial meltwater; sediments may have experienced several cycles of alluviation and erosion, but are now inactive due to burial or fluvial incision; basal gravels within these sediments commonly contain placer gold

Alluvial Terrace Sediments: sandy pebbles and cobble gravel deposited by streams having a fluvial source but graded to the margins of pre-Ried glaciers or glacial drainage; thickness 1 to 5 m

UNDIFFERENTIATED DRIFT: clastic, gravel, sand, silt and clay deposited from glacial ice, glacial streams, and glacially derived lakes; extensive weathering, poor exposure and poor sorting; interstratified into component glacial sediments; thickness commonly exceed 10 m and mask underlying bedrock topography; commonly colluviated and intergraded with colluvium; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

DRIFT: drift to gently sloping

DRIFT Modified by Landsliding: drift translated along failure plane into irregular steps and sub parallel scarp

FLUVIALLY INCISED DRIFT: formerly extensive areas of drift incised by closely spaced stream valleys

LATE PLEISTOCENE
Basal: columnar alkaline olivine basal and flow breccia; erosional remnants of formerly valley filling flows underlying terraces along lower Foothold Creek; thickness 10 m

FLUORENCE AND LATE MIOCENE
ALLUVIAL DEPOSITS: preglacial gravel and sand; highly dissected and deeply weathered

Piedmont and Basins Sediments: inclined fluvial surfaces which are found at a mid-slope position in unglaciated drainage systems; usually thinner than 5 m; formed as a result of limited aggradation of stream gravel and significant colluvial components of this unit; surface gravel contains highly developed subangular stream gravel deposits and angular bedrock fragments

High Level Terrace Sediments (includes White Channel Gravel and equivalent sediments): weathered pebbles to cobble gravel > 1 m thick; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering; within the Yukon River valley, terraces above the 500 m contour may be reworked features from the southward flowing paleo-Yukon River drainage system

PALEOZOIC AND MESOZOIC
Bedrock: schist, gneiss, ultramafic, granodiorite, monzonite, mafic, and basalt; includes areas of thin glacial cover, blockfields, and sorted stone polygons in slope areas

SYMBOLS

Geologic contact, defined, approximate, inferred
Open system pierce, collapsed open system pierce
Thermokarst collapse activity
Landslide movement direction in bedrock and colluvium
Scarps created by widespread landslide movement in drift
Terrace scarp (rills on steep side)
Degraded Cirque active during pre-Ried Glaciations
Degraded Active during pre-Ried Glaciations
Malwater channel; flow direction, unknown flow direction
Large malwater channel
All time (pre-Ried) glacial limit; defined, inferred
Cryoturbation terrace
Tor
Landform Streamlined by glacial ice
Vertebrate fossil locality
Stratigraphic section
Fault trace
Lineaments (fault, fracture, joint system) defined by linear drainage courses, aligned gaps in ridges, or aligned breaks in bedrock outcrops
Abandoned valley; paleoflow defined
Abandoned valley; paleoflow undefined
Paleoflow, suspected buried valley
Rock glacier

DESCRIPTIVE NOTES

The Los Angeles Creek map area lies within the Kooninka Plateau, an incised rolling upland predominantly underlain by Pleistocene and pre-Ried (RD), Anzures and bedrock metamorphic rocks of the Carmacks Group of Middle Cretaceous to Paleocene age distributed across the map area. They overlap, intrude or are in fault contact with the crystalline basement rocks. These volcanic rocks are partly truncated by the Kooninka Plateau surface. This relationship provides a timing lower age for the plateau. The valley of White River and Yukon River are incised 600 m into the plateau. The valley of White River, that drains an area much smaller than Yukon River, is up to two times the width of the Yukon River valley upstream from their confluence. This reflects the geologically recent downcut of Yukon drainage; the course of Yukon River above the White River confluence was formerly a minor tributary to the ancestral White River prior to the glacially induced reorganization of drainage in southern and central Yukon. The contemporary Yukon River drainage was diverted across a low drainage divide across the Dawson Range about 40 km to the southeast (Duk-Rodin et al. 2003). This apparently occurred during the first regional glaciation of southern and central Yukon Territory between ca. 3.1 to 2.8 million years ago (Frees et al. 2000, 2001).

Natural exposure of surficial deposits are rare in the map area and are most plentiful along Yukon River. Dense vegetation obscures much of the area. Permanently frozen ground is a rare feature of the surface and is generally of exposure with hard tools extremely difficult. Consequently, surficial sediments have been largely mapped from the interpretation of photographs.

The Los Angeles Creek map area has been seen glaciated with the exception of small cirque glaciers on the highest peaks. The glacial sediments are the Carmacks Group and are underlain by the Kooninka Plateau surface. The geology of the map area. Colluvium is the dominant sediment. It is formed by the breakdown of bedrock into regolith that is transported down-slope by gravitational processes such as solifluction and landsliding. With few exceptions, major sedimentary basins have occurred in the incised valleys of the Carmacks Group.

Placer deposits are confined to low terraces and valley bottoms. Terraces and valley bottom gravels may locally contain economically viable amounts of placer gold. Placer gravels along Fines Creek are the only ones to be successfully mined to date. However, placer potential of most of the map area is poorly known.

Strong winds during glacial periods and the Holocene have deposited extensive areas of loess in sheltered steep-sided valleys. This has been reworked and mixed with organic sediments to form loess accumulations called muck. They commonly contain extensive bodies of aggregated ice. Organic deposits in bogs and fens are extensive in valley bottom settings and also commonly contain massive lenses of ice.

REFERENCES

Beacock, H.B., 1942. Geology, Geological Survey of Canada map 71A, 1:253,460.
Duk-Rodin, A., Berendregt, R.W., White, J.M., and Singhvi, V.N., 2001. Geologic evolution of the Yukon River: implications for placer gold. Quaternary International, v. 82, p. 5-20.
Frees, D.G., Berendregt, R.W., Enkin, R.J., and Baker, J., 2000. Paleogeographic evidence for multiple late Pleistocene-early Pleistocene glaciations in the Kooninka area, Yukon Territory. Canadian Journal of Earth Sciences, v. 37, p. 857-877.
Frees, D.G., Aget, T., Duk-Rodin, A., Westgate, J., and White, J. and Smith, D., 2001. Drainage reversal and integration of the Pleistocene upper Yukon River (3.1-2.8) key new evidence; Astoria, Arctic Workshop, Ashcroft, Massachusetts.