

<b>Table 2. Characteristics of post-tectonic plutons</b>				
Name <sup>1</sup>	Area (km <sup>2</sup> )	% of total plutonic exposure	Mean specific gravity <sup>2</sup>	Age  (K-Ar) <sup>3</sup>
<b>Plutons southwest of Tintina Fault</b>				
Black River batholith	160	5.7	2.62	<b>66, 46.9, 53.9</b>
Tintina plug	7	0.2	2.64	-
White Creek stock	4	0.1	-	<b>99.2</b>
Bacon stock	8	0.2	2.64	-
Mt. Cook stock	< 1	-	-	-
Fox batholith	79	2.9	2.68	-
Big Salmon batholith	220	8.0	2.65	<b>92</b>
Nisutlin batholith:				
- NE arm	150	5.4	2.64	<b>96.0, 91.2</b>
- main SE part	600	21.7		<b>86.4</b>
- NW part	130	4.7		<b>88.1</b>
Total	800	31.8		
Dycer Creek stock	105	3.8	-	<b>76.4</b>
Quiet Lake batholith	1250	45.2	-	<b>85.3, 70.1</b>
Red Mtn. plug	50	1.8	-	-
<b>Total</b>	<b>2765</b>			
<b>Plutons northeast of Tintina Fault</b>				
Fyre Lake plug	85	18.4	2.64	-
Grass Lakes plug	55	11.9	2.72	-
North Lakes plug	62	13.4	2.76	-
Mink Creek plug	40	8.6	-	-
Money Creek plug	105	22.7	-	-
Fortin Lake stock	45	9.7	2.65	<b>99.6</b>
McEvoy Lake stock	70	15.2	2.72	-
<b>Total</b>	<b>462</b>			
<sup>1</sup> Pluton names modified to match those used by Yukon Geological Survey.				
<sup>2</sup> Mean specific gravity calculated from minimum of 5 samples				
<sup>3</sup> Age data reported in Appendix 3 and published reports				