


IRON
 Percentage in stream sediments




LEGEND

- DEVONIAN**
 - uDMcg** Grey weathering, resistant, thin- to very thick-bedded, massive, chert pebble conglomerate and medium- to coarse-grained, light- to dark-grey, chert-quartz arenite and wacke; minor brown weathering, blue-grey to black shale, siltstone, and slate
 - uDMps** Brown weathering, recessive, thin bedded, laminated, blue-grey to black shale, siltstone, and slate; minor grey-brown weathering, thin- to medium-bedded, fine- to medium-grained, chert-quartz arenite and wacke
 - muDpt** Black, gun-blue or silvery white weathering, thin bedded, siliceous, black shale, chert, and slate; merges with muDt to southwest by increase in proportion of chert
- ORDOVICIAN SILURIAN**
 - Sp** Orange weathering, resistant, thick bedded, dolomitic, silty, grey mudstone characterized by discontinuous wispy black lamination and locally by abundant small pyrite cubes
 - OSp** Black, gun-blue or silvery white weathering, recessive, black slate; minor thin interbeds of finely crystalline, black limestone and black chert
 - uEIOI** White to buff weathering, laminated or thin bedded, finely crystalline, blue-grey limestone; includes in upper part northeast of Howard's Pass, thin bedded, finely crystalline, nodular, silty limestone; local thin bedded to massive, pale green, lapilli tuff
- CAMBRIAN**
 - ImEp** Tan weathering, resistant, medium bedded, variably calcareous and dolomitic, blue-grey siltstone and mudstone; parallel lamination in grey to black disrupted to discontinuous wispy lamination
 - IEp** Brown to orange brown weathering, recessive, thin bedded, blue-grey slate and siltstone; minor fine grained subarkose to quartz arenite
 - IEcG** Lenticular bodies of white weathering limestone conglomerate and minor blue-grey finely crystalline limestone conglomerate clasts include fine grained blue-grey limestone, oolitic limestone, and archeocyatha; matrix is orange to grey weathering, fine grained, locally sandy limestone
- HADRYNIAN & CAMBRIAN**
 - HIcPs** Dark brown to rust weathering, thin- to thick-bedded, greenish grey siltstone; very fine grained quartz arenite and/or subarkose; slate; southwest of South Nahanni River - dark brown weathering, pale green to blue-grey slate and siltstone, and minor greenish grey, very fine grained, quartz sandstone
 - HIcP** Maroon, purple or green weathering, recessive slate, thin bedded or laminated in like colours; minor thin intervals of thin- to medium-bedded, fine grained, pale green, quartz arenite to subarkose and interbedded pale green to tan slate
 - HIcS** Orange, grey or tan weathering, thin- to medium-bedded, fine grained, pale green, quartz arenite to subarkose and interbedded pale green to tan slate
- HADRYNIAN**
 - Hsp** Grey to brown weathering, thin- to thick-bedded, coarse grained, calcareous, grey quartz arenite and subarkose; quartz pebble conglomerate; brown to pale green slate; minor thin bedded grey or white finely crystalline limestone; sandstone contains conspicuous blue quartz, minor plagioclase and orthoclase

SYMBOLS

- Fault (known, assumed)
- Geological Contact (known, assumed)
- Contact with Quaternary deposits
- 2.30..... Iron (pct) in stream sediments



INDEX MAP - LIEU DE LA CARTE

OPEN FILE
 DOSSIER PUBLIC
845-12
 1982
 GEOLOGICAL SURVEY
 COMMISSION GÉOLOGIQUE
 OTTAWA

This map has been reprinted from a scanned version of the original map
 Reproduction par numérisation d'une carte sur papier