
## LEAD (ppm)

G.S.C. OPEN FILE 868 YUKON AND NORTHWEST TERRITORIES, 1981 NAHANNI MAP (NTS 1051)

## Geochemical Symbol and Data Presentation

The concentration of each element is represented by the actual value plotted adjacent to the sample site represented by a "+" symbol. In addition to enhance visual impact, values over the 75th percentile are designated by grey solid squares which are symmetrically arranged so that they increase in size from the 75th to the 99th percentile. The actual concentration range represented by each symbol is illustrated below with a histogram.

In addition to 25 geochemical maps, each Open File contains an appendix consisting of a short discussion of the geochemistry, survey and analytical methodologies, listing of field and analytical data, and statistical data. The statistical data is provided for the total data set as well as for data subsets grouped on the basis of major stratigraphic units.



## SELECTED MINERAL DEPOSITS AND OCCURRENCES

- Stratabound Zn-Pb (Lower Silurian Age)
- ▲ Stratabound Zn-Pb-Ba (Devonian Age)
- Stratabound Barite (Devonian Age)
- ◆ Replacement Zn, Pb (age unknown)
- Vein Zn, Pb, Ag, Au, Sb (age unknown)
- ▼ Skarn W, Zn (Cretaceous)

Note: Further information on each occurrence or deposit is given in the Appendix which accompanies this open file.

> Geochemistry by W.D. Goodfellow Geological Survey of Canada Resource Geophysics and Geochemistry Division

## CONTRACTORS

Sample collection by Marshall Macklin Monaghan Ltd., Toronto, Uranium in sediment chemical analysis by Nova Track Ltd., Vancouver. Other sediment chemical analysis by Bondar-Clegg and Company, Ottawa

This map forms one of a series of 26 maps released by the Geological Survey of Canada on Open File 868. Each Open File consists of maps for 19 elements for stream sediments, 5 elements for stream waters, and 1 each for water pH and sample site location.


> The Director Geological Survey of Canada 601 Booth Street Ottawa, Ontario K1A OE8


The data are also available in digital form. For further information please contact:

The Director Computer Science Centre Department of Energy, Mines and Resources Ottawa, Ontario K1A 0E4



MINISTÈRE DE L'ÉNERGIE, DES MINES ET DES RESSOURCES






Mean magnetic declination 1982, 32058.1' East, decreasing 8.8 annually. Readings vary from 32044.2' in the SE corner to 3305.6' in the NW corner of the map.

LEAD (ppm) G.S.C. OPEN FILE 868 NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 51-1981 STREAM SEDIMENT AND WATER GEOCHEMICAL SURVEY YUKON AND NORTHWEST TERRITORIES, 1981 NAHANNI MAP (NTS 1051)

© Crown Copyrights reserved

Elevation in feet above mean sea level Base map drawn and printed by the Army Survey Establishment R.C.E. 1949-54



Ce document est le produit d'une · This document was produced numérisation par balayage by scanning the original publication.

LEAD (ppm) G.S.C. OPEN FILE 868 YUKON AND NORTHWEST TERRITORIES, 1981 NAHANNI MAP (NTS 1051)

Black to gun-blue weathering, massive, chert and shale clast granule to pebble conglomerate with mud matrix; contains minor quartz sand; clasts commonly matrix supported Black weathering, thin- to medium-bedded, black chert; minor black weathering, black, siliceous shale Black, gun-blue or silvery white weathering, thin bedded, siliceous, black shale, chert, and slate; merges Light grey weathering, resistant, thin- to thick-bedded, fine- to medium crystalline, dark grey limestone mmDI2 Orange weathering, recessive, thin bedded, finely crystalline, dark blue-grey limestone mDI Orange-brown weathering, thin- to medium-bedded, finely crystalline, light- to dark-grey limestone Light grey weathering, resistant, thin- to thick-bedded, fine- to medium crystalline, dark grey limestone, Dark grey weathering, recessive, thin bedded, platy, finely crystalline, black limestone; minor grey weathering, medium bedded, finely crystalline, grey limestone Dark grey weathering, thick bedded, finely crystalline black dolomite; white dolomite filling veins and Light grey weathering, medium bedded, fine- to medium crystalline light- to dark-grey dolomite; member in middle part of unit of dark grey weathering, medium-to thick-bedded, fine- to medium crystalline, in part crinoidal, dark grey dolomite; top of unit marked by alternating light and dark grey dolomite; Blue-grey weathering, resistant. thin- to very thick-bedded, grey crinoidal limestone characterized by abundant crinoid stem fragments with twin axial canals; massive fine- to medium crystalline, grey Dark grey reathering, thin- to medium-bedded, finely crystalline, black limestone Tan, buff or dark grey weathering, recessive, thin bedded, laminated, argillaceous, finely crystalline, black limestone; in the northeast, black weathering, finely crystalline, black, crinoidal limestone with crinoid stem fragments having twin axial canals occurs near top of unit Blue-grey weathering, thin bedded, finely crystalline, porcellaneous, black or dark blue-grey limestone Orange weathering, resistant, thick bedded, dolomitic, silty, grey mudstone characterized by discontinuous wispy black lamination and locally by abundant small pyrite cubes Black, gun-blue or silvery white weathering, recessive, black slate; minor thin interbeds of finelycrystalline, black limestone and black chert; merges with OSt to southwest by increase in proportion of chert, and with upper part of ueOI to east by increase in proportion of limestone Black weathering, thin- to medium-bedded, dark grey to black chert; rare black siliceous shale; minor tan to brown weathering, recessive dark grey shale at base White to grey weathering, thick- to very thick-bedded, massive, medium crystalline, grey dolomite, locally Grey to white weathering, medium- to thick-bedded, massive, fine- to medium crystalline, grey dolomite; in upper part minor thick beds of medium crystalline, black dolomite Brick red weathering, thin- to thick-bedded, maroon mudstone; orange to grey weathering, thick bedded, fine-to medium crystalline, light coloured dolomite; medium bedded, medium- to coarse-grained, dolomitic, grey quartz arenite; thick bedded, finely crystalline, blue-grey limestone White to orange weathering, massive, fine- to medium crystalline, grey dolomite Rust-brown weathering, resistant, pyritic, amydgaloidal basalt; grey and rust-grey weathering, fissile, green tuff; minor dolomite u601 Buff to grey weathering, recessive, thin bedded, finely crystalline, dark grey to black limestone Blue-grey weathering, thin bedded, finely crystalline, porcellaneous, black limestone, minor grey weathering, Grey to white weathering, thick bedded, massive, fine- to medium crystalline, grey to black dolomite; local dolomite breccia with large blocks of finely crystalline, grey dolomite in matrix of coarsely crystalline, white dolomite White to buff weathering, laminated or thin bedded, finely crystalline, blue-grey limestone; includes in upper part northeast of Howard's Pass, thin bedded, finely crystalline, nodular, silty limestone; local thin bedded to massive, pale green, lapilli tuff Tan\_to orange brown weathering, thin bedded, finely crystalline, blue-grey limestone, locally nodular; at base is minor thin bedded, fine grained, grey quartz arenite Light grey weathering, resistant, thick bedded, massive, fine- to medium crystalline, grey dolomite Tan to brown weathering, recessive, thin bedded, finely crystalline, grey limestone Orange weathering, thin- to thick-bedded, finely crystalline, locally sandy, cream, orange, or grey dolomite; minor medium- to thick-bedded, medium grained, white quartz arenite; minor purple weathering upper - bright orange weathering, thin- to thick-bedded, finely crystalline light coloured dolomite middle - purple weathering, recessive, thin- to thick-bedded, brown to purple siltstone and dolomitic siltstone, minor thin bedded, orange weathering dolomite lower - light orange to brown weathering, resistant, medium- to thick-bedded, medium grained, grey quartz arenite and interbedded brown siltstone; thin to thick interbeds of orange weathering dolomite towards top Grey to buff weathering, thin bedded, locally wavy bedded and nodular, finely crystalline blue-grey to black limestone; minor limestone conglomerate with rounded to subangular clasts of blue-grey weathering grey limestone and oolitic limestone in orange weathering, locally sandy, limestone matrix; upper 1/3 of IGId is white weathering, massive, finely crystalline, grey dolomite Tan weathering, resistant, medium bedded, variably calcareous and dolomitic, blue-grey siltstone and mudstone; parallel lamination in grey to black disrupted to discontinuous wispy lamination Brown to orange brown weathering, recessive, thin bedded, blue-grey slate and siltstone; minor fine grained Lenticular bodies of white weathering limestone conglomerate and minor blue-grey finely crystalline limestone; conglomerate clasts include fine grained blue-grey limestone, oolitic limestone, and archeocyatha; matrix is orange to grey weathering, fine grained, locally sandy limestone Dark brown to rust weathering, thin- to thick-bedded, greenish grey siltstone; very fine grained quartz arenite and/or subarkose; slate; southwest of South Nahanni River - dark brown weathering, pale green to blue-grey slate and siltstone, and minor greenish grey, very fine grained, quartz sandstone Buff weathering massive dolomite Maroon, purple or green weathering, recessive slate, thin bedded or laminated in like colours; minor thin intervals of thin- to medium-bedded, fine grained, pale green, quartz arenite to subarkose and interbedded pale green to tan slate Orange, grey or tan weathering, thin- to medium-bedded, fine grained, pale green, quartz arenite to subarkose and interbedded pale green to tan slate Grey to brown weathering, thin- to thick-bedded, coarse grained, caicareous, grey quartz arenite and subarkose; quartz pebble conglomerate; brown to pale green slate; minor thin bedded grey or white finely crystalline limestone; sandstone contains conspicuous blue quartz, minor plagioclase and orthoclase Geology by S.P. Gordey 1977, 1978, 1979, 1980 (with contributions from previous work by S.L. Blusson, J.A. Roddick, and L.H. Green (1967)) Geological boundary (defined, approximate, assumed or extrapolated beneath overburden). Fault, steeply dipping (defined, approximate, assumed or extrapolated beneath overburden; barb on downthrown side). No analytical result ....... Geology by S.P. GORDEY (1981), Geological Survey of Canada, Open File 780

LEGEND

Pelitic hornfels: red-rust to brown weathering, extremely well indurated, massive, fine grained

Tan weathering, thin bedded, ripple cross-laminated siltstone, fine grained sandstone, and shale

shale; minor fine- to medium-grained quartz arenite

Grey weathering resistant, massive, fine-to medium-grained quartz arenite

TRIASSIC

PERMIAN-

Grey weathering, resistant, medium-to coarse-grained, megacrystic (K-feldspar), biotite quartz monzonite

Carbonate hornfels: white to grey weathering, extremely well indurated, fine-to coarse crystalline; large tremolite porphyroblasts abundant in hornfelsed uEOL

Orange to grey weathering, thin bedded, locally lenticular, pale green to blue-grey chert; minor dark dreen

Brown weathering, recessive, thin bedded, blue-grey shale, black laminated quartz siltstone, and pale green

Grey weathering, resistant, thin- to very thick-bedded, massive, chert pebble conglomerate, and medium- to

Brown weathering, recessive, thin bedded, laminated, blue-grey to black shale, siltstone, and slate; minor grey brown weathering, thin- to medium-bedded, fine- to medium-grained, chert-quartz arenite and wacke

coarse-grained, light- to dark-grey, chert-quartz arenite and wacke; minor brown weathering, blue-grey to