

SULPHATE in water (ppm)

G.S.C. OPEN FILE 868
YUKON AND NORTHWEST TERRITORIES, 1981
NAHANNI MAP (NTS 1051)

Geochemical Symbol and Data Presentation

The concentration of each element is represented by the actual value plotted adjacent to the sample site represented by a "u" symbol. In addition to enhance visual impact, values over the 75th percentile are designated by grey solid squares which are symmetrically arranged so that they increase in size from the 75th to the 99th percentile. The actual concentration range represented by each symbol is illustrated below with a histogram.

In addition to 25 geochemical maps, each Open File contains an appendix consisting of a short discussion of the geochemistry, survey and analytical methodologies, listing of field and analytical data, and statistical data. The statistical data is provided for the total data set as well as for data subsets grouped on the basis of major stratigraphic units.

CONCENTRATION	PERCENTILES
142.9 TO 315.8	99TH TO MAX.
112.4 TO 142.8	98TH TO 99TH
74.3 TO 112.3	95TH TO 98TH
57.1 TO 74.2	90TH TO 95TH
34.3 TO 57.0	75TH TO 90TH

SELECTED MINERAL DEPOSITS AND OCCURRENCES

- Stratabound Zn-Pb (Lower Silurian Age)
- ▲ Stratabound Zn-Pb-Ba (Devonian Age)
- Stratabound Barite (Devonian Age)
- ◆ Replacement Zn, Pb (age unknown)
- Veih Zn, Pb, Ag, Au, Sb (age unknown)
- ▼ Skarn W, Zn (Cretaceous)

Note: Further information on each occurrence or deposit is given in the Appendix which accompanies this open file.

Geochemistry by W.D. Goodfellow
Geological Survey of Canada
Resource Geophysics and Geochemistry Division

CONTRACTORS


Sample collection by Marshall Macklin Monaghan Ltd.,
Toronto. Uranium in sediment chemical analysis by Nova Track
Ltd., Vancouver. Other sediment chemical analysis by
Bondar-Clegg and Company, Ottawa

This map forms one of a series of 26 maps released by the
Geological Survey of Canada on Open File 868. Each Open
File consists of maps for 19 elements for stream sediments,
5 elements for stream waters, and 1 each for water pH and
sample site location.

The Director
Geological Survey of Canada
601 Booth Street
Ottawa, Ontario
K1A 0E8

The data are also available in digital form. For further
information please contact:

The Director
Computer Science Centre
Department of Energy, Mines and Resources
Ottawa, Ontario
K1A 0E4



Mean magnetic declination 1982, 32°58.1' East,
decreasing 8.8 annually. Readings vary
from 32°04.2' in the SE corner to 33°5.6'
in the NW corner of the map.

SULPHATE in water (ppm)

G.S.C. OPEN FILE 868

NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 51-1981

STREAM SEDIMENT AND WATER GEOCHEMICAL SURVEY


YUKON AND NORTHWEST TERRITORIES, 1981
NAHANNI MAP (NTS 1051)

Scale 1:250,000

Kilometres 6 0 6 12 18 Kilometres
Miles 4 0 4 8 Miles
Universal Transverse Mercator Projection
© Crown Copyrights reserved

Elevation in feet above mean sea level

Base map drawn and printed by the
Army Survey Establishment R.C.E. 1949-54



1051

LEGEND

CRETACEOUS	Kqm	Grey weathering, resistant, medium to coarse-grained, megacrystic (K-feldspar), biotite quartz monzonite
	Wf	Pelitic hornfels: red-rust to brown weathering, extremely well indurated, massive, fine grained Carbonate hornfels: white to grey weathering, extremely well indurated, fine to coarse crystalline; large tremolite porphyroblasts abundant in hornfelsed uGf.
TRIASSIC	Tc	Tan weathering, thin bedded, ripple cross-laminated siltstone, fine grained sandstone, and shale.
PERMIAN	Pt	Orange to grey weathering, thin bedded, locally lenticular, pale green to blue-grey chert; minor dark green to brown weathering, pale green, splintery shale
CARBONIFEROUS	Cp	Brown weathering, recessive, thin bedded, blue-grey shale, black laminated quartz siltstone, and pale green shale; minor fine- to medium-grained quartz arenite
	Cs	Grey weathering resistant, massive, fine- to medium-grained quartz arenite
	uGms	Grey weathering, resistant, thin- to very thick-bedded, massive, chert pebble conglomerate, and medium- to coarse-grained, light- to dark-grey, chert-quartz arenite and wacke; minor brown weathering, blue-grey to black shale, siltstone, and slate
		Brown weathering, recessive, thin bedded, laminated, blue-grey to black shale, siltstone, and slate; minor grey brown weathering, thin- to medium-bedded, fine- to medium-grained, chert-quartz arenite and wacke
		Black to gun-blue weathering, massive, chert and shale clast granule to pebble conglomerate with mud matrix; contains minor quartz sands; clasts commonly lenticular
	muDpt	Black weathering, thin- to medium-bedded, black chert; minor black weathering, black, siliceous shale
		Black, gun-blue or silvery white weathering, thin bedded, siliceous, black shale, chert, and slate; merges with mud to southwest by increase in proportion of chert
DEVONIAN	mm02	Light grey weathering, resistant, thin- to thick-bedded, fine- to medium crystalline, dark grey limestone
	md01	Orange weathering, recessive, thin bedded, finely crystalline, dark blue-grey limestone
	mm011	Orange-brown weathering, thin- to medium-bedded, finely crystalline, light- to dark-grey limestone
	md011	Light grey weathering, resistant, thin- to thick-bedded, fine- to medium crystalline, dark grey limestone, in part crinoidal
	ld01	Dark grey weathering, recessive, thin bedded, platy, finely crystalline, black limestone; minor grey weathering, medium bedded, finely crystalline, grey limestone
	md0d	Dark grey weathering, thick bedded, finely crystalline black dolomite; white dolomite filling veins and vugs; sparse chert nodules
	ld0d	Light grey weathering, medium bedded, fine- to medium crystalline light- to dark-grey dolomite; member in middle part of unit of dark grey weathering, medium- to thick-bedded, fine- to medium crystalline, in part crinoidal, dark grey dolomite; top of unit marked by alternating light and dark grey dolomite.
	u01	Blue-grey weathering, resistant, thin- to very thick-bedded, grey crinoidal limestone characterized by abundant crinoid stem fragments with twin axial canals; massive fine- to medium crystalline, grey limestone; minor limestone breccia
	sd01	Dark grey weathering, thin- to medium-bedded, finely crystalline, black limestone
DEVONIAN, SILURIAN AND ORDOVICIAN		Tan, buff or dark grey weathering, recessive, thin bedded, laminated, argillaceous, finely crystalline, black limestone; in the northeast, black weathering, finely crystalline, black, crinoidal limestone with crinoid stem fragments having twin axial canals occurs near top of unit
		Blue-grey weathering, thin bedded, finely crystalline, porcellaneous, black or dark blue-grey limestone
	05pt	Orange weathering, resistant, thick bedded, dolomitic, silty, grey mudstone characterized by discontinuous wispy black lamination and locally by abundant small pyrite cubes
		Black, gun-blue or silvery white weathering, recessive, black slate; minor thin interbeds of finely crystalline, black limestone and black chert; merges with 05t to southwest by increase in proportion of chert, and with upper part of u01 to east by increase in proportion of limestone
		Black weathering, thin- to medium-bedded, dark grey to black chert; rare black siliceous shale; minor tan to brown weathering, recessive dark grey shale at base
	u05d	White to grey weathering, thick- to very thick-bedded, massive, medium crystalline, grey dolomite, locally containing abundant nodules of black or grey chert
	u05d	Grey to white weathering, medium- to thick-bedded, massive, fine- to medium crystalline, grey dolomite; in upper part minor thick beds of medium crystalline, black dolomite
SILURIAN, ORDOVICIAN AND CAMBRIAN		Brick red weathering, thin- to thick-bedded, medium mudstone; orange to grey weathering, thick bedded, fine- to medium crystalline, light coloured dolomite; medium bedded, medium- to coarse-grained, dolomitic, grey quartz arenite; thick bedded, finely crystalline, blue-grey limestone
	u0d	White to orange weathering, massive, fine- to medium crystalline, grey dolomite
	u01	Rust-brown weathering, resistant, pyritic, amygdaloidal basalt; grey and rust-grey weathering, fissile, green tuff; minor dolomite
		Buff to grey weathering, recessive, thin bedded, finely crystalline, dark grey to black limestone
	u001	Blue-grey weathering, thin bedded, finely crystalline, porcellaneous, black limestone, minor grey weathering, thin bedded, finely crystalline, grey dolomite
		Grey to white weathering, thick bedded, massive, fine- to medium crystalline, grey to black dolomite; local dolomite breccia with large blocks of finely crystalline, grey dolomite in matrix of coarsely crystalline, white dolomite
	u0101	White to buff weathering, laminated or thin bedded, finely crystalline, blue-grey limestone; includes in upper part northeast of Howard's Pass, thin bedded, finely crystalline, nodular, silty limestone; local thin bedded to massive, pale green, lapilli tuff
	u01	Tan to orange brown weathering, thin bedded, finely crystalline, blue-grey limestone, locally nodular; at base is minor thin bedded, fine grained, grey quartz arenite
	md	Light grey weathering, resistant, thick bedded, massive, fine- to medium crystalline, grey dolomite
	md	Tan to brown weathering, recessive, thin bedded, finely crystalline, grey limestone
	ld0d	Orange weathering, thin- to thick-bedded, finely crystalline, locally sandy, cream, orange, or grey dolomite; minor medium- to thick-bedded, medium grained, white quartz arenite; minor purple weathering, thin bedded, purple siltstone
CAMBRIAN		upper - bright orange weathering, thin- to thick-bedded, finely crystalline light coloured dolomite middle - purple weathering, recessive, thin- to thick-bedded, brown to purple siltstone and dolomitic siltstone; minor thin bedded, orange weathering, orange weathering, orange weathering, orange weathering lower - light orange to brown weathering, resistant, medium- to thick-bedded, medium grained, grey quartz arenite and interbedded brown siltstone; thin to thick interbeds of orange weathering dolomite towards top
		Grey to buff weathering, thin bedded, locally wavy bedded and nodular, finely crystalline blue-grey to black limestone; minor limestone conglomerate with rounded to subangular clasts of blue-grey weathering grey limestone and dolitic limestone in orange weathering, locally sandy, limestone matrix; upper 1/3 of 101 is white weathering, massive, finely crystalline, grey dolomite
	106p	Tan weathering, resistant, medium bedded, variably calcareous and dolomitic, blue-grey siltstone and mudstone; parallel lamination in grey to black disrupted to discontinuous wispy lamination
		Brown to orange brown weathering, recessive, thin bedded, blue-grey slate and siltstone; minor fine grained subarkose to quartz arenite
		Lenticular bodies of white weathering limestone conglomerate and minor blue-grey finely crystalline limestone; conglomerate clasts include fine grained blue-grey limestone, oolitic limestone, and archaocyath; matrix is orange to grey weathering, fine grained, locally sandy limestone
	H10ps	Dark brown to rust weathering, thin- to thick-bedded, greenish grey siltstone; very fine grained quartz arenite and/or subarkose; slate; southwest of South Nahanni River - dark brown weathering, pale green to blue-grey slate and siltstone, and minor greenish grey, very fine grained, quartz sandstone
		Buff weathering massive dolomite
CAMBRIAN AND MIDDLE DEVONIAN	H10p	Maroon, purple or green weathering, recessive slate, thin bedded or laminated in like colours; minor thin intervals of thin- to medium-bedded, fine grained, pale green, quartz arenite to subarkose and interbedded pale green to tan slate
	H10p	Orange, grey or tan weathering, thin- to medium-bedded, fine grained, pale green, quartz arenite to subarkose and interbedded pale green to tan slate
	H10p	Grey to brown weathering, thin- to thick-bedded, coarse grained, calcareous, grey quartz arenite and subarkose; quartz pebble conglomerate; brown to pale green slate; minor thin bedded grey or white finely crystalline limestone; sandstone contains conspicuous blue quartz; minor plagioclase and orthoclase

Geology by S.P. GORDEY 1977, 1978, 1979, 1980 (with contributions from previous work by S.L. Blusson, J.A. Roddick, and L.H. Green (1967))

Limit of outcrop.....
Geological boundary (defined, approximate, assumed or extrapolated).....

Fault, steeply dipping (defined, approximate, assumed or extrapolated beneath overburden; bars on downthrown side).....

No analytical result.....

Geology by S.P. GORDEY (1981), Geological Survey of Canada, Open File 780

SULPHATE in water (ppm)

G.S.C. OPEN FILE 868

YUKON AND NORTHWEST TERRITORIES, 1981
NAHANNI MAP (NTS 1051)

This document was produced
by scanning the original publication.

Ce document est le produit d'une
numérisation par balayage
de la publication originale.