A Geochemical Report on the KRYPTOS Property

submitted as Technical Report for YMEP Grant 20-003 on the MEGA Target Evaluation Module, Hardrock Type and an Assessment Report for filing work on the KRYPTOS claims

> Work performed on: **KRYPTOS 37, 39, 47-50** YE90243, YE90245, YE90253-YE90256

Work applied to: KRYPTOS 15-80 YE90221-YE90266, YE93631-650 Mayo Mining District

**Owner: Gordon Richards** 

Location 115P/01 & 02 Camp on claims at UTM 427,700E, 7,005,135N, Elev 945 m NAD 83, UTM Zone 8

Field work performed by Gordon Richards & Aaron Holway During the period August 4-10, 2020

Report written by Gordon Richards November 1, 2020

#### TABLE OF CONTENTS

SUMMARY		3
HISTORY		4
CLAIMS		5
GEOLOGY		6
GEOCHEMICAL SURVEY		7
General		7
MMI Soil Sampling		8
Rock Chip Sampling		9
Results		9
CONCLUSIONS		10
RECOMMENDATIONS		11
STATEMENT OF COSTS		11
STATEMENT OF QUALIFICATIONS		12
	TABLES all at back of report	

Table 1. Claim Status.

Table 2. Legend for Figure 3.

 Table 3. Rock Sample Descriptions with As, Au, Sb Results

 Table 4. Response Ratios for MMI Soil Samples MEGA 2020

**FIGURES** all at back of report

Figure 1. Location Map of Project.

Figure 2. Claim Map.

Figure 3. Local Geology Map.

Figure 4. 2020 MMI Results and Rock Samples Locations.

Figure 4A. MEGA 2020 Rock Sample Pit Locations.

Figure 5. Au Response Ratios for MMI Samples and Outline of Figures 4 and 4A.

### **DIGITAL COPIES:**

Table 3. Rock Sample Descriptions with As, Au, Sb results and with UTM xls

Table 4. Response Ratios for MMI Soil Samples with UTM. xls

Geochemical Data from SGS and Bureau Veritas Labs in xls, xlsx and PDF format

Figures 1-3 as png and pdf. This report in pdf.

### SUMMARY.

The general area of the KRYPTOS claims was prospected with the aid of YMEP grants awarded to G Richards in 2016, 2019 and to Jeff Mieras in 2017. The property is located on gentle to moderate slopes across the Klondyke Highway about 25 km south Stewart Crossing within NTS map sheets 115P01 & 02. Access was made by truck to a camp along the Klondyke Highway and then by foot about two km to the area of 2020 work as shown on Figures 4, 4A, and 5 from where soil sampling and rock sampling was conducted.

The geology of the area has been described on Canadian Geoscience Map 7 of southwestern McQuesten and parts of northern Carmacks by Ryan, J.J., Colpron, M., and Hayward, N., 2010. Figure 3. The area is shown on that map to be underlain by the Early Mississippian aged Reid Lakes Batholith that is a weakly Kspar-porphyritic, medium-grained granite to quartz monzonite intruding its own volcanic pile in the west portion of the claims in contact with metasedimentary rocks of the Finlayson Assemblage (?) in the east portion of the claims. Loess, about 25 cm thick, blankets most slopes. The claims lie entirely within Reid glaciated terrain with the possible occurrence of pre-Reid glaciated terrain along ridge tops.

Regional Geochemical Data (RGS) provided the original encouragement for prospecting in 2016 using MMI soil sampling. Results of the soil sampling identified four sizeable targets with porphyry geochemical signatures.

In 2017 the KRYPTOS 1-60 claims were staked June 12 to 14 and recorded June 14 to cover known anomalous zones identified from the 2016 work and their extensions. An MMI soil and black spruce twig sampling prospecting program was undertaken on the claims June 16 to 22 and July 1, 2017 to define the extent of known geochemically anomalous zones and search for additional targets.

Results of the 2016-2017 field work were successful in defining four pronounced multi-element anomalous zones with porphyry geochemical signatures. A poorly defined new zone of anomalous metal values occurred in the southeastern third of the claims. It is characterized by consistently very high response ratios for As, Sb, W, and Mn and numerous high response ratios for Au, Bi, Tl, Fe, Zn, Ti and Cs. One outcrop of micaceous quartzite was noted during the soil sampling and the area occurs within an area of metasediments described in Geoscience Map 7

In 2019 the KRYPTOS 61-80 claims were staked August 1, and recorded in Dawson Aug 2, 2019. MMI soil sampling over the new claims and part of the old claims was carried out from Aug 2 to 9, 2019.

Results showed two patterns of anomalous gold measuring two km long and one km long and up to 400 m wide encompassed within large coincident patterns of anomalous As, Sb, W, and Bi. 38 small (0.06 to 0.24 kg) rock chip samples were collected and assayed. Results contained low values of gold and other pathfinder elements leaving the source of the anomalous gold and other elements unexplained.

Work in 2020 involved digging 34 pits near and around soil samples with highly anomalous metal values in order to find and sample mineralized float that would explain the cause of the anomalous metals and help with planning future exploration on the property. Eighteen additional MMI soil samples were also collected in one area to limit the extent of the anomalous metal values. Results did not provide strongly anomalous gold and other metal values that would explain the geochemically anomalous patterns.

## HISTORY.

Previous exploration activity occurred in the 1980's by a Mr Jim Carson with the staking of both quartz and placer claims. Two of these previous claim blocks are the MEGALURUS labelled on the Figure 5 and extending over much of the area sampled in 2019 and the FIRELORD which is a small claim block along the Crooked Creek at the northeast end of the MEGALURUS claim block. Other claims staked by Carson occur north of the KRYPTOS claims along Crooked Creek.

Only minimal hand trenching was recorded. All of this hand trenching was on narrow quartz veins within schist and micaceous quartzites of the Finlayson Assemblage done north and east of the KRYPTOS claim block along steep slopes into Crooked Creek. Work is summarized in Minfile Report 115P 038. Assessment Report 019539 provides some detail to the sampling. Samples submitted by Carson report grades up to 0.36 oz/T Au from selected samples but reports by personnel working for Curragh Resources, Erickson Gold Mining Corp., and Noranda Exploration Company, Limited all reported no gold from samples collected over greater widths. One claim block, the FIREDEVIL staked in 1987 covers the biggest porphyry target defined by the 2016/2017 surveys although no outcrop is known to exist in this area. No record of any work that might have been done on this or the MEGALAURUS claim block have been found although considerable effort was spent trying to find any data on these claims.

Placer claims were staked over a two or three km length along the small creek flowing west immediately south of the KRYPTOS claims. No placer production was recorded or evident along the creek that was staked.

YMEP grants 16-056 and 19-056 awarded to Richards and 17-001 awarded to Jeff Mieras supported extensive MMI soil sampling across the KRYPTOS Claims. Results defined four patterns of strong multi-element geochemical anomalies in MMI soil samples with porphyry mineralization signatures. A fifth zone of anomalous metal values was defined by a 2 km diameter zone of anomalous As-Sb-W-Bi with two contained zones of anomalous Au measuring one and two km long and up to 400 m wide. YMEP grant 20-003 was used to conduct a small MMI soil survey used to define the limits of the anomalous zone and also to collect rock chip samples from angular rock in soil pits dug over and near to previously collected samples that were anomalous for Au. No strong assay results were obtained although there remains some encouragement for additional exploration work on the property.

## CLAIMS.

Table 1 is a list of all claims forming the property. The claims lie in the Mayo and Dawson Mining District with the Klondyke Highway forming the boundary between the districts. The Registered Owner is Gordon G Richards. The work described in this report was funded by a YMEP grant and Richards.

Claim Name	Grant No.	Expiry Date	Mining District
KRYPTOS 1-14	YE90207-YE90220	2025/06/15	Dawson
KRYPTOS 15-60	YE90221-YE90266	2025/06/14	Мауо
KRYPTOS 61-80	YE93631-YE93650	2025/08/02	Мауо

Tab	le 1.	Claim S	Status	befor	e apply	ying	work	in t	his	report
-----	-------	---------	--------	-------	---------	------	------	------	-----	--------

Certificate of Work is to be filed on all of the KRYPTOS claims lying within the Mayo District based on work described in this report.

Apply 2 years to KRYPTOS 15-60: 46 cl X 2yrs X \$100/cl-yr = \$9,200. Apply 2 years to KRYPTOS 61-80: 20 cl X 2yrs X \$100/cl-yr = \$4,000.

# **GEOLOGY**.

Bedrock geology is best described on Canadian Geoscience Map 7 of *Southwestern McQuesten and Parts of Northern Carmacks* by Ryan, J.J., Colpron, M., and Hayward, N., 2010. Figures 3 is a portion of that map covering the general area of the property. The claims area is shown on that map to be underlain by the Early Mississippian aged Reid Lakes Batholith in the west portion of the claims in fault contact with metasedimentary rocks of the Finlayson Assemblage (?) in the east portion of the claims.

The Reid Lakes Batholith is an 80 km long unmetamorphosed Early Mississippian aged batholith that intrudes its own volcanic pile. It is a weakly Kspar-porphyritic, medium-grained granite to quartz monzonite.

The Finlayson Assemblage is a Late Devonian to Early Mississippian metavolcanic and metasedimentary assemblage. The metasediments such as occur on the property are carbonaceous quartzite to mica-quartz schist, black to white quartzite, with schist and garnet schist interlayers; and rare black phyllite, possibly equivalent to Nasina Formation, or simply a carbonaceous member of the Snowcap Assemblage. Two closely spaced outcrops of the metasedimentary rocks were located in 2017 along the most north-easterly sample line. They were both dark grey micaceous quartzite. In August 2019 three more outcrops were located. An outcrop of quartz monzonite occurs on a small knoll at the west end of the recently staked claims near sample Y140. Two other outcrops of micaceous quartzite were located, one along a creek bank near sample site P124 and the other on the north facing hillside near sample site Y83. All three metasedimentary outcrops had near flat to very gently dipping schistosity.

Glaciation in the area of the property is described as Reid in age on several government maps although pre-Reid glaciation may have occurred on the ridge

top in the western half of the recently staked claims. Pre-Reid glaciation is possibly older than 500,000 years (Jeff Bond, personal communication, 2012). Reid glaciation began 200,000 years ago and ended about 50,000 years ago. Younger McConnell Glaciation which lies further east of the claims ended about 20,000 years ago.

Uppermost soil is an organic soil from almost absent to less than one cm thick on dryer slopes and in excess of 10 cm thick over gentle poorly drained slopes. Loess occurs on all slopes, generally about 20 to 30 cm thick beneath the organic soil. This loess is believed to have formed in late stages or soon after the end of McConnell Glaciation. Till is commonly found beneath the loess on the north facing and east facing slopes containing well rounded cobbles and smaller rocks of foreign origin. At higher elevations along the ridgeline till does occur beneath the loess but angular rock fragments believed to be of local origin are common in many soil pits and in a few pits are very abundant.

# **GEOCHEMICAL SURVEY.**

General.

On August 3, G Richards bought supplies, rented radios and organized gear for the work. On August 4, G. Richards and A. Holway drove from Whitehorse to the project area and set up a camp on a small spur road along the Klondyke Highway. From August 5 to 9 they collected MMI and rock samples. On August 10 they packed the camp and samples into their vehicle and drove to Whitehorse. On August 11 Richards sorted samples and shipped them to Vancouver, returned rented radios, payed invoices, and cleaned and stored camping gear. In Vancouver, Richards dropped the MMI samples at the SGS lab, examined and described all the rock samples, splitting them in half and sending half to Bureau Veritas Lab for analysis.

Time spent and samples collected: G Richards. Aug 3-11, **9 man days** A Holway, Aug 4-10, **7 man days 18 MMI soil** samples were collected **75 rock** samples were assayed Four sample series are shown on Figures 4 and 4A. "C" and "T" sample series were collected in 2016; "R" and "K" sample series were collected in 2017; P and Y sample series were collected in 2019; W and some R sample series were collected in 2020.

The 2020 MMI soil sampling program was conducted across a limited portion of the KRYPTOS claims to try and limit the extent of the highest Au response ratio, 227 in sample Y38, from the 2019 MMI soil survey and resample some of the high metal values 200 m east of Y38. Both goals were successful. Three samples around and close to Y38, W5 to W7 returned response ratios of 112, 129 and 509 indicating the possibility of significant underlying gold mineralization. Refer to Figure 4 for location of samples and Table 4 for UTM (NAD83, Zone 8) co-ordinates and response ratios for ten elements. A complete list of geochemical results is provided in attached digital files.

Thirty-three pits were dug across the property at and close to previous high Au response ratios in an attempt to find and sample angular gold mineralized angular float and possibly bedrock that would explain the anomalous gold patterns found in the previous soil surveys. No outcrop was found. Multiple rock samples were collected from each pit with only angular or subangular material sampled. Pits were from 0.5 to 1.0 m deep dug on steep to moderate slopes. Upon collection of rock samples for assay the pit walls were sloughed in and the pits backfilled to complete reclamation. Location of the pits are shown on Figure 4A and rock descriptions are provided in Table 3 with UTM (NAD83, Zone 8) coordinates and geochemical values for As, Au, and Sb. A complete list of geochemical results is provided in attached digital files.

### **MMI Soil Sampling.**

Watch and ring were removed prior to sampling. Pits were dug by shovel to a depth of 30 cm in order to expose the soil profile for sampling. The profile was scraped clean with a plastic scoop to remove any metal effect from the digging shovel. A continuous strip of soil was collected by plastic scoop over the interval of 10 to 20 cm below the top of true soil, placed in a pre-numbered zip lock baggie and placed in a 30 cm by 50 cm 2 mil plastic bag. Loess was present at nearly all sample sites and was the sample medium for most samples with a minor contribution from underlying till in some samples. Samples were kept cool until they were shipped to SGS Minerals Services in Vancouver for analyses.

In the SGS Lab, samples are not dried or prepared in any way. The MMI process includes analyses of an unscreened 50-g sample using multi-component extractants. Metals are determined by ICP-MS in the parts per billion range.

Response Ratios were calculated for Ag, As, Au, Bi, Fe, Mn, Pb, Sb, W, and Zn. Normally response ratios are calculated by dividing each value by the average of the lower quartile but because only 18 samples were collected the average from the 2019 soil sampling program was used, which could introduce errors that are believed to be small.

### Rock Chip Sampling.

Rock samples were shipped to Vancouver where they were described and broken in half with one-half given to Bureau Veritas Laboratories (BVL). BVL prepared the samples using their PRP70-550 where a 1 kg sample was crushed to  $\geq$ 70% passing 2mm followed by a 500gm sample pulverized to  $\geq$ 85% of -75 microns. A 15-gm split was analyzed by BVL's AQ201 technique using a modified aqua regia digestion with an ECP-ES/MS analysis. This provided a 37-element analyses with a suitable detection limit on critical elements including 0.5 ppb Au, 0.1 ppm Cu, 0.1 ppm Mo, 0.1 ppm Ag, 0.5 ppm As, 0.1 ppm Bi, 0.1 ppm Sb, subject to solubilities of mineral species present.

### **Results.**

MMI soil results confirmed the high response ratio, 227, from sample Y38 in the 2019 survey with three response ratios of 112, 129, and 509 collected within 100 m of Y38. The MMI results also confirmed the anomalous results from 2019 in samples collected 200 m east and yielded no anomalous results to the west thus confirming the extent of the two km anomalous zone described in the previous surveys.

All rock samples were either of quartz or more commonly of various forms of muscovite bearing carbonaceous quartzite. No outcrop was encountered in any of the soil pits although depth of pits varied from 0.5 to 1.0 m. Tills were encountered in all pits. Angular to subangular rock was collected from each pit for evaluation and later selected for assaying. Many of the quartzites contained lamellae of oxidized limonitic material and occasionally limonitic fractures. Seven quartzite samples returned gold values of 10.8 to 18.6 ppb Au, considered only weakly anomalous. Rock chip sample results are not considered high enough to explain the high Au response ratios in the MMI soil samples. Rock samples collected from a pit dug at Y38 at and close to the four very high MMI Au response ratios returned Au values of 11 and 18.6 ppb. Four samples of quartz from these pits returned gold values of 33.2, 26.8, 14.9 and 11.6 that are similarly not considered high enough to explain the high MMI Au response ratios.

Values of other metals notably As and Sb are moderately anomalous in many samples, Bi sporadically moderately anomalous, Tl weakly anomalous in some samples and W not anomalous in any samples although these four elements form widespread anomalous patterns in MMI soil samples over a three km diameter area.

Ca has a uniform consistent background of 0.02 to 0.10 %Ca. Three rock samples stand out with values of 4.85% in Y31E, 5.86% in Y31I, and 2.18% in Y38D that have corresponding Au values of 5.0, 17.6, and 18.6 ppb. This may indicate that the causative source of the high Au MMI response ratios could contain high calcium content.

Cause of the anomalous elements may be some style of epithermal gold mineralization given the anomalous Au and the number of traditional Au pathfinder elements included in the MMI anomalous suite. These pathfinder elements include As, Sb, Bi, Tl, and W.

# **CONCLUSIONS.**

Previous sampling in 2019 on the KRYPTOS claims defined two zones of anomalous gold in MMI soils, one measuring 3 km long by up to 300 m wide forming a horseshoe shape and the other one km long by up to 400 m wide all contained within a three km diameter pattern of nearly coincident anomalous As, Sb, Bi, Tl, W and other elements.

Bedrock throughout this zone is certainly metasediments, mainly carbonaceous quartzite, based on three outcrops and abundant similar angular float in soil pits. The target area lies within metasediments of the Finlayson Assemblage described on Geoscience Map 7. In 2019 no strongly mineralized samples were found although one sample, Y87, of phyllite/schist with weakly quartz bearing fractures yielded 15.9 ppb Au with moderately anomalous As and Sb but no anomalous values in any of the other pathfinder elements.

MMI soil sampling in 2020 confirmed the high Au response ratio, 227, from 2019 in sample Y38 with three samples up to 50 m away yielding Au response ratios of 112, 129, and 509. Other MMI soil samples confirmed the patterns of anomalous metals in samples from 2019.

Rock sampling in 2020 failed to provide results high enough to convincingly explain the anomalous gold patterns of MMI soil samples. Further exploration is warranted.

Epithermal gold mineralization may be present within the target area given the anomalous Au in soils and the number of traditional Au pathfinder elements included in this anomalous suite.

## **RECOMMENDATIONS.**

It is recommended that excavator trenching or percussion drilling be undertaken on and around some of the soil sample locations with the highest gold values in an attempt to find significant Au mineralization.

# **STATEMENT OF COSTS**

	TOTAL	\$ 13,674.55
Report: 10% of Sub Total		1,243.14
	Sub Total	12,431.41
Freight: Air North, Samples Whitehorse to Vancouver		166.11
YWCB 0.0459 X \$2100		96.39
Generator: 7 days @ \$10/day		70.00
Truck:Whitehorse-Stewart-Whitehorse: 900 km x \$0.60	/km	540.00
16 man-days @ \$100/man day		1600.00
Living Allowance: sample bags, food, sat phone, radios,	flagging, etc	
A Holway Aug 4-10; 7 days @ \$300/day		2100.00
G Richards Aug 3-11; 9 days @ \$500/day		4500.00
Wages:		
SGS Labs 590665 MMI samples		788.09
Bureau Veritas VANI373773 Rocks		2570.82
Geochem:		

Apply 2 years to KRYPTOS 15-60: 46 cl X 2yrs X \$100/cl-yr = \$9,200. Apply 2 years to KRYPTOS 61-80: 20 cl X 2yrs X \$100/cl-yr <u>= \$4,000</u>.

Total \$ 13,200

# STATEMENT OF QUALIFICATIONS.

I, Gordon G Richards, with business address at 6410 Holly Park Drive, B.C., V4K 4W6, do hereby certify that:

- 1. I hold a B.A.Sc. (1968) in Geology from The University of British Columbia, and an M.A.Sc. (1974) in Geology from The University of British Columbia.
- I have been practicing my profession as a geologist for over 40 years and as a consulting geological engineer from 1985 to 2018. I have work experience in western areas of the United States, Alaska, Canada, Mexico and Africa.
- 3. I have based this report on my own field work and supervision of field work by Aaron Holway during the period of August 3 to 11 and on the results generated by that field work.

Respectfully submitted,

Gordon G Richards





Figure 3. Local Geology of Mega Project within Kryptos Claims. Au-As-Sb-Bi-W anomalous zone was the target that was evaluated. It occurs within schists, quartzites, and phyllite of the DMF1 unit east of a northerly fault. Porphyry Cu-Au targets were not examined. See Table 2 for Legend.



Маяс

CONTRACT AND A TRACTACTION AND A TRACTACIAN TO CONTRACT DID HOLD		
eten and narts of northern Carmacks. Yukon: Geological Survey of Canada.	enOoM mot	səmunos
igure 🐔 taken from: Ryan, J.,, Colpron, M., and Hayward, N., 2010. Geology,	I tof bnesd	Table 1.
Quartzite to quartz-mica schiat; banded to massive, grey to white in colour; locally conglomeratic; commonly contains beds of micaceous quartz arenite; clastic in origin; quartz-muscovite-biotite schiat is possibly derived from siliceous siltstone; commonly finely interlayered with garnet-metapelite.	rsoa	
Amphibolite scrints to game-simprinolite; metabasite; usually gamet-horneblende-plagioclase or horneblende-plagioclase; with local chorite-biotite; probably derived from mafic volcanic to volcaniclastic rocks; some layers that are internally homogeneous may be mafic sills; more intermediate varieties can have rosettes of decussate; larger hornblende.	6DS3	
	I VIE DENO	
Carbonaceous quartzite to mica-quartz schiat; black to white quartzite, with schiat and garnet schiat interlayers; and rare black phyllite; possibly equivalent to Nasina formation, or simply a carbonaceous member of the Snowcap assemblage.	DMF1	
of the garnet-amphibolites assigned to the Snowce Amonto semiclastic to the stretching lineation; commonly medium green; possibly lower grade equivalent of the garnet-amphibolites assigned to the Snowcep Assemblage.	DMF2	
rinisyson essembiager (DMF1, DMF2) Greenstone - greenschist (acies metabasite: chlorite-actinolite schist:		
	DIRSJOV679M	
Intermediate to mafic granitoid (tonalite to diorite) sheets; intermediate to dark colour; homogeneous to layered.	RSIM	×
Monzogranite to granodiorite; equigranular; pink to orange; generally biotite-bearing (after hornblende?); homogeneous to layered.	RepM	
Simpson Kange suite (MgSR, MiSR, MagSR)		IAN,
	A francisco and	Δ
rake taut; easity contraed with undetormed post- I riassic intrusions.		
phenocrysts common in eastern extent; slightly foliated adjacent to Willow blebby to porphyritic smokey quarts; fresh magmatic hornblende and K-feldspar Keid Lake batholith: polyphase; undeformed to weakly foliated adjacent to Willow	равМ	
<b>Кеіd Lakes complex (</b> Мдbяר, Мдяר, Мvяר)		
NAI991221	SSIM YJAAA	
Intermediate to mafic, light-green, pyrite-chlorite schist; commonly exhibits a pitted surface indicative of coarse pyrite cubes having weathered out; primary volcanic textures locally preserved.	PK1	
Metafelsite, commonly porphyritic or augen-textured; possibly derived from felsic volcanic rocks or hypabyssal intrusions; locally derived from equigranular to augen monzogranite; locally exhibits decussate amphiboles pseudomorphed to chlorite-biotite; local coarse porphyroblastic garnet. Intermediate to mafic, light-green, pyrite-chlorite schist; commonly exhibits a pitted surface indicative of coarse pyrite cubes having weathered out; primary volcanic textures locally preserved.	РК1 РК2	50,
Klondike Schist (PK1, Pk2) Metafelsite, commonly porphyritic or augen-textured; possibly derived from felsic volcanic rocks or hypabyssal intrusions; locally derived from equigranular to augen monzogranite; locally exhibits decussate amphiboles pseudomorphed to chlorite-biotite; local coarse porphyroblastic garnet. Intermediate to mafic, light-green, pyrite-chlorite schist; commonly exhibits a pitted surface indicative of coarse pyrite cubes having weathered out; primary volcanic textures locally preserved.	PK1 PK2	50,
Sulphur Creek suite: quartz and K-feldspar porphyritic to augen monzogranite; strain varies from moderately foliated to gneissic (including porphyroclastic straight gneiss); biotite bearing; locally is the protolith to felsic Klondike Schist. Metafelsite, commonly porphyritic or augen-textured; possibly derived from felsic volcanic rocks or hypabyssal intrusions; locally derived from seculgranular to augen monzogranite; local coarse porphyroblastic garnet. Intermediate to mafic, light-green, pyrite-chlorite schist; commonly exhibits a pitted surface indicative of coarse pyrite cubes having weathered out; primary volcanic textures locally preserved.	ЬКJ ЬKS Ьâsc	50,
and metavolcanic rocks of the Klondike arc Sulphur Creek suite: quartz and K-feldspar porphyritic to augen monzogranite; straight gneiss); biotite bearing; locally is the protolith to felsic Klondike Schist. Metafelsite, commonly porphyritic or augen-textured; possibly derived from felsic volcanic rocks or hypabyssal intrusions; locally derived from equigranular to augen monzogranite; locally exhibits decusaate amphiboles pseudomorphed to chlorite-biotite; local coarse porphyroblastic garnet. Intermediate to mafic, light-green, pyrite-chlorite schist; commonly exhibits a pitted surface indicative of coarse pyrite cubes having weathered out; primary volcanic textures locally preserved.	PgSC PgSC	20,
and metavolcanic rocks of the Klondike arc Sulphur Creek suite: quartz and K-feldspar porphyritic to augen monzogranite; straight gneiss); biotite bearing; locally is the protolith to felsic Klondike Schist. Klondike Schist (Pk1, Pk2) Metafelsite, commonly porphyritic or augen-textured; possibly derived from felsic volcanic rocks or hypabyssal intrusions; locally derived from equigranular to augen monzogranite; locally exhibits decussate amphiboles pseudomorphed to chlorite-biotite; local coarse porphyroblastic garnet. Intermediate to mafic, light-green, pyrite-chlorite schist; commonly exhibits a pitted surface indicative of coarse pyrite cubes having weathered out; primary volcanic textures locally preserved.	PERMIAN Metaplutonic PgsC	50,
underformed but locally foliated plutons and/or dykes. and metavolcanic rocks of the Klondike arc Sulphur Creek suite: quartz and K-feldspar porphyritic to augen monzogranite; atraipht gneiss); biotite bearing; locally is the protolith to felsic Klondike Schiat (Pk1, Pk2) Metafelsite, commonly porphyritic or augen-textured; possibly derived from felsic volcanic rocks or hypabyssal intrusions; locally derived from geuugranular to augen monzogranite; local oranse porphyroblastic decusate amphiboles pseudomorphed to chlorite-biotite; local coarse porphyroblastic garnet. Intermediate to mafic, light-green, pyrite-chlorite schiat; commonly exhibits a pitted surface indicative of coarse porphyroblastic garnet. intermediate to mafic, light-green, pyrite-chlorite schiat; commonly out; primary volcanic textures locally preserved.	PERMIAN Metaplutonic PgSC PRZ	50,
Aishihik suite: granodionite to monzogranite (± quartz monzonite and quartz monzodiorite); commonly K-feldspar porphyntic and hornblende-bearing; common biotite ± chlorite alteration with secondary epidote; prominent magmatic epidote; intrudes Stikinia and Yukon Tanana terranes; generally and metavolcanic rocks of the Klondike arc Sulphur Creek suite; quartz and K-feldspar porphyritic to augen monzogranite; denisin varies from moderately foliated plutons and/or dykes. Klondike Schist (Pk1, Pk2) Metafelsite, commonly porphyritic or augen-textured; possibly derived from felsic volcanic rocks or hypabyssal intrusions; locally atribits decusate amphiboles equigranular to augen monzogranite; local coarse porphyroblastic gamet. pseudomorphed to chlorite-biotite; local coarse porphyroblastic gamet. Intermediate to mafic, light-green, pyrite-chlorite schist; commonly exhibits a pitted surface indicative of coarse porphyroblastic gamet. intermediate to mafic, light-green, pyrite-chlorite schist; commonly exhibits a pitted surface indicative of coarse porphyroblastic gamet. intermediate to mafic, light-green, pyrite-chlorite schist; commonly exhibits a pitted surface indicative of coarse pyrite cubes having weathered exhibits a pitted surface indicative of coarse pyrite cubes having weathered exhibits a pitted surface indicative of coarse pyrite cubes having weathered exhibits a pitted surface indicative of coarse pyrite cubes having weathered exhibits a pitted surface indicative of coarse pyrite cubes having weathered exhibits a pitted surface indicative of coarse pyrite cubes having weathered exhibits a pitted surface indicative of coarse pyrite cubes having weathered exhibits a pitted surface indicative of coarse pyrite cubes having weathered exhibits a pitted surface indicative of coarse pyrite cubes having weathered exhibits a pitted surface indicative of coarse pyrite cubes having weathered exhibits a pitted surface indicative of coarse pyrite cubes having weathered exhibits a pitted surface indicative of coarse py	EJgA PERMIAN PgsC PgsC	STIKINIA/QUESNELIA









То **GORDON RICHARDS GORDON RICHARDS** 6410 HOLLY PARK DR DELTA V4K 4W6 BC CANADA

Order Number	PO:	Date Received	12-Aug-2020
Project	Mega	Date Analysed	12-Aug-2020 - 21-Aug-2020
Submission Number	*BBY* MEGA/ 17 (MMI)	Date Completed	22-Aug-2020
Number of Samples	17	SGS Order Number	BBM20-03991

#### Methods Summary

Number of Sample Method Code	Description
17 G_LOG	Sample Registration Fee
17 G_WGH_KG	Weight of samples received
17 GE_MMIM	Mobile Metal ION standard package, ICP-MS

Authorised Signatory

John Chiang Laboratory Operations Manager

This document is issued by the Company under its General Conditions of Service accessible at https://www.sgs.com/en/Terms-and-Conditions.aspx. Attention is drawn to the limitation of liability, indemnification and jurisdiction issues defined therein. Any holder of this document is advised that information contained hereon reflects the Company's findings at the time of its intervention only and within the limits of Client's instructions, if any. The Company's sole responsibility is to its Client and this document does not exonerate parties to a transaction from exercising all their rights and obligations under the transaction documents. Any unauthorized alteration, forgery or falsification of the content or appearance of this document is unlawful and offenders may be prosecuted to the fullest extent of the law.

WARNING: The sample(s) to which the findings recorded herein (the "Findings") relate was(were) drawn and / or provided by the Client or by a third party acting at the Client's direction. The Findings constitute no warranty of the sample's representativeness of any goods and strictly relate to the sample(s). The Company accepts no liability with regard to the origin or source from which the sample(s) is/are said to be extracted. The findings report on the samples provided by the client and are not intended for commercial or contractual settlement puposes.

> - not analysed -- element not determined | I.S. insufficient sample | L.N.R. listed not received

22-Aug-2020 2:54PM BBM\_U0003273158

Page 1 of 10

MIN-M\_COA\_ROW-Last Modified Date: 05-Nov-2019

SGS Canada CA MIN Burnaby, BC 3260 Production Way, Burnaby, BC V5A 4W4 Burnaby CANADA t +1 (604) 638 2349 f

www.sgs.com

Member of the SGS Group (SGS SA)



### **ANALYSIS REPORT BBM20-03991**

Element	Wtkg	Ag	AI	As	Au	Ва
Method	G_WGH_KG	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM
Lower Limit	0.01	0.5	1	10	0.1	10
Upper Limit						
Unit	kg	ppb	ppm m / m	ppb	ppb	ppb
W1	0.17	47.8	27	<10	0.5	13600
W2	0.21	9.4	254	70	0.2	9080
W3	0.15	20.4	98	50	0.5	14500
W4	0.17	50.7	12	<10	0.3	2180
W5	0.22	43.8	19	120	15.0	3070
W6	0.21	33.3	27	150	17.3	1760
W7	0.18	37.2	17	90	68.1	2160
W8	0.32	15.3	33	40	0.3	1160
W9	0.31	69.5	18	10	1.7	13700
W10	0.15	81.6	144	50	0.9	13000
W11	0.32	35.8	192	100	1.1	8440
W12	0.23	25.3	148	440	3.5	6600
W13	0.13	10.1	145	10	1.6	7870
W14	0.23	37.0	86	40	0.4	7430
W15	0.32	18.9	97	130	0.3	8890
W17	0.18	56.8	153	10	1.3	8940
W18	0.18	33.8	282	50	0.9	5460
*Rep W8	-	13.8	31	40	0.2	1160
*Rep W18	-	35.3	278	40	0.9	5400
*Std AMIS0169	-	7.0	51	10	0.3	800
*Blk BLANK	-	<0.5	<1	<10	<0.1	<10

Element	Bi	Са	Cd	Ce	Со	Cr
Method	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM
Lower Limit	0.5	2	1	2	1	100
Upper Limit						
Unit	ppb	ppm m / m	ppb	ppb	ppb	ppb
W1	<0.5	554	26	272	54	<100
W2	2.8	59	11	410	92	200

- not analysed | -- element not determined | I.S. insufficient sample | L.N.R. listed not received

22-Aug-2020 2:54PM BBM\_U0003273158

Page 2 of 10

MIN-M\_COA\_ROW-Last Modified Date: 05-Nov-2019



### ANALYSIS REPORT BBM20-03991

Element Method Lower Limit Upper Limit	Bi GE_MMIM 0.5 	Ca GE_MMIM 2 	Cd GE_MMIM 1 	Ce GE_MMIM 2 	Co GE_MMIM 1 	Cr GE_MMIM 100 
Unit	ppb	ppm m / m	ppb	ppb	ppb	ppb
W3	1.4	324	4	345	62	100
W4	<0.5	423	7	76	15	<100
W5	<0.5	476	6	65	39	<100
W6	<0.5	423	2	119	19	<100
W7	<0.5	515	3	56	11	<100
W8	<0.5	382	10	43	19	<100
W9	<0.5	342	2	15	6	<100
W10	<0.5	290	18	360	59	100
W11	1.1	112	13	1050	142	100
W12	3.7	91	8	300	86	100
W13	<0.5	221	29	210	207	<100
W14	0.7	228	9	361	149	<100
W15	2.8	161	11	346	115	100
W17	<0.5	275	8	298	156	<100
W18	0.8	18	42	312	73	<100
*Rep W8	<0.5	380	10	42	19	<100
*Rep W18	0.8	17	45	296	78	<100
*Std AMIS0169	<0.5	31	1	580	72	<100
*Blk BLANK	<0.5	<2	<1	<2	<1	<100

Element	Cs	Cu	Dy	Er	Eu	Fe
Method	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM
Lower Limit	0.2	10	0.5	0.2	0.2	1
Upper Limit						
Unit	ppb	ppb	ppb	ppb	ppb	ppm m / m
W1	0.5	2380	38.3	19.6	12.5	27
W2	3.5	450	42.4	17.5	13.1	231
W3	1.4	690	49.9	26.1	15.5	116
W4	1.0	1160	30.7	22.1	8.2	34

- not analysed | -- element not determined | I.S. insufficient sample | L.N.R. listed not received

22-Aug-2020 2:54PM BBM\_U0003273158

Page 3 of 10

MIN-M\_COA\_ROW-Last Modified Date: 05-Nov-2019

SGS Canada CA MIN Burnaby, BC 3260 Production Way, Burnaby, BC V5A 4W4 Burnaby CANADA t +1 (604) 638 2349 f

www.sgs.com



### **ANALYSIS REPORT BBM20-03991**

Element Method Lower Limit Upper Limit	Cs GE_MMIM 0.2 	Cu GE_MMIM 10 	Dy GE_MMIM 0.5 	Er GE_MMIM 0.2 	Eu GE_MMIM 0.2 	Fe GE_MMIM 1 
Unit	ddd	ppb 1110	45 O	hbn	ppb 4.0	ppm m / m
VV5	0.5	1140	15.0	9.0	4.0	21
W6	2.1	580	27.8	15.9	6.5	47
W7	0.6	900	14.6	8.4	3.7	27
W8	0.7	490	7.8	4.5	2.0	53
W9	1.2	270	7.0	3.6	3.1	13
W10	4.1	290	83.3	45.5	21.7	119
W11	5.1	730	114	58.4	30.6	121
W12	8.2	370	44.6	25.1	11.2	267
W13	1.2	800	179	109	21.4	117
W14	2.8	550	53.4	30.6	14.8	114
W15	7.2	700	45.7	26.2	11.6	174
W17	0.7	860	169	103	27.8	182
W18	3.4	430	69.0	34.4	13.8	195
*Rep W8	0.6	480	7.2	4.2	1.9	55
*Rep W18	2.8	440	68.3	35.2	13.5	181
*Std AMIS0169	7.0	2950	22.3	10.0	8.5	35
*Blk BLANK	<0.2	<10	<0.5	<0.2	<0.2	<1

Element	Ga	Gd	Hg	In	к	La
Method	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM
Lower Limit	0.5	0.5	1	0.1	0.5	1
Upper Limit						
Unit	ppb	ppb	ppb	ppb	ppm m / m	ppb
W1	1.5	51.2	<1	<0.1	3.6	89
W2	20.7	47.3	<1	0.3	10.4	223
W3	8.4	60.9	<1	<0.1	3.4	208
W4	1.4	32.7	<1	<0.1	5.2	95
W5	0.8	16.6	<1	<0.1	6.6	31
W6	2.1	28.9	<1	<0.1	6.9	70

- not analysed | -- element not determined | I.S. insufficient sample | L.N.R. listed not received

22-Aug-2020 2:54PM BBM\_U0003273158

MIN-M\_COA\_ROW-Last Modified Date: 05-Nov-2019

SGS Canada CA MIN Burnaby, BC 3260 Production Way, Burnaby, BC V5A 4W4 Burnaby CANADA t +1 (604) 638 2349 f

www.sgs.com



### **ANALYSIS REPORT BBM20-03991**

Element Method Lower Limit Upper Limit Unit	Ga GE_MMIM 0.5  ppb	Gd GE_MMIM 0.5  ppb	Hg GE_MMIM 1  ppb	In GE_MMIM 0.1  ppb	K GE_MMIM 0.5  ppm m / m	La GE_MMIM 1  ppb
W7	0.6	15.9	<1	<0.1	4.7	30
W8	1.1	8.2	<1	<0.1	4.4	17
W9	0.6	8.5	<1	<0.1	4.2	14
W10	6.3	86.6	<1	0.2	19.2	190
W11	12.4	130	<1	0.2	11.1	568
W12	10.2	48.6	<1	0.2	17.6	169
W13	1.4	113	<1	0.1	13.6	84
W14	3.5	61.2	<1	<0.1	7.4	196
W15	8.0	51.1	<1	0.2	10.8	179
W17	1.8	140	<1	0.2	6.2	136
W18	7.3	62.5	<1	0.3	11.7	157
*Rep W8	1.2	7.7	<1	<0.1	4.4	16
*Rep W18	6.5	60.5	<1	0.3	11.3	146
*Std AMIS0169	9.2	32.7	<1	<0.1	40.3	338
*Blk BLANK	<0.5	<0.5	<1	<0.1	<0.5	<1

Element	Li	Mg	Mn	Мо	Nb	Nd
Method	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM
Lower Limit	1	0.5	100	2	0.5	1
Upper Limit						
Unit	ppb	ppm m / m	ppb	ppb	ppb	ppb
W1	8	46.9	2000	5	<0.5	163
W2	21	8.7	2200	6	8.1	218
W3	8	26.6	2300	5	2.5	235
W4	12	9.5	1900	2	<0.5	131
W5	2	24.0	1500	<2	<0.5	53
W6	16	24.1	800	<2	<0.5	101
W7	4	27.4	200	<2	<0.5	50
W8	6	13.9	1500	3	<0.5	28

- not analysed | -- element not determined | I.S. insufficient sample | L.N.R. listed not received

22-Aug-2020 2:54PM BBM\_U0003273158

MIN-M\_COA\_ROW-Last Modified Date: 05-Nov-2019



### **ANALYSIS REPORT BBM20-03991**

Element Method Lower Limit Upper Limit Unit	Li GE_MMIM 1  ppb	Mg GE_MMIM 0.5  ppm m / m	Mn GE_MMIM 100  ppb	Mo GE_MMIM 2  ppb	Nb GE_MMIM 0.5  ppb	Nd GE_MMIM 1  ppb
W9	6	10.0	500	3	<0.5	27
W10	21	29.9	5700	4	1.3	287
W11	19	15.7	6100	5	3.2	593
W12	23	12.5	5400	10	4.4	180
W13	<1	45.3	5800	<2	<0.5	219
W14	5	32.3	4500	4	1.1	258
W15	20	22.8	5100	9	3.9	202
W17	<1	52.1	1100	<2	<0.5	299
W18	4	6.6	2800	3	2.7	224
*Rep W8	4	13.3	1400	2	<0.5	28
*Rep W18	4	6.7	2800	3	2.4	200
*Std AMIS0169	<1	29.9	3000	3	2.6	295
*Blk BLANK	<1	<0.5	<100	<2	<0.5	<1

Element Method	Ni GE MMIM	P GE MMIM	Pb GE MMIM	Pd GE MMIM	Pr GE MMIM	Pt GE MMIM
Lower Limit	5	0.1	5	1	0.5	0.1
Upper Limit						
Unit	ppb	ppm m / m	ppb	ppb	ppb	ppb
W1	1040	0.2	37	<1	33.6	<0.1
W2	235	8.7	703	<1	53.7	<0.1
W3	298	1.0	259	<1	53.5	<0.1
W4	301	0.3	37	<1	28.9	<0.1
W5	705	0.2	21	<1	11.1	<0.1
W6	334	0.4	29	<1	22.1	<0.1
W7	431	0.3	20	<1	10.5	<0.1
W8	455	0.8	58	<1	6.3	<0.1
W9	101	0.4	11	<1	5.5	<0.1
W10	514	3.3	154	<1	63.4	<0.1

- not analysed | -- element not determined | I.S. insufficient sample | L.N.R. listed not received

22-Aug-2020 2:54PM BBM\_U0003273158

Page 6 of 10

MIN-M\_COA\_ROW-Last Modified Date: 05-Nov-2019



### **ANALYSIS REPORT BBM20-03991**

Element Method Lower Limit Upper Limit Unit	Ni GE_MMIM 5  ppb	P GE_MMIM 0.1  ppm m / m	Pb GE_MMIM 5  ppb	Pd GE_MMIM 1  ppb	Pr GE_MMIM 0.5  ppb	Pt GE_MMIM 0.1  ppb
W11	524	5.4	528	<1	148	<0.1
W12	317	8.3	681	<1	43.2	<0.1
W13	1120	0.6	1060	<1	40.2	<0.1
W14	383	2.4	462	<1	58.4	<0.1
W15	390	7.0	490	<1	48.7	<0.1
W17	1810	0.7	924	<1	57.2	<0.1
W18	255	4.6	579	<1	50.4	<0.1
*Rep W8	442	0.8	55	<1	5.9	<0.1
*Rep W18	276	4.3	590	<1	46.2	<0.1
*Std AMIS0169	331	2.5	76	<1	80.2	<0.1
*Blk BLANK	<5	<0.1	<5	<1	<0.5	<0.1

Element Method Lower Limit Upper Limit Unit	Rb GE_MMIM 1  ppb	Sb GE_MMIM 0.5  ppb	Sc GE_MMIM 5  ppb	Sm GE_MMIM 1  ppb	Sn GE_MMIM 1  ppb	Sr GE_MMIM 10  ppb
W1	19	1.9	28	40	<1	2300
W2	77	6.4	55	44	1	240
W3	56	10.6	43	50	<1	1000
W4	27	2.0	30	26	<1	1360
W5	15	31.6	14	12	<1	1160
W6	29	81.4	22	22	<1	940
W7	16	26.2	16	12	<1	1310
W8	23	8.9	9	6	<1	1110
W9	18	3.9	16	7	<1	960
W10	71	12.6	128	67	<1	820
W11	111	9.8	109	116	<1	440
W12	106	26.7	56	38	<1	380

- not analysed | -- element not determined | I.S. insufficient sample | L.N.R. listed not received

22-Aug-2020 2:54PM BBM\_U0003273158

Page 7 of 10

MIN-M\_COA\_ROW-Last Modified Date: 05-Nov-2019



PO:
Mega
*BBY* MEGA/ 17 (MMI)
17

Element Method Lower Limit Upper Limit	Rb GE_MMIM 1 	Sb GE_MMIM 0.5 	Sc GE_MMIM 5 	Sm GE_MMIM 1 	Sn GE_MMIM 1 	Sr GE_MMIM 10 
Unit	ppb	ppb	ppb	ppb	ppb	ppb
W13	34	3.0	72	62	<1	1510
W14	68	6.6	43	53	<1	900
W15	103	14.2	73	42	<1	690
W17	80	2.4	86	86	<1	1880
W18	108	4.2	45	48	<1	200
*Rep W8	23	9.7	9	6	<1	1110
*Rep W18	101	3.7	42	46	<1	200
*Std AMIS0169	212	0.6	44	44	2	80
*BIk BLANK	<1	<0.5	<5	<1	<1	<10

Element Method Lower Limit Upper Limit Unit	Ta GE_MMIM 1  ppb	Tb GE_MMIM 0.1  ppb	Te GE_MMIM 10  ppb	Th GE_MMIM 0.5  ppb	Ti GE_MMIM 10  ppb	TI GE_MMIM 0.1  ppb
W1	<1	6.6	<10	55.5	40	<0.1
W2	<1	7.1	<10	76.2	2320	0.4
W3	<1	8.2	<10	81.7	820	0.3
W4	<1	4.5	<10	49.7	70	<0.1
W5	<1	2.4	<10	53.1	20	<0.1
W6	<1	4.2	<10	144	60	<0.1
W7	<1	2.3	<10	93.2	10	<0.1
W8	<1	1.2	<10	83.9	70	<0.1
W9	<1	1.1	<10	25.6	40	<0.1
W10	<1	12.6	<10	54.7	420	0.1
W11	<1	18.3	<10	105	1180	0.3
W12	<1	7.0	<10	113	1120	0.4
W13	<1	22.5	<10	41.3	20	0.1
W14	<1	8.4	<10	74.1	250	0.1

- not analysed | -- element not determined | I.S. insufficient sample | L.N.R. listed not received

22-Aug-2020 2:54PM BBM\_U0003273158

Page 8 of 10

MIN-M\_COA\_ROW-Last Modified Date: 05-Nov-2019



Order Number	PO:
Project	Mega
Submission Number	*BBY* MEGA/ 17 (MMI)
Number of Samples	17
Number of Samples	17

Element	Та	Tb	Те	Th	Ti	TI
Method	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM	GE_MMIM
Lower Limit	1	0.1	10	0.5	10	0.1
Upper Limit						
Unit	ppb	ppb	ppb	ppb	ppb	ppb
W15	<1	7.3	<10	126	770	0.4
W17	<1	23.9	<10	70.8	30	0.2
W18	<1	10.0	<10	47.0	610	0.2
*Rep W8	<1	1.1	<10	85.1	70	<0.1
*Rep W18	<1	10.0	<10	43.8	520	0.2
*Std AMIS0169	<1	4.0	<10	55.5	390	1.1
*Blk BLANK	<1	<0.1	<10	<0.5	<10	<0.1

Element Method Lower Limit Upper Limit	U GE_MMIM 0.5 	W GE_MMIM 0.5 	Y GE_MMIM 1 	Yb GE_MMIM 0.2 	Zn GE_MMIM 10 	Zr GE_MMIM 2 
Unit	ppb	ppb	ррь	ppb	ppb	ppb
W1	49.8	<0.5	216	14.3	160	56
W2	22.4	1.5	173	11.2	380	107
W3	33.3	0.9	277	19.1	150	72
W4	24.6	<0.5	205	20.8	60	22
W5	43.1	<0.5	90	7.1	70	11
W6	53.8	<0.5	165	14.7	70	25
W7	45.9	<0.5	82	7.0	40	13
W8	39.3	<0.5	41	3.9	200	23
W9	16.4	<0.5	34	3.1	30	39
W10	37.9	0.6	439	36.2	240	71
W11	49.6	1.1	567	41.3	300	151
W12	25.7	1.0	235	19.3	560	256
W13	78.6	<0.5	1060	72.3	1110	41
W14	42.3	<0.5	268	24.3	350	115
W15	38.6	0.7	227	21.7	450	264
W17	88.4	<0.5	975	75.4	150	71

- not analysed | -- element not determined | I.S. insufficient sample | L.N.R. listed not received

22-Aug-2020 2:54PM BBM\_U0003273158

Page 9 of 10

MIN-M\_COA\_ROW-Last Modified Date: 05-Nov-2019



Order Number	PO:
Project	Mega
Submission Number	*BBY* MEGA/ 17 (MMI)
Number of Samples	17

Element Method Lower Limit Upper Limit Unit	U GE_MMIM 0.5  ppb	W GE_MMIM 0.5 	Y GE_MMIM 1  ppb	Yb GE_MMIM 0.2  ppb	Zn GE_MMIM 10  ppb	Zr GE_MMIM 2  ppb
W18	24.9	<0.5	321	21.4	470	79
*Rep W8	36.9	<0.5	40	3.8	200	23
*Rep W18	24.1	<0.5	313	21.2	490	70
*Std AMIS0169	18.6	1.1	87	7.0	150	37
*Blk BLANK	<0.5	<0.5	<1	<0.2	<10	<2

- not analysed | -- element not determined | I.S. insufficient sample | L.N.R. listed not received

22-Aug-2020 2:54PM BBM\_U0003273158

Page 10 of 10

MIN-M\_COA\_ROW-Last Modified Date: 05-Nov-2019

SGS Canada CA MIN Burnaby, BC 3260 Production Way, Burnaby, BC V5A 4W4 Burnaby CANADA **t** +1 (604) 638 2349 **f** 

www.sgs.com

Member of the SGS Group (SGS SA)