



105K/2 NE  
YUKON  
SCALE 1:25 000

1:250,000  
0 1000 2000 3000 4000  
1:250,000


CONTOUR INTERVAL 100 FEET  
Elevations in feet above Mean Sea Level  
North American Datum 1983  
Transverse Mercator Projection

True north  
Magnetic north  
Magnetic declination  
Annual change decreasing 4.1"



Topographic data provided by  
SURVEY AND MAPPING BRANCH  
DEPARTMENT OF ENERGY, MINES  
AND TECHNICAL SERVICES  
Copyright law reserves the Queen  
in Right of Canada  
ONE THOUSAND METRE  
Universal Transverse Mercator Grid  
ZONE 8

A



LEGEND

#### INTRUSIVE ROCKS

##### EOCENE

quartz-feldspar porphyry  
white-weathering, aphanitic to fine-grained, locally flow-banded quartz-feldspar porphyry; commonly contains phenocrysts of smoky grey quartz, biotite and white feldspar

##### CRETACEOUS

granite to granodiorite undifferentiated  
grey, resistant, generally medium- to coarse-grained, locally megacrystic, undifferentiated Tay River plutonic suite or Anvil plutonic suite granite to granodiorite

Tay River plutonic suite  
Orchay phase - biotite + hornblende granite to granodiorite

Anvil plutonic suite  
Mount Mye phase - biotite-muscovite granite; locally foliated

##### PERMIAN?

gabbro, harzburgite, serpentinite  
mafic and ultramafic intrusive rocks; locally extensively sheared and serpentinitized  
Ps - serpentinite; Phz - harzburgite; Pg - gabbro

##### ORDOVICIAN-SILURIAN

gabbro  
dark green, locally magnetic, coarse- to fine-grained, massive to foliated gabbro; subvolcanic dykes and sills to Menzies Creek basalts (OSMCb); enclosing phyllites locally display thin contact metamorphic aureoles  
pyroxenite  
dark green, locally magnetic, coarse-grained, massive to foliated, variably serpentinitized pyroxenite; subvolcanic dykes and sills to Menzies Creek basalts (OSMCb); enclosing phyllites locally display thin contact metamorphic aureoles

##### LAYERED ROCKS

##### YUKON-TANANA TERRANE

##### TRIASSIC

Faro Peak formation  
resistant, massive, polymictic conglomerate; clasts include quartzite, chert, limestone and serpentinite; matrix contains detrital muscovite  
dark grey carbonaceous, locally calcareous shale or siltstone interbedded with medium to dark grey, fine-grained limestone  
interbedded cherty argillite, chert, sandstone and mafic greywacke or conglomerate  
massive, dark green, fine-grained to aphanitic basalt; may be equivalent to Anvil Range Group basalt

##### PALEOZOIC

metasedimentary and metavolcanic rocks  
medium to dark grey, locally gritty, muscovitic meta-quartzite to quartzose schist; contains bands of greywacke, gabbro, phyllite; rarely contains eclogite lenses  
grey to tan, massive limestone or dolostone  
medium to dark olive green, chloritic phyllite to amphibolite; locally displays relict equigranular igneous texture; locally includes ultramafic rocks and/or eclogite (Pggs)  
felsic orthogneiss or paragneiss

##### SLIDE MOUNTAIN TERRANE

##### PERMIAN

Campbell Range formation  
Epizonal, locally hercynitic, dark green, resistant, massive, poorly foliated basalt or brecciated basalt; contains lesser grey, green, red and black bedded chert, and pale green epivolcaniclastic sandstone or conglomerate

##### EARLY CARBONIFEROUS-PERMIAN

Rose Mountain formation  
pale green, tan-weathering, bedded phyllitic chert interbedded with lesser maroon chert and argillite, especially near top of unit; also contains minor black bedded chert, black chert-pebble conglomerate, siltstone, limestone and argillite

##### DEVONIAN-PERMIAN

undivided Rose Mountain formation and Mount Aho formation  
dark grey to black, pale green, and maroon noncalcareous argillite and bedded chert with lesser siltstone, sandstone, chert-pebble conglomerate and limestone

##### DEVONIAN-EARLY CARBONIFEROUS

Mount Aho formation  
silvery cream, tan-weathering, bedded phyllitic chert with light grey barite beds  
dark grey to black, noncalcareous, siliceous argillite and bedded chert with lesser siltstone, sandstone, chert-pebble conglomerate and limestone  
pale green, noncalcareous argillite and bedded chert with lesser shale chip and siltstone breccia, grey sandstone and chert-pebble conglomerate; locally contains maroon argillite and bedded chert

##### ANCIENT NORTH AMERICA


##### DEVONIAN-EARLY CARBONIFEROUS

Earn Group  
dark grey to black, noncalcareous, siliceous argillite with lesser siltstone, sandstone, chert-pebble conglomerate and limestone

##### SILURIAN

siltstone  
dark grey to black, platy, tan-weathering, thinly laminated, dolomitic siltstone  
tan- to orange-weathering, dolomitic, bioturbated, silty mudstone

##### COMPILATION SOURCES



B



#### ORDOVICIAN-DEVONIAN

quartz sandstone and dolostone  
Massive, medium-grained, quartz sandstone interbedded with pale tan-weathering limestone or dolostone

Road River Group  
Duo Lake Formation  
dark grey to black, gabbroic argillite; contains lesser medium to pale grey siltstone and fine sandstone, medium grey limestone and basalt flows


##### Menzies Creek formation

undivided dark grey green, foliated basalt; includes massive and pillowed, locally amygdaloidal flows and heterolithic or monolithic breccias with lesser limestone, argillite and tuff  
dark grey green, locally amygdaloidal, massive and pillowed basalt with minor monolithic basalt breccia, volcanoclastic sandstone, siltstone and tuff  
dark grey green, monolithic basalt breccia with lesser volcanoclastic sandstone, siltstone and tuff, and massive and pillowed flows  
grey to off-white limestone locally interbedded with orange-weathering dolostone

#### CAMBRIAN-ORDOVICIAN

Vangorda formation  
soft, silvery grey, carbonaceous phyllite with lesser medium crystalline, grey marble, dark grey to black phyllite and dark green gabbro dykes and sills (OSg)  
pale green and dark purplish brown, thinly banded calc-silicate rock with lesser black schist, marble and dark green gabbro dykes and sills (OSg)  
black, locally calcareous, carbonaceous phyllite or schist; commonly contains thin quartzose siltstone interbeds, interbedded with dark green gabbro dykes and sills (OSg)  
pale to dark grey, foliated marble

#### SYMBOLS



#### UPPER PROTEROZOIC-CAMBRIAN

Mount Mye formation  
brownish grey, noncalcareous, pervasively foliated phyllite; locally indistinctly bedded; contains minor siltstone, marble, calc-silicate rock, carbonaceous phyllite and dark green gabbro dykes and sills (OSg)  
brownish grey, noncalcareous, pervasively foliated muscovite-biotite schist; may contain sauroite, garnet, andalusite, or kyanite; locally indistinctly bedded; contains minor siltstone, marble, calc-silicate rock, carbonaceous phyllite and dark green gabbro dykes and sills (OSg)  
pale green and dark purplish brown, thinly banded calc-silicate rock; contains marble and siliceous marble beds and dark green gabbro dykes and sills (OSg); lithologically similar to Vangorda calc-silicate rock  
dark to pale grey, medium crystalline marble; typically contains abundant boudins of calc-silicate rock and/or quartz; locally contains coarsely crystalline garnet-pyroxene skarn  
black phyllite or schist; locally contains lenses and beds of black carbonaceous limestone and dark green gabbro dykes and sills (OSg)

MINERAL OCCURRENCES Yukon MINFILE				
105K 27	★	SPUR	Exploration Target	
105K 38	★	VALRAY	Exploration Target	
Dekker, R. 2003. Yukon MINFILE - a database of mineral occurrences. Yukon Geological Survey, CD-ROM.				

ISOTOPIC AGE DATES					
Sample	Date	System	Mineral	Comments	Ref
GSC80-82	88.349.9 Ma	K-Ar	biotite	intrusion cooling age	(3)
GSC80-86	97.823.4 Ma	K-Ar	hornblende	intrusion cooling age	(3)

#### REFERENCES

- Gorday, S.P., 1990. Geology of Tenas Creek (105K/1), Swim Lakes (105K/2), and Faro (105K/3) map areas, Yukon Territory. Geological Survey of Canada, Open File 2040 (1:50 000 scale).
- Gorday, S.P. and Innes, S.E.B., 1987. Geology, Stikton Lake and Tay River map areas, Yukon Territory. Geological Survey of Canada, Map 19-1987 (3 sheets, 1:250 000 scale).
- Hunt, P.A. and Roddick, J.C., 1991. A compilation of K-Ar ages, Report 20. Geological Survey of Canada, Paper 89-2, p. 113-145.
- Jennings, D.S., Jason, G.A., Hanson, D.J. and Francis, J.P., 1978. Geology Anvil District Map Area. Unpublished Cyprus Anvil Mining Corporation internal company report (1:100 000 scale).
- Jennings, D.S., Jason, G.A., Hanson, D.J. and Francis, J.P., 1979. Geology Anvil District Map Area. Unpublished Cyprus Anvil Mining Corporation internal company report (1:12 000 scale).
- Piagge, L.C., 2004. Bedrock geology compilation for the Anvil District (parts of 105K/2, 3, 5, 6, 7, and 11), central Yukon. Yukon Geological Survey, Bulletin 15.
- Reed, C.V., 1968. Winter 1968 Moose Lake area diamond drilling program, geology and diamond drilling report. Unpublished Assessment Report #02020, Energy, Mines and Resources, Yukon Government.
- Svensen, P.H., 1967. Geophysical, geological, geochemical & drilling work to February 14, 1967. Unpublished Assessment Report #017933, Energy, Mines and Resources, Yukon Government.
- Terpelmen-Klutt, D.J., 1972. Geology and origin of the Faro, Vangorda, and Swim concordant zinc-lead deposits, central Yukon. Geological Survey of Canada, Bulletin 208, 75 p.
- Woolham, R.W., 1996. Report on a combined helicopter-borne electromagnetic and magnetic survey, Faro, Yukon, NTS 105K/2, 3, 5, 6, 7. Unpublished Aerodart Inc. company report for Anvil Range Mining Corporation (1:24 000 scale).

#### RECOMMENDED CITATION

Piagge, L.C., 2004. Geological map of Swim Lakes (NTS 105K/2 NE), central Yukon (1:25 000 scale). Yukon Geological Survey, Geoscience Map 2004-16, also Plate 16 in Bulletin 15.  
This map accompanies the bulletin.  
Piagge, L.C., 2004. Bedrock geology compilation of the Anvil District (parts of 105K/2, 3, 5, 6, 7, and 11), central Yukon. Yukon Geological Survey, Bulletin 15.  
An earlier version of this map was published as Open File 2000-5 by Exploration and Geological Services Division, Yukon Region, Indian and Northern Affairs Canada.

The legend shown here is for the entire Anvil District (shown in Plate 2 - Geoscience Map 2004-2). Rock units not present in this map area are not coloured in this legend.  
Digital cartography and drafting by L.C. Piagge, Yukon Geological Survey.  
Any revisions or additional geological information known to the user would be welcomed by the Yukon Geological Survey.

Paper copies of this map, the accompanying report and Yukon MINFILE may be purchased from the Geoscience Information and Sales, c/o Whitehorse Mining Recorder, P.O. Box 2702 (R-102), Whitehorse, Yukon, Y1A 2C6. Phone 867-667-5200, Fax 867-667-5150, Email geosales@gov.yk.ca.

A digital PDF (Portable Document Format) file of this map may be downloaded free of charge from the Yukon Geological Survey website at [www.geology.gov.yk.ca](http://www.geology.gov.yk.ca).

Keep this map in a dark area to keep colours from fading.

Yukon Geological Survey  
Energy, Mines and Resources  
Yukon Government

Plate 16  
Geoscience Map 2004-16  
Geological Map of Swim Lakes  
(NTS 105K/2 NE)  
Central Yukon (1:25 000 scale)

compiled by  
L. C. Piagge