



LEGEND

QUATERNARY

HOLOCENE

ORGANIC DEPOSITS - peat and woody material; occurring as a flat to gently sloping plain; overlie lacustrine, till or poorly drained glacioluvial and alluvial deposits but rarely form a dominant geologic unit. Most common in upper Blind Creek valley and as a veneer (<1 m thick) in old growth forests. Permafrost is commonly present within 1 m of the surface. Localized peat development occurs in more poorly drained organic deposits.

O - O - organics - consisting of woody sedge peat, variable thickness. White River ash accumulations are commonly associated with poorly drained peaty areas.

ALLUVIAL DEPOSITS - sand, silt and pebbles with minor cobbles deposited in modern drainages. Common in Blind Creek and its major tributaries.

Ap - alluvial plain - silt, sand and pebbles with minor reworked cobbles and boulders occurring as bars, overbank floodplain or low terrace deposits, 0 - 10 m thick; floodplain subject to periodic floods. Small valley alluvial plains may not be mapped at this scale.
Af - alluvial fan - coarse sand, pebbles, cobbles and mudflow deposits, up to or >10 m thick. Appear as vegetated, often peat covered, landforms developed during post-glacial sedimentation.
Ax - complexes of Ap and Af undivided. Common when a stream is unconfined and also in narrow valleys where side-entry alluvial fans cannot be differentiated from an alluvial plain.

PLEISTOCENE AND HOLOCENE (UNDIVIDED)

COLLUVIAL DEPOSITS - diamicton, gravel, shattered bedrock, and lenses of sand and silt derived from bedrock and surficial sediments by physical and chemical weathering processes. Transport of dislodged debris occurs as surface creep or by mass wasting processes. Permafrost and seasonal freeze-thaw processes often initiate and enhance colluviation. Common on hillslopes north of Blind Creek.

Cv - colluvium veneer - conforms to bedrock topography, <1 m thick.
Ca - colluvium apron - coalescing colluvial fans at the base of a slope, >1 m thick.

Cz - mass wasting - includes slumping, debris slides and rockfalls. Debris slides in till and colluviated till are most common in the map area.

LATE PLEISTOCENE (WISCONSINAN) - MCCONNELL GLACIATION

GLACIOLACUSTRINE DEPOSITS - well-stratified sand, silt and minor clay deposited in lakes impounded by glacial ice; may have a smooth or kelted surface pattern due to melting of buried glacial ice. Sediments form poorly drained areas with peaty blankets. Thermokarsting is common. Small deposit on western border of the map area.

Lb - glaciolacustrine blanket - 1 - 30 m thick.

GLACIOLUVIAL DEPOSITS - stratified to massive; poorly to well sorted; gravel and sand with minor silt and cobbles; deposited by meltwater originating from glacial ice. Common in Blind Creek, the unnamed tributary to Blind Creek from the northeast and the unnamed valley in the north part of the map area that drains into Top River.

Gp - glacioluvial plain - 3-10 m thick.
Gt - glacioluvial terrace - <10 m thick.
Gx - glacioluvial complex - 1 - 30 m thick, composed of deposits of outwash, glaciolacustrine and minor till deposited in an ice contact environment. Hummocky topography is associated with this depositional setting. Crevasse fillings were mapped in an unnamed valley in the north part of the map area.

GLACIAL DEPOSITS (Ill) - unsorted clay, silt, sand, pebbles and cobbles with minor boulders; deposited by or from glacial ice and occurs as subglacial veneer and blanket deposits. Till is common south of Blind Creek and in the north part of the map area. Colluviated till veneers are more common in the middle part of the map area where there is more relief.

Tv - till veneer - conforms to underlying topography, <1 m thick.

Tb - till blanket - gently to moderately sloping plain controlled by bedrock or underlying surficial deposits, >1 m thick.

Tx - till complex - till blanket or veneer composed of meltout till and minor ice contact glacioluvial deposits.

LOWER CAMBRIAN TO CRETACEOUS

BEDROCK - The map area is underlain by metasedimentary and metavolcanic rocks of North American affinity and intrusives of the Anvil Plutonic Suite. North American rocks underlie most of the map area and include the Lower Cambrian Mt. Mye formation, the Cambrian to Lower Ordovician Vangorda formation, the Lower Ordovician Masego Creek formation and undivided Ordovician to Devonian rocks of Root River and Earn Group affinity. Late- and post-metamorphic mid-Cretaceous intrusives of the Anvil Plutonic Suite cut the metamorphic stratigraphy along the west side of the map (Jennings and Jilson, 1986).

R - bedrock - common on plateau summits and ridges north of Blind Creek and in the northwest corner of the map along valley sides.

COMBINED MAP UNITS

The surficial geology unit(s) are shown first followed by the terrain modifiers. Combined surficial geology units are used where, for reasons of scale, two or more deposits cannot be delineated individually. The dominant unit (>50 % of polygon coverage) is shown first and the subordinate units (<50 % of polygon coverage) are shown second and third. A dot separates the surficial units and a dash separates the terrain modifier from the surficial geology.

TERRAIN MODIFIERS

SUB-ARCTIC, ALPINE AND PERIGLACIAL PROCESSES

Pf - permafrost within 1 m of surface
K - normal k
S - solifluction

FLUVIAL PROCESSES

Active - recently active part of floodplain

EROSIONAL PROCESSES

G - gullying - areas of rapid erosion

SYMBOLS

Geological boundary (defined, assumed)	
Glacial meltwater channel	
Moraine ridge	
Esker	
Cirque	
Mass movement failure - slow to moderate (i.e. creeps or slumps)	
Mass movement failure - moderate to fast (i.e. slides or avalanches)	
Crevasse filling	
Aligned landform	
Till geochemistry sample (ppm)	
Drill hole locations (lengths in meters)	

REFERENCES

JENNINGS, D.S. and JILSON, G.A., 1986. Geology and subsurface geology of Anvil Range, Yukon. In: Mineral Deposits of Northern Cordillera, Proceedings of the Mineral Deposits of Northern Cordillera Symposium, MORIN, J.A. (Ed.), Canadian Institute of Mining and Metallurgy, Special Volume 37, p. 319-361.

RECOMMENDED CITATION

BOND, J.D., 1999. Surficial geology map and till geochemistry of Blind Creek (105K/7 SW), central Yukon (1:25,000 scale). Exploration and Geological Services Division, Yukon Region, Indian and Northern Affairs Canada, Open File 1999-6.

Digital cartography and drafting by P.S. Lipovsky, Yukon Geology Program.

Any revisions or additional geological information known to the user would be welcomed by the Yukon Geology Program.

Copies of this map may be purchased from Geoscience Information and Sales, c/o the Whitehorse Mining Recorder, Indian and Northern Affairs Canada, Room 102-300 Main St., Whitehorse, Yukon, Y1A 2B5, Ph 867-667-3266 Fax 867-667-3267.

Keep this map stored in a dark area to prevent map colours from fading.

This map was released Nov. 1999, and was revised Dec. 13, 1999.

Indian and Northern Affairs Canada
Exploration and Geological Services Division
Yukon Region

Open File 1999-6

SURFICIAL GEOLOGY MAP AND TILL GEOCHEMISTRY
OF BLIND CREEK (105K/7 SW), CENTRAL YUKON
by

J.D. Bond
Yukon Geology Program
Geoscience Office

SURFICIAL GEOLOGY MAP AND TILL GEOCHEMISTRY OF BLIND CREEK (105K/7 SW), CENTRAL YUKON SCALE 1:25 000

CONTOUR INTERVAL 100 FEET

Elevations in Feet above Mean Sea Level
North American Datum 1983
Transverse Mercator Projection

1000 0 1000 2000 3000 4000 Meters

Use diagram only to obtain numerical values
APPROXIMATE MEAN DECLINATION 1917
FOR CENTRE OF 105K MAP
Annual change decreasing 4.7

Topographic base produced by
SURVEYS AND MAPPING BRANCH,
DEPARTMENT OF ENERGY, MINES
AND RESOURCES.

Copyright by Her Majesty the Queen
in Right of Canada

ONE THOUSAND METRE
Universal Transverse Mercator Grid
ZONE 8