

Yukon Regional Mineral Potential by Deposit Models

Introduction

The data presented in these mineral potential maps are the results from four separate regional mineral potential assessments initiated by the Yukon Government from 1999 to 2001. The assessments were designed to assist in land use planning exercises, but also may be of interest to the mineral exploration industry. The 18 maps are in PDF format, and each illustrates the mineral potential of a different deposit model, as indicated by the file names. An index tract map and table contain information on the number and type of mineral deposit models that were assessed for each tract. This document provides detailed information on the purpose, methodology and limitations of the mineral assessment process.

Regional Mineral Potential Assessments

Regional mineral potential studies have been completed over the majority of Yukon Territory (with the exception of the northernmost Yukon and southwest of the Alaska Highway). Regional mineral potential was assessed in four phases (Fig. 1).




Figure 1: Regional mineral potential assessments

These regional mineral resource assessments were conducted using a quantitative method for prediction of undiscovered deposits that was developed by the United States Geological Survey (USGS). This method is based on 39 mineral deposit types (i.e., mineral deposit models of Cox and Singer, 1986) and their probability of being hosted in a particular geological environment. The British Columbia Geological Survey (BCGS) modified the deposit models defined by the USGS and added others to best fit the geological and metallogenic setting of the southern Canadian Cordillera (Lefebure and Ray, 1995; Lefebure and Hoy, 1996). For the Yukon assessments, the deposit models utilized by the BCGS were further modified to incorporate Yukon deposits (Fonseca and Abbott, in press). This method is best suited for regions such as Yukon where vast tracts of land commonly lack complete geological characterization and may contain a variety of mineralization styles. Although this method of mineral assessment is not without shortcomings, it yields reproducible and unbiased results.

Mineral potential

The mineral potential of a region describes the probability for the existence of undiscovered metallic mineral deposits. This mineral potential is based on the current state of scientific knowledge, and its accuracy is dependent upon the availability and quality of geoscientific data (also supplemented by the mineral exploration history records). Regional mineral resource assessments utilize the following geoscience and mineral exploration data: (1) bedrock geology maps at 1:250000 and 1:50000 scale; (2) regional airborne geophysical surveys; (3) regional stream sediment, lake sediment, and till surveys (RGS); and (4) exploration history (Deklerk, 2002). These regional assessments were based on existing, publicly available data. Mineral potential of a region is a “snapshot in time” and should be re-evaluated when there is a significant advance in the knowledge of the geology and the mineral deposit types in the region, or when new base data (e.g., RGS data) becomes available.

Assessment Methodology

Each mineral resource assessment consists of seven phases: (1) compilation; (2) definition of tracts; (3) preparation of deposit models; (4) assessment workshop; (5) data entry; (6) statistical simulation, and (7) ranking.

Compilation

Yukon Digital Geology (Gordey and Makepeace, 1999) was used as the geological base map at 1:250000 scale. The overall accuracy of this compilation on a regional scale is considered to be very good, although the geology in some areas is based on studies done as long as 60 years ago. The Yukon Digital Geology compilation includes many recent 1:50000-scale maps produced by the

Yukon Geological Survey (YGS), and 1:250000-scale maps produced by the Geological Survey of Canada (GSC).

Regional stream sediment geochemical surveys (RGS) have been completed over a large part of the Yukon Territory. Median values were calculated for 21 diagnostic elements, and multiples of the medians were reported on 1:250000-scale geochemical maps for each element. At the time of the mineral assessments, geochemical coverage was absent or incomplete in the following 1:250000-scale map sheets: NTS 95C and 95E in southeast Yukon; NTS 106B, 106C, 106E, 106F, and 106L in northeast Yukon; and NTS 116F, 116G, 116H, 116I, 116J, 116K, 116N, 116O and 116P in north Yukon. RGS coverage has improved considerably since the completion of the regional mineral assessments, especially in the north Yukon (Fig. 2).




Figure 2: Yukon RGS coverage

Aeromagnetic coverage is available for most of the Yukon (Fig. 3). There is little or no geophysical coverage for NTS 106C, 106D, 106E and 106F in northeast Yukon. Most flight lines in the southern Yukon are at 0.8-km spacing. Flight lines in the north Yukon (north of $\sim 65^\circ$) are at 2-km spacing. Digital data was captured by digitizing contoured analog data, because most surveys are 1950-1960 vintage. Coloured maps illustrating the variations in the aeromagnetic total residual field were provided for each of the assessments.

Mineral occurrences from the Yukon MINFILE database (anomalies, showings and deposits) were plotted on geological and geochemical maps to highlight

areas of known mineralization and past exploration activity. Summaries and original descriptions of the mineral occurrences in each assessment area, which include deposit type, status, commodities, work history, and geological description, were provided as supplements to the geology and geochemistry maps.




Figure 3: Yukon airborne geophysical coverage

Tracts

The Yukon Territory was divided into four large regions (each corresponding to a distinct mineral assessment phase) based on the large scale geological environment (e.g., Selwyn Basin). The area of each assessment phase was separated into a large number of tracts of approximately equal area (~1000 km²). Tracts were defined on the basis of the regional geology. Tract boundaries are most commonly geological contacts (more specifically faults, lithologic contacts, or limits of Quaternary cover). A few tracts were assigned arbitrary boundaries, such as drainage patterns or roads, in order to maintain similar areas.

Digital deposit models

Tonnage and grade curves for 44 metallic mineral deposit types were utilized for the regional assessments. The number of tracts for each of the four assessments, and the deposit models assessed for each tract are summarized as a separate table, which is included with this document.

Assessment workshops

Assessment workshops took place following the data compilation for each of the four phases. Five industry geologists (hereafter referred to as “the estimators”) with considerable field experience and knowledge of the geology and mineral deposit models applicable to each region participated in the assessment workshops. The following procedure was used for each of the four assessments: (1) for each tract, the estimators decided on the mineral deposit models that could potentially occur; (2) for each mineral deposit model, and for each individual tract, the estimators evaluated the percent probability (from 100 to 0) of discovering new deposits of that type in that tract; (3) for each tract, the estimators recorded their confidence (from 100 to 0) in the current knowledge of the geology; and, (4) for each mineral deposit model, and for each tract, each estimator distributed 100 points between the other four estimators to evaluate the knowledge and experience of each individual estimator. No estimates were made for non-metallic minerals such as diamonds, asbestos, emeralds, and rhodonite. Likewise, potential for placer gold deposits and gravel deposits was not evaluated.

Statistical simulation and ranking

Data provided by the estimators were entered into a spreadsheet. Measurements of tract confidence and confidence level for undiscovered deposits were digitized in AutoCAD, and then copied to the spreadsheet. The data were then converted to a single evaluation for each tract/deposit model combination. The Monte Carlo Mark 3b simulator used the data to produce metal tonnages at the 90%, 50%, 10%, 5% and 1% confidence level intervals for each tract. The tonnages represent a combination of all possible mineral deposit models that could potentially occur within a given tract. These tonnages are then converted to dollar values using 10-year average prices for each of the commodities that are dictated by the relevant mineral deposit models. A “confidence index” is derived from each of these dollar values by dividing the dollar value that corresponds to each confidence interval by the tract area. A “confidence score” is calculated for each of the confidence level intervals by sorting and ranking the confidence index for each tract (i.e., the lowest confidence index has a score of 1, and the highest has a score equal to the total number of tracts). A final confidence score referred to as “sum score” is then calculated for each tract using the individual confidence scores weighted according to the 90%, 50%, 10%, 5% and 1% confidence level

intervals. The sum score value is then ranked from highest to lowest, and defines the rank intervals used on the mineral potential map.

For this compilation, the data provided by the estimators from all four regional assessments were used to calculate, in the same manner as described above, the potential for each tract to host a particular deposit type (i.e., a new “sum score” was calculated for every tract that was assessed for a given deposit model). This value is used to rank the relative potential for each deposit type throughout the Yukon.

Mineral potential maps by deposit models

The mineral potential of the entire Yukon is ranked on the following maps using 18 individual deposit models. Of the 44 deposit models utilized in the 4 regional assessments, these 18 deposit types were deemed the most beneficial for publication as mineral potential maps. Relative rankings are from higher to lower and are illustrated using three categories for purposes of simplicity and ease of display. The index tract map shows the regions covered by each assessment phase, and the numbered tracts within each region. The subsequent maps show the relative potential, from higher to lower, for each tract to contain a specific deposit type. Every tract that was assessed for a given deposit model is ranked, and therefore tracts defined during different assessment phases are now ranked relative to one another. Tracts that were not assessed for a given deposit model are not ranked, and are displayed as white tracts on the respective mineral deposit model map. It should be emphasized, however, that no tract has zero potential and it still may be possible for a mineral deposit of a specific type to exist within a tract not assessed for that deposit model.

Limitations of Regional Mineral Assessments

The primary limitation of mineral potential studies is that they are based on geological knowledge and data that was available at the time of the assessments. Rankings are subject to change as more data becomes available and geological knowledge improves. Although the estimators recorded their confidence in the current knowledge of the geology for each tract, it was not possible to integrate this information into the simulator. Furthermore, there may be potential in Yukon for deposit models that have not yet been recognized. Most commonly, tracts with limited baseline data were ranked as lower potential. For example, many tracts in the North Yukon were either not assessed or were found to have lower potential for most mineral deposit types. This is, at least partly, because of the relatively low level of geological knowledge and lack of baseline data (e.g., RGS) at the time of the North Yukon assessment.

Mineral potential assessments are also limited by the quality of the data on which they were based. For example, RGS data collected in 1976 does provide important information, but has not benefited from recent advances in the science

of geochemistry and may prove to be unreliable for certain elements due to improvements in our understanding in how to collect and analyze samples. The number, locations, and types of mineral occurrences (from the Yukon MINFILE database), although controlled primarily by geology, also depend on the amount of exploration work done, which in turn depends on ease of access, price of commodities, and other non-scientific issues. Also, information pertaining to geology and mineral deposit models from the MINFILE database may require updating, particularly where derived from properties not recently worked.

Despite the limitations, quantitative regional mineral assessments yield reproducible and unbiased results. The deficiencies are a direct consequence of the fact that the mineral potential of a region is a “snapshot in time” and should be re-evaluated when there is a significant advance in the knowledge of the geology and the mineral deposit types in the region, or when new base data (e.g. RGS data) becomes available.

Acknowledgements

This compilation is the result of the extensive efforts of former Yukon Government mineral assessment geologists: in particular A. Fonseca, with large contributions from D. Héon, R. Hulstein and R. Stroshein. Geologists M. Baknes, E. Balon, G. Bidwell, M. Burke, R. Carne, A. Doherty, G. Gilson, M. Jones, P. Holbek, B. Mawer, R. Stroshein, Paul MacRobbie, Jean Pautler, and M. Stammers are thanked for sharing their considerable knowledge of Yukon geology, mineral exploration history, and deposit model profiles during the assessment workshops.

References

- Cox, D.P. and Singer, D.A. (eds.), 1986. Mineral Deposit Models. U.S. Geological Survey, Bulletin 1693, 379 p.
- Deklerk, R., 2002 (comp.) Yukon MINFILE 2002. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada.
- Fonseca, A. and Abbott, G. (comps.), 2003 (in press). Yukon Mineral Deposit Models. Yukon Geological Survey, Open File.
- Gordey, S.P. and Makepeace, A.J. (comps.), 1999. Yukon Digital Geology. Geological Survey of Canada, Open File D3826 and Exploration and Geological Services Division, Yukon Region, Indian and Northern Affairs Canada, Open File 1999-1(D).
- Lefebvre, D.V. and Ray, G.E. (eds.), 1995. Selected British Columbia Mineral Deposit Profiles, Volume I – Metallics and Coal. British Columbia Ministry of Energy, Mines, and Petroleum Resources, Open File 1995-20, 136 p.

Lefebure, D.V. and Höy, T. (eds.), 1996. Selected British Columbia Mineral Deposit Profiles, Volume 2 – Metallic Deposits. British Columbia Ministry of Energy, Mines, and Petroleum Resources, Open File 1996-13, 172 p.