



1:250 000-scale topographic base data produced by CENTRE FOR TOPOGRAPHIC INFORMATION, NATURAL RESOURCES CANADA

ONE THOUSAND METRE GRID Universal Transverse Mercator Projection North American Datum 1983 Zone 8 CONTOUR INTERVAL 100 FEET Elevations in metres above Mean Sea Level

W skarn/porphyry Weighted sums model (Principal Component Residuals) Sheet 12 of 13



Use diagram only to obtain numerical values APPROXIMATE MEAN DECLINATION 2015 FOR CENTRE OF MAP

Table with 3 columns: 11SI (CARMACK), 105L (GLENLYON), 105K (TAY RIVER), 115H (AIBIK LAKE), 105E (THIS MAP), 105F (QUIET LAKE), 105G (DEADASH RANGE), 105D (WHITEHORSE), 105C (TESLIN)

INTRODUCTION

New geochemical data from re-analysis of archived stream sediment samples have been assessed using weighted sums modeling and catchment basin analysis, as described in the methodology report that accompanies this map (Mackie et al., 2015).

sediments. One uses data levelled by the dominant geology mapped within each catchment. The other uses residuals calculated from regression against principal components. Weighted sums models (WSM) have been generated using the processed data. Importance rankings used in the WSM for a variety of deposit types are summarized in Table 2. Each model is optimized for a specific deposit type however multiple deposit types may be represented in a given model due to similarities in elemental abundances and associations.

SAMPLING AND ANALYSIS PROGRAMS

Regional stream sediment and water samples from the Lake Laberge map area (NTS 105E) were collected at a reconnaissance scale in 1988 as part of the National Geochemical Reconnaissance program under the Canada-Yukon Minerals Development Agreement (Hornbrook & Friske, 1989).

Exploratory data analysis of both raw element data and principal components shows that the distribution of many commodity and pathfinder elements is related to lithological variation. For example, the first principal component, accounting for ~27% of the total variation, shows high positive loadings for Sr, Ca, LOI, Se, Hg and Cu, and high negative loadings for Ce, Th, La, Li, Rb and Pb.

MINERAL OCCURRENCES

A variety of types of base and precious-metal mineralization have been documented in the map area as summarized in Table 1 (Yukon MINFILE, 2015). The most notable occurrences are classed as Cu-Ag-Pb-Zn skarn (Laberge prospect, Dycer and D'Abbadie showings), Polymetallic Ag-Pb-Zn-Au vein (Loon Prospect, RK, Deet, Livingston and Sylvia showings) and Cu-Mo porphyry (TUV Prospect).

The effectiveness of historical sampling coverage has been assessed empirically using graphs of WSMs plotted against catchment surface area to determine the ideal maximum catchment size (12 km<sup>2</sup>).

WEIGHTED SUMS MODELING

As described in the report accompanying this map (Mackie et al., 2015), two approaches have been used to subdue the influence of background lithological variation and secondary absorption on the composition of stream

Table 2: Importance rankings for weighted sums models using residuals on principal components.

Table with 3 columns: Target Deposit Type, Other Deposit Types, and 20 elemental columns (Mn, Fe, Co, Ni, Cu, Mo, Zn, Pb, Ag, Au, As, Ba, Cd, Sn, Sb, Te, Hg, Tl, Bi, W). Rows include Polymetallic Ag-Pb-Zn skarn, VMS (Zn-rich), Porphyry Cu-Mo, Intrusion-related Au, Epithermal Au-Ag, and W skarn.

\*Polymetallic Ag-Pb-Zn type includes vein and manto styles; SEDEX = sedimentary exhalative; VMS = volcanic-hosted/associated massive sulphide deposits. Raw data following a log10 transformation.

Table 1: List of Mineral Occurrences for NTS map sheet 105E (Yukon MINFILE, 2015)

Table with 5 columns: Number, Name, Type, Status, and Commodities. Lists various mineral occurrences such as Livingsston, TUV, Loon, Laberge, Ruth, Packers, Walsh, Semenov, Illusion, Cassier Bar, Sylvia, Corduroy, Lore, Mustard, Bacon, Klusha, Salmon, Hitckens, Ovas, Enof, Lake, Germ, Preston, Rank, Deet, Milner, Braburn Lime, Egypt, Richthofen, Reef, Slne, Rk, Nickeline, Dycer, Trenchice, Cros, Napua, Brenda, Little Bear, Mendozna, Teraktu, Fone, Debicki, Little Violet, Coughlan, and Aurier.

RECOMMENDED CITATION

MACKIE, R., ARNE, D. AND PENNIMPEDE, C., 2016. Weighted sums model for W skarn/porphyry deposits using principal component residuals. In: Enhanced interpretation of stream sediment geochemical data for NTS 105E. Yukon Geological Survey, Open File 2016-9, scale 1:250 000, sheet 12 of 13.

Catchment basin polygons generated by the Yukon Geological Survey (J. O. Bruce).

Any revisions or additional geological information known to the user would be welcomed by the Yukon Geological Survey.

Paper copies of this map and the accompanying report may be obtained from the Yukon Geological Survey, Energy, Mines and Resources, Government of Yukon, Room 102-300 Main St., Whitehorse, Yukon, Y1A 2B5. Ph. 867-667-3201, Email geology@gov.yk.ca.

A digital PDF (Portable Document File) file of this map may be downloaded free of charge from the Yukon Geological Survey website: http://www.geology.gov.yk.ca.

Yukon Geological Survey Energy, Mines and Resources Government of Yukon

Open File 2016-9

Weighted sums model for W skarn/porphyry deposits using principal component residuals (NTS 105E) Sheet 12 of 13

by

Rob Mackie, Dennis Arne, and Chris Pennimpe

REFERENCES

- Hornbrook, E.H. and Friske, P.W., 1989. National Geochemical Reconnaissance stream sediment and water geochemical data, southern central Yukon (105E). Geological Survey of Canada, Open File 1960.
Jackman, W., 2015. Regional stream sediment geochemical data, Lake Laberge area, southern Yukon (NTS 105E). Yukon Geological Survey, Open File 2015-7.
Mackie, R., Arne, D. and Brown, O., 2015. Enhanced interpretation of regional stream sediment geochemical data from Yukon: catchment basin analysis and weighted sums modeling. Yukon Geological Survey, Open File 2015-10.
Yukon MINFILE, 2010. Yukon MINFILE - A database of mineral occurrences. Yukon Geological Survey, www.data.geology.gov.yk.ca, accessed May 2015.